April 4th, 1920: Getting rapidly worse; orthopnoea; oedema of lungs; cannot lie down at all, and has to be nursed in a chair. On April 12th I persuaded Dr. Whyte, a resident, to administer ether to the patient in the upright position. As soon as the patient was "under" he was placed on his face over an air-bag, in which position he was able to breathe sufficiently well for me to proceed with the operation. I stripped both kidneys in twenty-four minutes from the first cut to the last stitch. There was advanced oedema of the perirenal fat. The capsule was thin and friable, and very adherent to the kidney substance, and stripped badly. There was no shock, and the same night the patient was able to sleep quietly lying on his back, propped up by pillows. In a few days the oedema had much diminished and the dyspnoea had disappeared. The improvement was only maintained for a few weeks, and then the symptoms returned again, though the dyspnoea did not become so bad.

He went to the infirmary, where he died of phthisis and uraemia on September 6th, 1920. There was no *post-mortem* examination.

CASE III.

Male, aged 18 years. Admitted to the London Hospital under the care of Dr. Hadley, June, 1920, with a history of six months' general oedema and pallor, and headaches.

On admission, general oedema, pallor. Urine: Albumin $\frac{1}{2}$; casts; polyuria, as much as 90 oz. in twenty-four hours. Systolic blood pressure 170 to 180 mm. of mercury.

Operation July 22nd, 1920. Dr. Whyte administered open ether. I stripped the right kidney in thirteen minutes, the left in nine minutes. Capsule stripped easily. Kidneys enlarged, pale and mottled and fatty. No oedema of perirenal fat. Lymph oozed out of kidney on stripping. Portion removed for section.

The patient improved at once. In October, 1920, he was able to go about and looked upon himself as well. There is, however, still a little albuminuria, and the legs swell very slightly at times.

CASE IV.

Male, aged 20 years. Admitted to the London Hospital July, 1920, under Dr. Hadley, with a history of general oedema since six weeks before admission.

On admission, general oedema, pallor not marked; no vomiting; no fever; no headache. Urine: Solid with albumin; specific gravity 1020-1025; oliguria, 18 to 32 oz. in twenty-four hours. Systolic blood pressure 130 mm. of mercury. No improvement after five weeks' medical treatment.

Operation on July 22nd, 1920. Dr. Probyn Williams administered ether. Each kidney stripped in eleven minutes. Fluid poured out of perirenal fat. Capsule thickened, stripped badly, and tore pieces of kidney out. Marked fibrosis. Kidneys not enlarged; capsule pale because of the oedema of capsule itself; the cortex appeared to show areas of fatty degenerations, but was not diffusely and completely degenerated as in the other cases.

Three weeks later the oedema had disappeared, the urine had increased to normal amount, and the albumin had diminished. The patient was getting up in September, but then got chilled when sitting about the ward, and relapsed.

RECONSTRUCTIVE ANEURYSMORRHAPHY IN THE THIRD PART OF THE AXILLARY ARTERY.

BY

C. JENNINGS MARSHALL, M.S., M.D., F.R.C.S.,

ASSISTANT SURGEON AND ORTHOPAEDIST, KING'S COLLEGE HOSPITAL; SURGEON, MANOR HOUSE HOSPITAL.

A MAN, aged 25, wounded in March, 1918, was, after three months in hospital and *dépôt*, returned to his unit. He was demobilized in 1919, and resumed his occupation as a baker. At that period he had, at most, occasional aching in the axilla. In February, 1920, he noticed a swelling in the armpit which increased slowly in size; it was accompanied by a feeling of heaviness in the left upper limb after use. After a short time he began to experience pain radiating down to the index and middle fingers and to lose power in the index. After six months' delay he was admitted to the Manor House Hospital.

It was seen that the patient had sustained a through-and-through bullet wound of the left axilla. A fine scar above the lower margin of the pectoralis major one inch and a half from its insertion indicated the entry, and a slightly larger one a little outside the inferior angle of the scapula the exit. In the line of the artery was an aneurysm the size of a large walnut, its upper third lying under cover of the anterior axillary fold. The expansion was forcible, but not great in extent; the tumour could not be emptied by pressure owing to the pain referred to the index. There was no venous hum. The adjacent tissues were not thickened—the tumour was quite discrete—very little inflammation or effusion had evidently accom-

panied the wound. There was slightly impaired sensation of the ends of the index and middle fingers; flexion of the index was very weak. The peripheral circulation of the limb did not differ from that on the other side.

Operation.

Open ether was administered by Dr. Bookey, and Mr. A. E. Woodall, F.R.C.S. (resident surgeon), assisted.

The arm was fully abducted, and an incision four inches long was made in the line of the artery, its centre over the aneurysm, and its upper end curving inwards towards the chest wall. The deep fascia was divided to the full extent of the wound, and the axillary artery was exposed one inch above the upper limit of the aneurysm; the minimum amount of dissection was done to permit the application of a Crile's artery clamp at this spot. The vessel was then treated similarly below the sac; the two

clamps lay above the anterior circumflex and the profunda artery respectively. The aneurysm was next defined by sharp dissection. Only at the lower and inner part was there any difficulty; elsewhere the tumour required only a few touches of the knife to free it. When the vein had been separated at the lower and inner part it was seen that the inner head of the median was bound to the aneurysm by dense fibrous tissue; when this was divided, however, the nerve showed no break of continuity. The sac was then defined up to its point of origin from the artery, to which it was moderately adherent up to its neck, and then opened by a longitudinal incision. At once it became apparent that the collaterals needed control. The branches arising between the clamps were quickly defined by blunt dissection, the flow from the sac being checked by pressure, and a clamp placed to include them; very slight pressure secured complete haemostasis. The

opening from the artery was oval ($\frac{1}{2}$ in. by $\frac{1}{4}$ in.); at the junction of vessel and sac there was a ring of almost cartilaginous density, as became evident later on an attempt to pass the needle at this spot. The sac, which was free from clot, had the thickness of blotting paper; its interior looked smooth and shiny everywhere. Two branches had been clamped—the posterior circumflex just above the level of the opening, and the anterior at its lowest level, each at a part of the artery opposite to the opening into the aneurysm. A test was made to determine the relative importance of these in the collateral circulation. With the posterior vessel clamped the flow from the anterior was the feeblest possible trickle. That from the posterior was considerable, steady, and in quantity about equal to that from a radial vein.

It was estimated that to suture the edges of the opening, allowing for a fair depth of apposition of the intima, would diminish the diameter of the vessel, compared with the part above, to about two-thirds, which seemed adequate. The sac was therefore cut away, leaving only a thin rim round the neck. The whole area was kept moist by normal saline. With a non-cutting fine needle a continuous suture was introduced, beginning above and finishing below the opening, passing through the arterial wall immediately adjoining the thick ring; the stitching was mattress in type, every other stitch returned slightly to give a locking effect. The material was very fine silk.

The clamp on the collaterals was then released and fairly free oozing occurred, the fineness of the silk having prevented any but the slightest tension in approximation. The clamp was again tightened and a stronger silk suture was introduced, the needle traversing the artery wall on either side and returning similarly through the rim of sac, thus embracing the hard ring and securing apposition of

its opposite sides. For further safety these stitches were made to cross in pairs. The clamps were released in order, first from the branches, then the distal, then the proximal, and, as the severest test, the distal clamp was tightened again. There was not the slightest oozing. In view of the exposure of the site of suture to strain, a graft of fascia lata was fixed as a collar round the vessel at this point, with its fibres transverse to the line of the artery, and with slits to allow the passage of the circumflexes. Above and below a loose stitch in the fascia near by prevented telescoping. The wound was then closed and dressed with gauze wet with 1 in 2,000 flavine. At the close of the operation the radial pulses were equal.

After-History.

Convalescence was uneventful. There was pain referred to the index and middle fingers for three days. The wound healed by first intention, and physiotherapy appropriate to the nerve condition and to the axillary scar was adopted from the twelfth day.

Three months later there was no sign of yielding; at the site of the suture was a thickening due to the presence of the graft. The brachial pulses were equal and the index showed almost complete recovery. Pulsation, though slight, indicated that thrombosis and obliteration of the artery had not occurred at the point of suture.

FRACTURE OF FEMUR: A SIMPLE METHOD OF EXTENSION.

BY

B. F. PENDRED, M.R.C.S., L.R.C.P.,
LOUGHTON, ESSEX.

FRACTURE of the femur in private practice is always looked upon with apprehension, and even in hospital work with some degree of anxiety.

One great reason for this attitude is the necessity for having a complicated apparatus made up of iron framework, a mast and guy-ropes, all requiring frequent adjustment and refitting, to the pain and distress of the patient. This, perhaps, may not be true of fractures below the middle of the shaft, as in these cases fair results may be obtained by use of foot extension with the limb lying straight along the bed.

The method I am about to advocate is for fractures above the middle, where the upper fragment is flexed forwards. As this fragment is too short to be maintained in any other position than that allowed by the muscles attached to it, it becomes necessary for the lower fragment to be brought into line with the upper, at the same time keeping up extension in that line. In such a case the patient is placed in bed on his back, the head and shoulders being supported by two or three pillows. The knee of the injured limb is raised until the femur is at the required angle, which is generally about 30 to 40 degrees to the horizon. It is maintained in this position with a knee pillow, or better still by a pyramid of sand-bags. A ring-shaped air-cushion is slipped over the foot, drawn up to the knee and inflated as fully as possible, the foot being allowed to fall back on the bed. A strong bandage is tied round the air-cushion exactly opposite the knee and the other end carried to a pole or mast fitted to the foot of the bed; strong traction is made in the line of the upper fragment, lower fragment, diameter of cushion, bandage and point on mast, until the ring becomes definitely oval, when the bandage is fixed to the mast. It will be seen that the extension is procured by a pneumatic pad in the flexure of the knee, and that the strain is transmitted to the femur through the head of the tibia and the ligaments of the knee-joint. These being rigid structures serve the purpose well.

Pneumatic pressure at the point indicated is justifiable, because all the great nerves and blood vessels dip deeply here under the heads of origin of the gastrocnemii, which being massive and fibrous in this situation are well adapted to support continuous pressure and protect underlying structures.

The advantages attendant on this method are:

1. Simple apparatus. A few pillows, a long piece of stick, a bandage and a ring air-cushion or inner tube of a motor tyre. Such things can be obtained almost anywhere at very short

notice, and there is no need to keep apparatus of various sizes in stock.

2. Ease of application. No skilled assistance required.

3. Once fixed adjustment is very seldom required. The bed-clothes are arranged so that the ring-cushion is uncovered and visible, a glance at its shape informs one immediately of the degree of tension and inflation.

4. The foot hanging over the pneumatic pad acts as a kind of rudder and prevents rotation of the lower fragment, or if desired it can be pushed inwards or outwards, rotating the lower fragment in any required direction.

5. There being no pulleys, guy-ropes, weights, or iron frame, there is very little necessity for disturbing the patient, which saves much anxiety and suffering. A bed slipper is easily adjusted.

6. The thigh is supported underneath by sand-bags, and others may be used at the sides if required to give a sense of security, but these are not necessary. The thigh can therefore be fully exposed, permitting easy examination, dressing or massage of suitable type, all without any disturbance of the fragments whatever.

I have treated only one case by this method:

A boy, aged 10 years, sustained a simple fracture of the right femur above the middle of the shaft. He remained in the apparatus for four weeks with the very slightest adjustments. At the end of that time, union being quite strong, the limb was removed from all restraint. At first any attempt to straighten the limb caused pain in the knee-joint, but full flexion and extension returned without assistance within forty-eight hours. The only evidence of pressure by the pneumatic pad was a very superficial blister, which no doubt would have been obviated by binding lint over the rubber before application. The shortening was under a quarter of an inch. At no time was there any numbness, tingling, or oedema of the leg or foot.

The extreme simplicity and efficiency of this method, as compared with many devices advocated, are my excuse for seeking publication.

AN UNUSUAL CASE OF MYELOCELE.

BY

A. J. COKKINIS, M.B., B.S. LOND., F.R.C.S. ENG.,
ILFORD.

In practically all the standard textbooks myelocoele is stated to be incompatible with life, the infant being either stillborn, or dying within a day or two. Bland-Sutton holds a somewhat different view. He states:

"Many children in which this condition is present are still-born: the few that survive their birth rarely live longer than three days, the continual leakage of cerebro-spinal fluid being sufficient to explain the invariable brevity of their lives." (*Tumours: Innocent and Malignant*. 1917.)

He thus admits the possibility of survival beyond the first day or two.

In inquiries, admittedly limited, which I have made on this subject, I have been unable to trace a single case in which the duration of life exceeded a period of three days. I have therefore thought that the following case, in which life was prolonged for twenty-three days, merited publication.

I was called to see the patient, a male infant, an hour after its birth, at the Ilford Maternity Home, during the early hours of September 18th, 1920. The reason for the early call was that the infant seemed very feeble and had difficulty in breathing. On arrival I found a well-nourished, good-sized baby, which soon revived and cried lustily. He had a well-shaped head, with no evidence of hydrocephalus.

The lower limbs were held rigidly flexed to a right angle at the hips, and rigidly and completely extended at the knees. The legs resisted straightening, and were paralysed, wasted, and, as far as could be made out, anaesthetic. There was double talipes equino-varus, the varus element markedly predominating, with a well-marked crease on the soles, corresponding to the transverse tarsal joint.

Looking at the back, there was a large, raw, circular area, somewhat depressed, and corresponding to a deficiency in the neural arches and failure of closure of the neural groove, opposite the lower three lumbar and upper two sacral bodies. The open spinal cord was seen in the middle line at the upper part of the defect, while the nerves of the cauda equina could be distinguished in the lower part. The myelocoele narrowed and deepened as it was traced upwards into the funnel-shaped termination of the central canal. Cerebro-spinal fluid was constantly, though slowly, dripping out of the apical opening, bathing the raw area, and then running in drops over the buttocks. The surrounding skin was rendered as far as possible aseptic, and a dressing applied. The next day the flow of cerebro-spinal fluid was freer, and some congestion developed round the myelocoele. During the first week the child's condition remained good. The raw area gradually contracted down and became deeper, while the inflammatory redness subsided.