bone could be felt high up with a soft blunt probe. An x-ray photograph showed large frontal sinuses divided into loculi with a fairly sharp outline on either side. There was a definite general haze round the glabella, but no indication of an opening in the anterior wall of the left sinus. The antra were clear on transillumination.

On October 18th, 1920, the pulsating swelling was opened by a vertical incision and a large quantity of glairy muco-pus escaped. Through this opening the sinuses were explored and were found to reach 2 in. above the orbital rim, and to the external angular processes. The nasal fossa was easily entered with a blunt probe on the right but not on the left. A Killian incision was made. This disclosed a complete absence of the roof of the orbit and extensive caries with the dura exposed above. The nasal fossa was now easily entered on the left. The carious ethmoidal cells were curetted away and a large drainage orifice made into the nose. Gauze wicks were led through from the upper incision and the inner extremity of the lower incision to the nose, and from the body of the sinus to the outer angle of the wound. In spite of careful drainage and vigorous antisyphilitic treatment it became obvious after some time that it was essential to remove the anterior wall of the left sinus and let the skin fall in. This was done on January 7th, 1921, good drainage was established right and left, and after careful curetting the wound was lightly packed. The patient made an uninterrupted recovery. The deformity was, of course, considerable.

CASE IV.

A. C., Anglo-Indian, aged 51, was admitted on October 22nd, 1920, with proptosis of the right eye, with swelling and oedema of the lid and brow, and acute pain. There was no impairment of vision. The first symptoms he noticed were bulging of the eye and double vision. A month previous to this he got a knock on the brow, and a week before exposed himself by sitting up shooting. Shortly after this severe pain started. When seen at hospital he presented a typical picture of orbital cellulitis. An incision was made in the superior fornix and pus evacuated, but this did not relieve the condition, and a fluctuating swelling appeared over the outer third of the eyebrow. Nasal examination revealed oedema of the middle turbinate and pus coming from the frontal sinus. No x-ray photograph was obtained. On November 13th, 1920, the frontal sinus was opened by the intranasal route. This did not relieve the condition, and a week later the external operation was done. There was caries of the orbital rim and a large pocket of pus under the periosteum of the orbital roof, which was perforated and opened into the frontal sinus. The latter was small and communicated with the superior meatus with a very angular fronto-nasal duct. Free drainage was established into the middle meatus, and gauze drains led through. About a month later a large sequestrum was removed from the outer part of the orbital rim. The patient frequently complained of severe neuralgic pain. In spite of a negative history of syphilis and a negative Wassermann reaction he was put on mercury and iodides. He improved steadily, but still complained of supraorbital pain. He was discharged at his own request, without diplopia, and with diminished proptosis, but there was considerable interference with the movement of the upper lid due to adhesions to the periosteum at the orbital rim.

CASE V.

This was a case of very extensive ulceration of the orbit and frontal sinuses, due to gummatous disease, superimposed upon which was myiasis. This case has already been fully recorded, as it was of interest also from an entomological point of view, the adult flies having been hatched out from the maggots.¹

CASE VI.

M. R., European, aged 42, was admitted on December 31st, 1920, complaining of acute supraorbital pain on the left side, which had prevented him from sleeping for some time. He had very slight swelling of the lid and thickening of the supra-orbital nerve. There was pain on pressure against the orbital rim and the nerve was excessively tender. The vision was 6/5 for both eyes and the fundi were normal. There was a negative history of syphilis, no clinical signs, but a positive Wassermann reaction. An x-ray photograph was not obtained. Transillumination was negative. He gave a history of a very severe fall from a motor bicycle five months before, with temporary unconsciousness and bleeding from the ear. The exact location of the latter was somewhat uncertain, and he had a perfectly normal drum. There was no history of influenza or acute catarrh. The nasal fossa looked healthy, but on passing a cannula into the fronto-nasal duct a small quantity of muco-pus was evacuated. He was given an injection of novocain, followed by alcohol, into the supraorbital nerve, which relieved his pain. On December 31st, 1920, the anterior end of the middle turbinate was removed with scissors and snare and the ostium of the frontal sinus enlarged. The anterior ethmoidal cells were also broken down. He developed a third nerve paralysis three days after operation. He was put on injections of mercuric cream and novarsenobillon on alternate weeks and rapidly got better. On January 21st, 1921, the paralysis had almost gone and there was no diplopia. He had no pain and his nasal fossa and sinus were apparently healthy.

REFERENCE.

¹ Indian Medical Gazette, February, 1921, p. 58.

THE ISOLATION OF DIPHThERIA BACILLI.

BY

DENYS R. WOOD, F.I.C.,

PUBLIC ANALYST AND BACTERIOLOGIST, SOMERSET COUNTY COUNCIL.

It is sometimes necessary to obtain the diphtheria bacillus in pure culture in order to test its pathogenicity, as, for instance, when found in swabs taken from lesions other than the throat and nose, such as the ear, the eye, and sores on other parts of the body, and occasionally from the throat and nose also in the case of very persistent carriers.

A useful medium for this purpose is described in Browning's *Applied Bacteriology* as follows:

Sheep serum, sterilized by heating on several occasions for one hour at 57° C.	5 c.cm.
1 per cent. telluric acid in distilled water	0.9 c.cm.
Peptone water agar (roughly neutral to litmus) to ...	100 c.cm.
The serum and telluric acid solution are added to the melted agar at 50° to 55° C. before pouring into Petri dishes.	

In my experience sheep serum, when the blood is obtained from the abattoir, is seldom sterile when heated at 57° C. for one hour on seven or eight consecutive days, and the ether and chloroform methods of sterilization are equally unreliable, in fact, the only reliable method appears to be filtration through a Chamberland filter. It is therefore a great advantage to be able to dispense with this constituent.

Diphtheria bacilli grow exceedingly well on Gordon and Hine's legumin tryptagar,¹ and when this medium is used no sterile serum is necessary. The addition of 0.3 c.cm. of sterile 1 per cent. telluric acid to 10 c.cm. of this agar gives a very satisfactory medium. Plates of this medium are poured, and when set brushed with a dilute emulsion of the growth containing the diphtheria bacilli and incubated at 37° C. for forty-eight hours, by which time the characteristic appearance of the colonies of the different organisms is most distinct.

The majority of organisms other than the diphtheroids reduce the telluric acid and produce blackish colonies; this is very marked with the staphylococci. The staphylococci do not reduce the telluric acid, their colonies are characteristically greyish-white, small (usually less than 1 mm. diameter), very flat and very translucent. The colonies of the diphtheroids are greyish-white, semi-translucent, slightly convex, about 1 mm. in diameter, with a slightly darkened central spot. By the use of this medium I have selected as many as five colonies of *B. diphtheriae* in succession from a plate containing numerous colonies of other organisms.

The morphology of the diphtheria bacillus when grown on legumin tryptagar differs from its morphology when grown on Loeffler's serum; the bacillus stains more uniformly, the characteristic beads are not evident, and club-shaped forms are common. With a little practice, however, it can readily be recognized and distinguished from Hoffmann's bacillus grown on the same medium. It is unfortunately not possible to distinguish between the colonies of Hoffmann's bacillus and those of true diphtheria bacilli. Douglas's trypsin broth² with the addition of pea-flour extract and 0.5 per cent. Witte's peptone (to help toxin production), is very suitable for the cultivation of *B. diphtheriae* for animal inoculation. Almost invariably a good growth is obtained—an essential condition of the experiment—while with ordinary broth this is far from being the case, the growth more often being only scanty. Glucose litmus legumin trypsin broth is very suitable for the observation of acid productions from this sugar.

REFERENCE.

¹ Vide BRITISH MEDICAL JOURNAL, November 18th, 1916; also Reports of the Medical Research Council, Special Report Series No. 19. ² Loc. cit.

A SMALL Italian surgical hospital with an out-patient department has been opened at Aleppo by the National Italian Association.

THE sixteenth Congress of the German Orthopaedic Society will be held in Berlin from May 18th to 20th, when the principal subjects for discussion will be bone and joint tuberculosis and rickets.