and CO₂ output of a normal individual living a normal life vary together. This, we suggest, is due to the influence of food, and that if this be excluded the CO₂ cost per hour of this constant work remains the same at any period.³ Benedict and Cathcart's well-known experiments on a trained bicyclist in the post-absorptive condition, show clearly that severe constant work requires the same amount of CO₂ exhalation per second at the end of three hours of work as during the first hour.⁴ Using a bicycle ergometer and four-minute samples of expired air, we have obtained similar results (Table II). It is only after taking food (Tables III and IV) that our subjects show a rise of CO₂ exhalation similar to the increase shown in Waller and De Decker's charts.

TABLE III.—Showing Effect of Food whilst Resting.
(Subject C.P., resting. Ten-minute samples of expired air.)

Date.		Heat Loss.	CO ₂ (c.cm. per sec.).
April 20, 1921	One hour after food ...	1.13	3.2
March 30 ..	" " " " ...	1.16	3.0
April 20, 1921	Two hours after food ...	1.43	3.7
March 30 ..	" " " " ...	1.47	4.0

TABLE IV.—Showing Effect of Food whilst at Work.
(Subjects C.P. and L.H., working on ergometer. Four-minute samples of expired air.)

	Heat Loss.			CO ₂ (c.cm. per sec.).		
	L.H.	C.P.	C.P.	L.H.	C.P.	C.P.
Two hours before food	3.3	4.9	4.23	12.7	12.3	11.8
One hour before food	3.5	4.8	3.88	11.9	12.0	11.1
Food taken.						
One hour after food ...	3.9	4.2	4.9	13.1	11.5	12.6
Two hours after food	4.2	5.1	4.86	14.9	14.3	14.7

It is well known that a fatigued or uncomfortable subject carries out a given amount of work at a greater cost than when normal, but it does not seem reasonable that the colliers and shoemakers under consideration are so unskilled and of such poor calibre as to show signs of fatigue after a couple of hours' constant work, when this work is of the degree specified.

REFERENCES.

¹ *Journal of the Royal Army Medical Corps*, November, 1918.
² Publication No. 261, Carnegie Institute, Washington, 1918. ³ Lusk, *Science of Nutrition*, Third Edition, p. iii. ⁴ *Muscular Work*, p. 66. The Carnegie Institute, Washington, 1913.

COMPARATIVE SYSTOLIC BLOOD-PRESSURE READINGS IN THE ARM AND LEG IN AORTIC INCOMPETENCE.

BY

O. K. WILLIAMSON, M.D.CANTAB., F.R.C.P.LOND.,

PROFESSOR OF MEDICINE ELECT, UNIVERSITY COLLEGE, JOHANNESBURG;
PHYSICIAN TO THE CITY OF LONDON HOSPITAL FOR DISEASES
OF THE CHEST; LATE ASSISTANT PHYSICIAN TO THE
WESTMINSTER HOSPITAL.

THE higher systolic reading by the blood-pressure instrument in the leg compared to that in the arm, when both are at the level of the heart, appears to have been so generally accepted as a characteristic and even diagnostic sign in aortic incompetence that it would seem to serve a useful purpose briefly to review the subject, so as to consider whether or not this conclusion is a sound one, and, if not, what are the fallacies which underlie it.

Leonard Hill, with the co-operation of Martin Flack and W. Holtzmann,¹ first made the observation that in all cases of aortic regurgitation there is a marked difference between the reading of the systolic arterial pressure in the arm and in the leg, that in the leg being the higher, the observation being taken in recumbency. These observers further state that "when such patients are lying quiet in bed this difference is a diagnostic sign of aortic regurgitation." They found that the difference of pressure is not abolished by the method of oscillating the pressure up and down near the obliteration pressure. The explanation which they give of the difference is that the "big systolic waves are conducted better

down the leg arteries, these being slightly thicker and more contracted and rigid." In a later paper Hill and Rowlands² brought forward additional cases illustrating the same phenomenon. H. A. Hare came to similar conclusions to those of Hill.³ H. D. Rolleston⁴ confirmed the observation, and among other conclusions found that the difference between the maximum systolic blood pressure in the arm and leg is most marked in uncomplicated and compensated aortic regurgitation, but "is much less marked in aortic regurgitation when the compensation is failing, and in cases of combined aortic regurgitation and mitral disease"; further, that "in recent aortic regurgitation before compensation is established the difference is slight or may be absent." J. R. Murray⁵ published records showing similar differences. J. Heitz⁶ recorded observations on 51 cases concerning this phenomenon, and whilst finding it in the majority of cases, disagrees with Hill as to its constancy. He, moreover, draws attention to the important fact that the difference is to be found in other classes of cases. He says: "L'élévation de pression dans les artères tibiales peut être constatée chez la plupart des sujets atteints d'insuffisance aortique" . . . but "elle peut manquer dans un certain nombre de cas." Again: "L'élévation de la pression systolique aux membres inférieurs peut se constater avec une certaine fréquence chez des sujets hypertendus et non atteints d'insuffisance aortique." He has also found it in abdominal aortitis. He concludes that the sign "ne présente donc pas de valeur diagnostique absolue."

Such a difference of pressure readings may certainly occur in other conditions than aortic incompetence. Thus I have observed it in high-pressure cases associated with arterio-sclerosis,⁷ and, according to Clifford Allbutt, this observation was first made by Volkmann.⁸ Further, MacWilliam has found the same difference in some patients with sclerotic arteries.

The conclusions drawn from my own observations are as follows:

1. Although higher systolic readings in the leg than in the arm appear to occur in the majority of cases of aortic incompetence, there is a large minority in which they do not obtain.

2. There is no evidence that such cases, when they do exist, are the result of the actual aortic incompetence rather than of the commonly associated conditions of hypertonic contraction or hypertrophy of the muscular coat, with or without accompanying sclerotic changes, conditions which also occur independently of aortic incompetence.

3. In support of the arterial explanation of such differences is the fact that the large majority of cases of the series in which the arteries were thickened showed the difference, being a much larger proportion than obtained in the total number of cases of the series (all of which had aortic incompetence); and further, that in the adult cases of aortic incompetence, nearly all of which had thick arteries, the phenomenon was more than twice as common as in the children, in none of whom were the arteries thickened.

4. If due to the aortic incompetence it would seem reasonable to suppose that the difference of reading should show some relationship to the degree of incompetence, which for practical purposes may be considered as measured by the pulse pressure. No parallelism, however, exists between these arm-leg differences and the corresponding degrees of pulse pressure.

5. In the few cases in which this was tried, in the majority the difference of pressure was abolished by repeated recompression, a fact decidedly in favour of the arterial explanation. (In cases of a higher leg reading than that in the arm in arterial disease it has been frequently found that the difference is abolished by such recompression.)

Method of Observation and Apparatus.

This paper being based on observations which were spread over many years, different instruments were used, but these were one or other of the modifications of the Riva-Rocci method—namely, Stanton's instrument, Oliver's compressed air haemomanometer, Martin's instrument, and that of French.

The armlets used were at least 12 cm. breadth, except in a few of the children, in whom an armlet of 10 cm. was used for the arm observations. When not otherwise stated, the armlet for the lower extremity was placed on the calf. In the few cases in which the thigh was used for compression a specially broad armlet (18 cm. broad) was used. All readings were taken on reducing the pressure after obliteration of the pulse. Unless otherwise stated, the tactile method was the one adopted. In the readings by this method the radial artery was used for the upper extremity, the dorsalis pedis or posterior tibial for the

Cases.

	Sex.	Age.	Diagnosis.	Arm Reading.	Leg Reading.	Difference.	Pulse Pressure.	History.	Remarks—Arteries, Pulse, etc.
Adults.									
1	M.	39	Aortic incompetence, (?) abdominal aneurysm	mm. *173	mm. 208	mm. 35	60 mm. approx.‡	Rheumatic fever, syphilis	Arteries thickened; pulse collapsing.
2	M.	53	Aortic incompetence, dilated aorta	209 184 184 135	234 200 237 135	+25 16 53 0	89 mm. approx.‡	—	Arteries markedly thickened; collapsing pulse.
3	M.	55	Aortic incompetence, dilated aorta	*182	265	83	106 mm. approx.‡	—	Arteries markedly thickened; pulse collapsing.
4	M.	53	Aortic incompetence	195 155	200 190	+5 35	—	Rheumatic fever (recent)	Arteries thickened; pulse collapsing.
5	M.	42	Aortic incompetence	142	139	-3	—	—	Arteries thickened; pulse collapsing.
6	F.	39	Aortic incompetence	198 208 207	208 242 243	10 34 36	72 mm.‡	—	Arteries thickened.
7	M.	32	Aortic incompetence and stenosis	118	215	197	67 mm.‡	(?) Syphilis	Arteries somewhat thickened; pulse somewhat collapsing; artery full between beats.
8	M.	16	Aortic incompetence and stenosis, mitral incompetence	80	110	130	—	—	Arteries normal.
9	M.	40	Aortic incompetence and mitral stenosis (?)	143	131 (thigh)	-12	—	—	Arteries normal; pulse collapsing; temperature at or nearly 100° F.
10	M.	31	Aortic incompetence	*143	148 (thigh)	5	99 mm.	No past illnesses	Temperature about 100° F.
11	M.	40	Aortic stenosis and incompetence	128 154 auditory	135 (thigh) 180 (thigh) 180 auditory	7 26	95 mm.	—	Arteries markedly thickened.
12	M.	43	Aortic incompetence; myocardial degeneration	160 approx. *187	180 approx. 270	20 approx. 83	—	Rheumatic fever and syphilis	Pulse collapsing; arteries markedly thickened; last observation caused pain.
13	M.	37	Aortic incompetence and stenosis	120 approx. 120	100 approx. 80	-20 approx. -40	40 mm. approx.	—	Arteries markedly thickened; pulse slightly collapsing; artery full between beats.
14	M.	31	Aortic incompetence and stenosis; mitral incompetence	145	145	0	115 mm.	Rheumatism and chorea	Arteries thickened; pulse collapsing.
15	M.	39	Aortic stenosis and incompetence; mitral stenosis and incompetence	115	170	155	63 mm.	—	Arteries somewhat thickened; pulse slightly collapsing; artery full between beats.
16	M.	70	Aortic incompetence	210 auditory 190 auditory	250 (thigh) 228 (thigh) 228 auditory	+40 38	130 mm.	—	Arteries thickened.
Children.									
17	M.	12	Aortic incompetence	*95	120	25	35 mm. approx.	—	Arteries normal; pulse perhaps slightly collapsing.
18	M.	15	Aortic incompetence, mitral stenosis and incompetence	133 119 114	122 128 114	-11 9 0	76 mm. approx.	—	Arteries normal; pulse collapsing.
19	M.	10	Aortic incompetence, mitral stenosis and incompetence	*122	122	0	58 mm. approx.	Rheumatic fever	Arteries normal; pulse rather collapsing; artery full between beats.
20	M.	13	Aortic stenosis and incompetence, mitral stenosis and incompetence	*107	97	-10	61 mm. approx.	—	Arteries normal; pulse slightly collapsing; artery full between beats.
21	M.	10	Aortic incompetence and stenosis	125	145	20	70 mm.	—	Arteries normal; pulse collapsing; artery full between beats.
22	M.	11	Aortic incompetence	*135	125	-10	—	—	Arteries normal.
23	M.	13	Aortic incompetence and stenosis, mitral stenosis and incompetence	110	135	25	—	—	Arteries normal.
24	M.	12	Aortic incompetence and mitral incompetence	104 *110	108 90	4 -20	82 mm. approx.	Chorea	Arteries normal. At first observation, pulse not collapsing, compensation fair; at second observation, pulse collapsing, compensation good.

* Indicates that the arm and leg observations were not made synchronously.

† Indicates that the difference was not abolished by recompression.

‡ Auditory method, point of disappearance of sound.

† Indicates that the difference was abolished by recompression.

§ Method of maximum oscillation for diastolic.

lower. Dr. George Oliver's auditory tambour attached to a binaural stethoscope was used for the auditory readings. The tambour for the arm readings was placed at the bend of the elbow, and for those on the lower extremity at the popliteal space.

In the auditory readings the point of appearance of the audible throb was taken as the systolic reading, and unless otherwise stated the point at which the loud note becomes suddenly dull before disappearing was taken as the diastolic reading. In a few of the early auditory readings the diastolic index was taken as the point of disappearance of the sound (Cases 6 and 7), and in the first observations (Cases 1, 2, 3) the approximate visual method of the point of maximum oscillation was used. The pulse pressure was taken as the difference between the systolic and diastolic readings in the arm.

Corresponding observations on the leg and arm are recorded on the same line, and, unless otherwise stated, were made synchronously, one manometer being connected by means of a T-shaped tube with both armlets. Observations of any case on different lines were made on different days. The patients were at rest in the recumbent position, and the results are recorded in millimetres of mercury.

Details.

This paper is based on observations of 24 cases of undoubted and, indeed, well marked aortic incompetence. In 14 of these a difference obtained in the arm and leg readings. In the 10 remaining cases there was either a higher arm than leg reading or there was no difference in favour of the leg reading. Two of the cases which did not show the difference had temperatures of 100° F. or nearly 100° F. at the time of examination. Rolleston,⁴ in his observations, found that in two cases temperatures of 101.4° F. and 100.4° F. respectively caused the difference to diminish greatly or to disappear. It is possible, therefore, that the slight degree of fever may have explained the absence of the difference in my two cases, so that I have thought it well to exclude these two cases from the list. This leaves 22 cases to be considered, of which 14 were adults and 8 children of from 15 years downwards. Of these 22 cases, 14 showed the difference, leaving 8 in which the sign did not obtain.

It is to be noted that 10 out of the 14 cases showing the difference had thick arteries (all except Nos. 8, 17, 21, and 23), or of all the 13 cases of the series having thick arteries 10, or all but 3 (Nos. 5, 13, and 14), showed the difference, a much larger proportion than that which obtained in the total number of cases (14 out of 22) all of which had, of course, aortic incompetence. Further, in the 14 adult cases 11 showed the difference, whereas by contrast it was observed only in 3 of the 8 children, in none of whom were any arterial changes found. These two points suggest very strongly the arterial origin of the phenomenon.

In regard to the relationship of the degree of pulse pressure to the arm-leg difference, there was no correspondence between the two. Taking the 6 highest pulse pressures, in 4 of them, Nos. 2, 3, 11, and 16, there was a marked difference, whilst, on the other hand, in the 2 others, Nos. 10 and 14, no appreciable difference was found. In 3 out of 5 cases in which it was tried, the difference of pressure was abolished by repeated recompression, which again is as we have seen in favour of the arterial explanation, and is at variance with Hill's finding aortic regurgitation.

It will be seen, if my results be compared with those of the various writers on this subject that they are similar to those of Heitz, but differ from those of the others, and it may be pointed out that Heitz's conclusions appear to be based, to judge from the published records, upon a considerably larger number of cases than those of the other observers.

Whilst of course I freely acknowledge that the limited number of cases upon which the present paper is based is insufficient to form the basis of general dogmatism, it is certainly quite sufficient to show that the phenomenon is nothing like a constant sign of aortic regurgitation. Finally, I would submit with some insistence that future observations should include in each case a note upon the condition of the arteries, an omission as to which has deprived past records of much of their value.

REFERENCES.

- ¹ *Heart*, vol. i, p. 73, 1909-10. ² *Ibid.*, vol. iii, p. 219, 1911-12. ³ *Therapeutic Gazette*, 1910, Third Series, xxxvi, p. 457. ⁴ *Heart*, vol. iv, No. 1, 1912. ⁵ *BRITISH MEDICAL JOURNAL*, March 28th, 1914. ⁶ *Paris Méd.*, 1917, xxiii, 112-122. ⁷ Clinical Observations on the Influence of the Vessel Wall on So-called Arterial Blood-Pressure Readings, *Proc. Roy. Soc. Med.*, May, 1909. ⁸ *Haemodynamik*, Leipzig, 1850.

An Address

ON

ELEMENTARY PSYCHOLOGY IN GENERAL HOSPITAL TEACHING.

DELIVERED BEFORE THE ULSTER MEDICAL SOCIETY,
MARCH, 1921.

BY

WILLIAM CALWELL, M.D.,

PHYSICIAN, ROYAL VICTORIA HOSPITAL, BELFAST; MEDICAL OFFICER
IN CHARGE, CRAIGAVON HOSPITAL FOR NEURASTHENIC PENSIONERS,
BELFAST.

In August, 1920, I was instructed to examine a woman aged between 60 and 70, and report on her mental and general condition; she had had a stroke in 1912 and another two years later. She was bedridden. Both arms and legs were strongly flexed, the knees nearly touching the hips, the heels the hips, and the arms were crossed over the abdomen; she had no control over bowel or bladder; her articulation was very defective, resembling an exaggerated mumbling lisp; orientation for time and space was hopelessly poor; she said she was 43, and that it was the year 1909; her memory was quite unreliable, her ideas extremely limited; she knew nothing about her affairs except that she got a certain sum a week from a deceased sister, and had been robbed of her furniture, and in making these statements she exhibited tearful but short-lived emotion.

I had no difficulty in making a report. In a subsequent letter I was asked whether I could give an opinion as to the time during which this patient had been incapable of managing her affairs, and whether there was any ground for forming an opinion as to what the mental capacity of the patient was likely to be after the seizure in 1912.

In trying to answer these questions I realized that my knowledge was very limited. It could be put into two statements: (1) After a stroke there is mental weakness, which may be of so slight a degree as to be unnoticed, or may amount to well marked secondary dementia, where the patient has returned to second babyhood; and (2) that there is a question how far aphasia meant intellectual incompetence. I answered that it was impossible to form a reliable opinion.

So I turned to our textbooks and to what medical literature I had time to consult. Some simply mentioned the question of examination of the mental condition of our patients, some gave a rough outline, a few lay emphasis on the "cerebral" features of arterio-sclerosis, by which I presume they meant "mental" or "psychic" features. This is the "arteriopathic dementia" of the alienist. It is interesting and instructive to note how even some of our best textbooks do not refer, or refer in an inadequate manner, to the difference between cerebral physiology and psychology. The textbooks on mental diseases are not included in this statement. Many of our manuals of medicine and monographs on the diseases of the nervous system consider elaborately the questions of cerebral pathology and of aphasia; to illustrate the latter subject the pages abound with diagrams which, rather unfortunately, recall from our unconscious memory the forty-seventh proposition of the first book of Euclid, and we are apt to associate with these diagrams the same childlike faith in the definiteness of proof and of absolute reliability that we learned in our boyhood to associate with the famous right-angled triangle; normal or morbid psychology is not touched upon.

Charcot, in his classical clinical lectures, takes up a very interesting attitude: the study of cerebral haemorrhage, he says, forms an introduction to the study of diseases of the encephalon. The sudden tearing of the tissue realizes in some degree the simplicity sought in experiments on animals, but he omits the mental condition. His further lectures were unpublished.

Our later textbooks and manuals on insanity do supply the want; whether in an altogether satisfactory manner or not is a legitimate subject of debate, but the effect of this separation is that the student and most medical men do not attempt to make even an elementary psychological examination of a patient, unless this patient is actually insane, and then only so far as to ascertain whether insanity exists or not. But a patient's mentation is a function of his brain; this function is continually being carried on; it is a function—not perhaps from a vital point of view, but certainly from a social point of view—which is incontestably the most important, and it should not be relegated to an odd half-dozen of lines.

Cerebral physiology is yet very incomplete, and so we should teach the elements of psychology to the student as part of the clinical examination of the most important function. At present it is left to the native wit of the doctor to size the mentality of a patient. Should we not train him in scientific methods? Would we give up our scientific and regular, orderly, means of physical examination because the native wit of Hippocrates enabled him to do so much? Let us rather correlate cerebral physiology and psychology to make as complete an examination of the functioning of the nervous system as our present knowledge allows, and obtain practical aid in the examination of our patients.

An examination of some of our excellent manuals on clinical examination and case-taking showed the same deficiency; the reflex movements of the big toe were strongly in evidence, but the powers of memory, of ideation, the depth of emotion, the occurrence and kind of dreams, and the bias of mind, the powers of expression, of calculation, were passed by unnoticed or barely noticed; Hutchison and Rainy in their *Clinical Methods* give a page on which is printed a special scheme for cases presenting mental symptoms drawn up by Henry Head for use in the wards of a general hospital. A student, however, is told in other places to practise examination on healthy lungs, hearts, urine, etc. Why should he not practise a little normal psychology, and so be better fitted to examine and recognize psychological abnormalities when they present themselves?

A student should have a certain amount of both normal and morbid psychological training before going to the asylum; he will be able to benefit by his training in that