but there is considerable danger of tearing the uterus by operation during the spasm.

Incision of the ring per vaginam is too dangerous to be attempted. Version is absolutely contraindicated; it is, moreover, often impossible.

The application of forceps is very rarely helpful. Where the ring is around the child's neck the blades almost invariably slip. If they do succeed in dragging the head down to the vulva it simply carries the ring down with it, unless the latter has been torn in the process. A case has been reported where the ring had formed around the shoulders of the second of twins, and where very gradual traction with forceps caused a tear of the vagina, cervix, and lower uterine segment into the broad ligament. The application of forceps, therefore, is not advisable in the case of a living child. Where the child is dead forceps with cleidotomy and steady traction will doubtless be sometimes successful.

Craniotomy also has but a limited field, even within that class where the child is already dead. Embryotomy has succeeded in a few cases, in others it has been impossible to deliver even after evisceration.

There is no doubt that where the child is dead the ring in the majority of cases will yield to slight steady traction if that part of the foetus endeavouring to enter the ring be cone-shaped. Division of the clavicles will therefore be of use where the shoulders are obstructed. Again, in that case of breech presentation reported by Lady Barrett to which I have already referred, a hydrocephalic head was successfully extracted through the ring after puncture of the spinal canal.

There remains but Caesarean section. Mr. Munro Kerr has said very recently that probably Caesarean section will in future be more frequently performed for the relief of this condition. And it surely is the ideal treatment for obstinate cases if only the uterus be not already infected. Where infection has already occurred the section should be followed by hysterectomy, or, if the child be dead, a hysterectomy alone is indicated.

With regard to treatment I am impressed by the following facts:

1. The difficulty, danger, and high mortality attending active interference.
2. That the only abnormal condition is a localized spasm of the uterine muscle.
3. That deep anaesthesia diminishes muscular spasm.
4. That repeated examinations increase spasm and are therefore contraindicated.
5. That in the case I have reported deep and prolonged anaesthesia alone resulted in the spontaneous delivery of a healthy child.
6. That in the previous confinements of this patient forceps were applied on every occasion, with an infant mortality of 50 per cent.

As a result I consider that the correct treatment for this condition is *deep anaesthesia, if necessary prolonged, and nothing else*. This treatment will, I believe, very nearly eliminate the necessity for Caesarean section.

BIBLIOGRAPHY.

- Holland: *Journ. Obstet. and Gyn. Brit. Emp.*, 1911, xix, p. 526.
 De Lee: *Principles and Practice of Obstetrics*, 1913, p. 574.
 Andrews and Maxwell: *Proc. Roy. Soc. Med.*, 1908, i, Obstetrical and Gynaecological Section, pp. 138-50.
 Clifford White: *Ibid.*, 1913, vi, pp. 70-85.
 Barrett: *Ibid.*, p. 86.
 Jardine: *Journ. Obstet. and Gyn. Brit. Emp.*, 1911, xx, p. 11.
 Hedley: *Pract. Encyclop. of Mid. and Dis. of Women*, ed. Fairbairn, 1921, p. 204.
 Galabin: *Manual of Midwifery*, 1897, p. 754.
 Tweedy: *Practical Obstetrics*, p. 263.
 Willett: *Proc. Roy. Soc. Med.*, 1912, v, p. 335.
 Wilhelm, J.: *Missouri Med. Assoc.*, 1910, xvi, p. 295; quoted *Surg., Gyn., and Obstet.*, Abst., January, 1920, p. 56.
 Munro Kerr: *BRITISH MEDICAL JOURNAL*, 1921, October 1st, p. 519.

RECTAL ADMINISTRATION OF TARTARATED ANTIMONY IN BILHARZIASIS.

BY

H. F. WILSON, M.C., M.B., B.CH.,

DAVID GORDON MEMORIAL HOSPITAL, LIVINGSTONIA, NYASALAND.

ALTHOUGH a number of papers have appeared recently on the treatment of *Schistoma haematobium* by tartar emetic, the method invariably advocated has been intravenous injection. For any parasitic infection of the general circulation this could hardly be improved upon, as the drug enters the blood stream in such quantities and at such precise times as the introducer desires. But where it is desired to destroy a parasite whose habitat does not generally extend beyond the portal circulation it seems more reasonable to introduce the

parasiticide directly into the radicals of that system, so that it may meet with what it is intended to kill in greater concentration than if injected into a superficial vein and accordingly distributed to all parts of the body before reaching its object.

As, however, accessible veins of the portal circulation are not available for injection, we are thrown back upon the absorbing powers of the rectal mucous membrane, reached by solutions of the tartarated antimony given as small enemata. The drug absorbed in this way enters the superior haemorrhoidal vein, where, in cases with rectal symptoms, the worms are understood to lay their eggs.¹ The free intercommunication between this vein and the vesical plexus allows of the drug passing to the bladder wall, in which oviposition is the rule in cases with haematuria or other vesical symptoms. That part of the dose which does not find its way into the vesical veins passes up to the portal vein itself, where the pairing of the worms takes place, and so reaches the liver, where the younger immature forms are found, and in this way exerts its maximum action upon all stages of the creature's life cycle before being merged into the general circulation, in which worms are only found, Kartulis tells us, in the inferior vena cava; this is the first part to be reached by the antimony.

Whether the portal system can be "sterilized" by the administration of a few large doses in this way, and the treatment reduced to, at most, one or two administrations, must be left for further investigation. The purpose of this preliminary note is to draw attention to the tolerance of the rectum to what would be considered very large doses if given in any other way, and in particular to emphasize the convenience of this method where the personal idiosyncrasy of the patient, or, in women and children, the difficulty of locating and entering a vein, makes the intravenous method undesirable.

Indeed, as an alternative it must always be of practical value to practitioners who, for whatever reason, do not go in for intravenous work, or who find the necessary sterilization of solutions, needles, etc., an undesirable call upon their time, apart from the need for giving an intravenous injection personally. Finally, experience of the high lying plateaux of Central Africa goes to show that, both in the rains and in the cold season which succeeds them, many days are most unsuitable for finding veins in the subcutaneous tissues of naked and shivering patients; at times weeks have passed without days sunny enough to give what may be called "good veins." At such times, when many cases of bilharziasis are waiting treatment, the method advocated here is a real boon.

Four cases only will be cited to illustrate the possibilities of intrarectal treatment.

CASE I.—Child with Old-standing Bilharzia Infection of the Bladder Cured after Fourteen Intrarectal Injections.

John, a half-caste boy aged about 7, who had come up country from Southern Rhodesia, eighteen months before, complained of suprapubic pain and of passing blood at the end of micturition; these symptoms were of some four or five months' duration. There was a history of a streak of blood in the stool, which had defied treatment for some time, but at that time no search seems to have been made for schistosome eggs in the faeces. Examination of uncentrifuged specimen of urine revealed terminal spined ova in astonishing numbers; the specimen was red and loaded with urates. No eggs were found in faeces. Treatment was commenced with an injection of 1 grain in 100 c.cm. of water at body temperature. A few drops of tincture of opium were added, but as there was no intolerance this was not repeated. Two days later a dose of 2 grains was given in the same way, and thereafter for three weeks the drug was given every second day in gradually increasing doses—2 grains, 3 grains, 4 grains, 5 grains, etc. During the first week the urine cleared of blood, but after eleven doses a few shrivelled eggs were still to be seen. The 12-grain dose was repeated twice, and two days later 14 grains were given, after which no sign of an egg was found on examining the last part of a specimen passed in the morning on rising and centrifuging the sediment after some hours' standing. At no time in the course of treatment did the boy show any sign of nausea, nor were his appetite or spirits in any way affected. He received in all 104 grains.

CASE II.—Child with Enlargement of Liver, Ascites and Anaemia: Schistosome Ova in Faeces: Cured after Fourteen Injections.

Simiyon, a boy of perhaps 6, was brought in with a much-distended belly, containing a quantity of free fluid; the liver was two fingerbreadths below the costal margin. The boy was anaemic and listless.

Examination of the faeces showed numbers of ova, chiefly lateral-spined in type. He was treated *pari passu* with Case I, and in the same doses up to 12 grains; this was then followed by one of 16 grains. No eggs were found thereafter; the abdomen had decreased in size so as to be no larger than that of the run of native children of his age, and although the liver was still easily palpable

the free fluid had disappeared. Under tonics the anaemia improved, and he was dismissed after six weeks a totally different boy, having received in all 108 grains without showing any tendency to nausea or vomiting.

CASE III.—Lad with Haematuria Cured in Eight Attendances.

Naaman, an apprentice carpenter, aged about 17, complained of passing blood in the urine for the previous month or two. The urine was smoky, and on centrifugalization showed ova of bilharzia. He came every other day for three weeks, receiving doses of 2 grains, 4 grains, 6 grains, etc. The urine cleared after the third injection, and by the time that he had received eight doses, amounting in all to 68 grains, the urine was found to be free of ova. He spent his midday rest hour receiving the injection and lying down, but otherwise was not away from his ordinary work.

CASE IV.—Old Man with Pronounced Haematuria Cured after Four Injections Administered in Five Days.

This old man could give no history of his complaint, but the urine was of a thick port-wine appearance with a sediment of urates. Ova were found easily. The first dose was 5 grains; this was followed two days later by a dose of 8 grains; the urine was then amber in colour without a trace of blood. On the next two days he received successively 10 grains and 12 grains; treatment was then discontinued as the centrifugized specimens of urine examined contained no more ova. He had been given 35 grains in the five days without showing any symptom of intolerance.

The treatment was in all these cases carried out in the following manner: The bowels were caused to move daily in the morning by giving if needed a mild laxative at night. Some three hours afterwards the patient lay down upon his left side, and the dose, made up to about one or two ounces by the addition of warm water, was administered by the hospital orderly by means of a funnel attached to a No. 12 rubber worm catheter, which after being lubricated was passed well up into the rectum; it is advisable to compress it firmly between the finger and thumb as it is being withdrawn so that the whole of the solution be introduced. The patient has been directed to lie down for a couple of hours, but whether this is essential has not been finally settled. In some cases the bowel must be cleared first by a soap and water enema, but this is not the rule by any means.

Conclusions.

1. The results of intrarectal administration of tartarated antimony are in no way inferior to intravenous.
2. It economizes time, and is free from all risk.
3. Nausea and vomiting occur less frequently than when the drug is run into the general circulation.
4. Case IV suggests the possibility of reducing the number of attendances for treatment considerably.

REFERENCE.

¹ Castellani and Chalmers: *Manual Trop. Med.*, second edition, p. 482.

SOME FUNCTIONS OF THE SUPRARENAL GLANDS.

BY
EVELYN E. HEWER,

LECTURER IN HISTOLOGY AND ASSISTANT LECTURER IN PHYSIOLOGY
AT THE LONDON (ROYAL FREE HOSPITAL) SCHOOL OF
MEDICINE FOR WOMEN.

THE medullary portion of the suprarenal gland (derived, with the cells of the sympathetic ganglia, from the neuroblasts of the central nervous system) is well known to be concerned with the elaboration of an internal secretion which contains as one of its active principles the substance known as adrenaline; but the part played in the animal economy by the cortex (developed from the mesothelium of the genital ridge) is still obscure. In view of the vital importance of this organ some experimental results recently obtained in rats seem to be worth recording. Even though there is no evidence that the suprarenal gland of rats is functionally homologous with that of man, yet there is no evidence to the contrary, and some of the findings are most suggestive.

The experiments consisted of feeding or inoculating white rats of varying ages with different preparations of suprarenal cortex for varying periods, and then killing, and examining various organs histologically. The preparations used were as follows, and I am much indebted to these firms for their courtesy and assistance in obtaining these preparations:

1. Adcortogen (by Carnrick Co.), which was dried cortex, and contained some adrenaline.
2. Cortex lipoids (by British Organotherapy Co.), which was a liquid extract containing only the lipoids.
3. Cortex extract B (by Carnrick Co.), which was a watery extract of cortex, containing no lipoids but some adrenaline.

4. Cortex extract C (by Carnrick Co.), which was the same as B, but free from adrenaline.

5. Adrenaline (by Parke, Davis). This was used to obtain control animals for those treated with cortical preparations which were found to contain adrenaline.

In all, 118 animals were used, and the following are the most striking of the results obtained. It is not proposed to give here all the details of the experiments, as it is hoped that the work will be published in full elsewhere.

The general health of the animals was not affected, except rarely when using adrenaline-containing extracts. But an extraordinary formation and deposition of fat was obtained in most of the animals, thus confirming the observations of others that the suprarenal cortex plays a part in controlling the fat metabolism of the body, which may thus become perverted in hyperactivity of the gland. In addition, nearly all the rats developed a curious condition of the coat; the fur was thick, fine and glossy (as it should be in health), but came out very freely on handling; the hair follicles microscopically showed degeneration. This condition was not due to the adrenaline, and points to a connexion between suprarenal cortex and calcium metabolism, as perversion of the latter is known to give rise to nutritional disturbances of the skin.

The organs to which attention was directed more particularly were those of internal secretion, and the following are the most conspicuous of the results obtained.

Thymus.—The gland was obviously affected by the treatment. There was most marked increase of invading mast cells; some of these cells are apparently always present in this animal, but the significance of this great increase is not clear. The gland was apparently responding to stimulation in the direction of involution, as indicated by an increase in the Hassall's corpuscles and chromophobe complexes, normally so scanty in the rat. In many cases, also, there were present large clear eosinophil cells of endothelial type invading the gland by the lymph channels, and sometimes completely destroying the normal structure. The appearance of these cells, both here and in the haemolymph glands, can hardly be a specific effect of the treatment, but probably indicates some degree of lowered resistance, the lymph glands being the organs first attacked.

Pancreas.—The gland appeared normal, but two points of interest may be mentioned. First, in the series where adrenaline was being given, a definite thickening of blood vessel walls, both here and in the spleen, was frequently found; secondly, lymphocytes were usually present in the islets. This lymphocyte increase was not confined to the pancreas, but was found in many organs, and is interesting in view of the increase commonly found in Addison's disease. The suprarenal gland, as a whole, appears to stimulate the appearance of lymphocytes in the tissues (no blood counts were made); perhaps there may be some connexion between the sexual immaturity of status lymphaticus and the influence exerted by the suprarenal cortex on the gonads.

Spleen.—Generalized lymphocytic increase was well shown, and the vascularity and blood pigments were very constantly increased. Adrenaline appears to affect the organ adversely; the capillaries were found engorged with blood, lymphocytes were frequently found below the endothelium of the blood vessels, and the typical structure was lost. Increase of giant cells was frequently observed, but no evidence of tuberculous infection.

Male Gonads.—The treatment at first hastened the genital development, but, when continued for some time, in every case degeneration of the seminiferous tubules occurred. The breakdown was very marked, but after a long interval some repair was usually found. The interstitial tissue was also affected, the lipoid constituent causing a cellular decrease and the non-lipoid constituents an increase; in this way a balance may be kept by the full action of the gland. The interstitial tissue contained a large quantity of eosinophil exudate in nearly every case.

Female Gonads.—Sexual precocity in the female rat is difficult to determine experimentally. Degeneration of the older follicles was always found; the younger follicles appeared normal, and therefore were either more resistant to treatment or had undergone repair. The interstitial cells were constantly increased.

Suprarenals.—Evidence was obtained that normally tells atrophy at the medullary border, and that regeneration occurs from the zona glomerulosa; this change was found to be accelerated in pregnancy, and could be stimulated by treatment with cortical preparations more easily in the female.