surrounded by fibrous tissue, and the whole encapsuled by a laminated capsule, is irresistible. When the mass of choroid plexus, or arachnoid tuft, is completely invested and growth arrested, the sequestered epithelium becomes impregnated with cholesterol sand. Hence the picturesque names—psammoma, cholesteatoma, and pearl tumour; some gleam like mother-of-pearl.

The infiltration of these tumours with cholesterol resembles the deposition of lime salts in dead uterine fibroids.

A psammoma infiltrated with cholesterol is a dead tumour, but it rarely dies before it has wrecked the life of the man or woman by damaging brain or spinal cord.

The clinical symptoms produced by these extraordinary tumours sometimes permit of exact localization; often they are complicated tangles, difficult to unravel even with the help, separately or combined, of a neurologist, otologist, ophthalmologist, radiologist, and a physician. In this lecture I am mainly concerned with pathologists.

VILLI IN GENERAL.

Villi of various kinds play a large part in our lives, mainly for good, sometimes for evil. Soon after conception the chorion of the embryo becomes shaggy with exquisite villi, and although the villi are extremely delicate structures they are able to erode the maternal structures with which, in the normal course of events, they come in contact, embed themselves and develop a placenta.

It occasionally happens that they fail to form a placenta, change into peculiar grape-like bodies and transform into what is commonly known as a "hydatid mole." Some years ago obstetricians were startled by the discovery that such erratic changes in the villi sometimes lead to disastrous consequences in fertile women, for these altered villi behave like malignant tumours. They erode blood vessels, colonize the blood, disseminate and quickly cause death. This particular change is now known as chorionic cancer.

There is another variety of villus which occurs only under pathological conditions. Occasionally the ovary is occupied

FIG. 4.—Psammoma of the fourth ventricle. From a woman, aged 32. (Museum, Royal College of Surgeons.)

FIG. 5.—Structure of a psammoma.

by a cyst, usually unilocular, and its walls are covered by crops of papillomas closely resembling, in structure and appearance, the grape-like bodies of hydatid moles. The papillomas are able to erode the cyst wall and project on its outer surface.

There are two varieties of ovarian papillomatous cysts. One is clinically innocent. The other exhibits malignant properties, for it is not only erosive but it recurs locally and disseminates. I have published an account of one which set up secondary deposits in the mediastinum, and the papillomatous buds eroded the sternum and appeared as papillomatous cysts under the skin. When the papillomas erode the walls of the cyst the minute collections of shed epithelium become engrafted on the peritoneum, and warts, sometimes in thousands, arise from them. When examined microscopically such implanted warts are often found to be encapsuled by peritoneal exudation, and thus recall in miniature the encapsuled psammomas of the meninges.

CONCLUSION.

The study of these conditions, I venture to think, helps us to understand the nature of the psammoma. This tumour begins in an overgrown clump of choroid plexus, or arachnoid tufts. The epithelium capping the clumps is bathed in cerebro-spinal fluid, and becomes enclosed in a capsule of plastic exudate which stratifies and organizes. It is known that peritoneal exudation will encapsule sterile foreign bodies, ovarian papillomas, calcified epiploic appendices, intestinal and subserous uterine fibroids, and even a coagulated collection of sterile fluid or blood clot.

The villi of the arachnoid erode bone, as the Pacchionian foramina prove; chorionic villi, normal and abnormal, can penetrate the walls of the uterus, and ovarian papillomas in a favouring environment erode bone.

Surely this is sufficient evidence to justify the removal of psammomas from the miscellaneous group at present labelled endotheliomas.

THE TREATMENT OF GASTRIC ULCER.*

BY

SIR WILLIAM HALE-WHITE, K.B.E., M.D.,
CONSULTING PHYSICIAN TO GUY'S HOSPITAL.

THE treatment of gastric ulcer labours under two great preliminary difficulties. The first is that it is often diagnosed as present when it is not. That this bears on treatment is shown by the very many gastro-enterostomies that have been performed for a supposed ulcer when none has existed; sometimes the gastro-enterostomy has had to be undone later.

Among the conditions the symptoms of which have often been erroneously ascribed to gastric ulcer is disease of the teeth and gums; cure of this may lead to the disappearance of symptoms thought to be due to a gastric ulcer. In the middle of the last century it was believed that young women suffered from haematemesis due to vicarious menstruation, and in one large textbook this was stated less than thirty years ago. When this belief was discarded, young women with dyspepsia and haematemesis were said to have a gastric ulcer, but gradually doubt was thrown on this view. Then, although these young women rarely died of their disease, a

few deaths occurred, and no ulcer was discovered. Some surgeons opened the stomach for this bleeding and observed that the blood did not proceed from an ulcer but from a general oozing over a considerable area. To this condition the name "gastrostaxis" has been given. It is commonest in young females, is occasionally seen in young men, and rarely kills. The opinion has been expressed that this condition is one of minute multiple ulcers, but there is evidence of its being a distinct clinical entity. Many cases of gastrostaxis are diagnosed as ulcer, and, as they almost always recover, the treatment of gastric ulcer may be thought to be more successful than it is.

There is no doubt that disease of the appendix, of the gall bladder, and even a stone in the kidney, may cause symptoms that are set down erroneously to gastric ulcer. Duodenal ulcer, carcinoma of the stomach, and various functional dyspepsias may be mistaken for it. All these conditions lead to errors of diagnosis sufficiently often to make most opinions expressed as to the results of treatment of no value. Other diseases are occasionally confounded with gastric ulcer, but not frequently enough to invalidate our views as to treatment.

When we consider the individual symptoms of gastric ulcer we also find ourselves in difficulties. Haematemesis may be completely absent, even when the ulcer is large, and is frequently seen in other diseases. The same is true of occult

* Being the introductory remarks at a debate on the treatment of gastric ulcer at the Royal Society of Medicine, February 2nd, 1922.

blood. Some sufferers from ulcer never have pain, many only complain of it at long intervals. It is common enough in various other conditions. Vomiting is an inconstant symptom, indeed very frequent sickness is strongly against the presence of an ulcer. Neither deep nor superficial tenderness are reliable symptoms, and it is generally allowed that an examination of the gastric contents may be of no help. My object is not to imply that the diagnosis of gastric ulcer is impossible—indeed, a skilful clinician can often infer its presence pretty confidently—but to show that the conditions which may be mistaken for it are so numerous and common, and that its own symptoms are so variable, that most statistics as to the success or failure of medical treatment are quite untrustworthy.

Examination by the x rays will often reveal an ulcer itself, it will also show the spasm of the gastric muscle (incisura) opposite to it. By this means we may obtain also further indications of the presence of an ulcer, such as hour-glass spasm, diffuse spastic distortion and delay in emptying from pyloric spasm, organic hour-glass stomach from cicatrization of an ulcer, and increased peristalsis from organic pyloric stenosis. Many radiologists supply us with beautiful pictures that are invaluable, but those produced by some are of little use. It may be that an ulcer although present is not demonstrated by the x rays, but that does not alter the fact that the only way in which we can be so sure of the existence of an ulcer that the case should be used in statistics employed to estimate the efficacy of medical treatment, is to see the ulcer with the x rays. It must be actually seen, for although the various manifestations of spasm are often highly suggestive of its presence, yet each of them may be evident when no ulcer exists. What is sorely needed is a series of cases showing on an x ray plate an ulcer, and examined again with the x rays six months later to see if the ulcer is healed.

Our second difficulty is concerned with causation. We knew how to treat a few diseases—for example, malaria and syphilis—before the cause was discovered, but generally treatment is imperfect until the cause is apparent. Clinical observations and experimental work indicate that a gastric ulcer is due to the local effect on the stomach of bacteria or toxins produced in another part of the body, and that when the gastric patch damaged by them has been formed the excavation of the ulcer is promoted and its healing retarded by the acidity of the gastric juice. It is probable that the incriminating factor is not always the same and that there are different varieties of ulcer due to different causes. Whether ulcers in human beings are formed by endogenous gastrotoxins is not known, but judging by Bolton's experiments it is possible. Still we do not yet know the cause of a gastric ulcer with anything like the certainty with which we are able to say that a diphtherial ulcer is caused by the diphtheria bacillus or a syphilitic ulcer by the *Spirochaeta pallida*. Putting aside traumatic ulcers and those due to irritant poisons, only twice have I been able to say, even after death, that I really knew the cause of an ulcer; once there were definite tubercles on its peritoneal covering and I saw the beautiful specimen exhibited here by Dr. McNee showing syphilitic spirochaetes in the wall of an ulcer. Nevertheless there is sufficient evidence for the view above mentioned for it to be strongly advisable, in every case of gastric ulcer, to remove any distant possible source of infection, be it in the teeth, appendix, or gall bladder, disease of any of these being commonly associated with gastric ulcer. More rarely such a source may be elsewhere—for example, tonsils or nose.

We are much encouraged in the treatment of ulcers of the stomach by the undoubted fact that they show a natural tendency to spontaneous cure. Those inadvertently caused by operations get well, those produced by experiments on animals heal under appropriate conditions, many seen in the *post-mortem* room show signs of healing, indeed they may be found quite healed; the fact that an ulcer by its cicatrization causes an hour-glass stomach or pyloric stenosis indicates that the attempt at cure has been made.

Perforation from an ulcer is much more common in women than in men; *post-mortem* evidence shows us that in women ulcers are more frequently multiple than in men. Therefore, it appears that we know of two distinct varieties, one multiple in many cases, a frequent cause of perforation and found chiefly in women, usually between the ages of 20 and 30; the other usually presenting itself clinically as a chronic ulcer, but not so often seen to perforate nor so frequently multiple, and found chiefly in men between the ages of 30 and 40. It is very likely that there are subdivisions of each variety

depending upon the infective agent. The first is usually spoken of as an acute ulcer, the second as a chronic. This terminology is unfortunate and confusing; the words "acute" and "chronic" merely refer to time. It is believed that some in the group named chronic may begin as acute, but if there are two distinct groups with different characteristics, should the disease be of the first variety it cannot be changed to the second because it has lasted a long while.

Faber of Copenhagen² also describes two distinct diseases under the term ulcer of the stomach. The first, corresponding to acute ulcer, is ulcer of the corpus or "woman's ulcer"; it is commonest in young women, is usually situated on or near the lesser curvature; apart from perforation the symptoms are not severe and in many cases unrecognized, the ulcer getting well spontaneously. The second, corresponding to chronic, he calls juxta-pyloric ulcer; it is commonest in men, its symptoms are frequently severe and often need surgical interference. But these names, although better than acute and chronic, are far from perfect, for the first variety is not confined to women nor the second to men, nor is the ulcer invariable in its situation in either. It would tend to clear thinking to abolish all these names, which are attempts at sharp cut descriptive distinctions which do not exist, and to call the first commonly spoken of as acute, ulcer A, and the second, ulcer B.

The first usually does not reveal itself on x -ray examination, and generally gives only slight symptoms; indeed, there may be none until perforation or severe haemorrhage takes place. Therefore it is often very difficult to diagnose, and especially so because, as we have already seen, the diagnosis of gastric ulcer has many pitfalls. Consequently it is impossible to gather accurate statistics as to the result of treatment. All we can say is that there are a number of patients, especially young women, who would a few years ago have been said to be suffering from a gastric ulcer, but some of whom we now believe owe their symptoms to other conditions already mentioned at the beginning of these remarks; others may have what is called an acute ulcer, a woman's ulcer, or ulcer A.

What shall we do with them? First search for a source of infection and, if possible, remove it; this may lead to the disappearance of the symptoms. If it does not, the outlook is still good, for the natural tendency of this class of ulcer is to heal completely. Put the patient to bed; rest in itself is good, and our diet will not contain enough calories for much work. The food must not be such—for example, meat or meat extractives—as will lead to the secretion of much acid; it must not be mechanically or chemically irritating; it must be capable of passing quickly and easily through the stomach; it must not stimulate this organ to excessive contraction, and must be sufficiently nutritious. I am accustomed to employ milk and raw egg, beaten up in it, in the proportion of two eggs to each pint of milk, giving $7\frac{1}{2}$ fluid ounces of this every two hours during the sixteen hours the patient is awake, and an extra feed if she happens to wake in the night. It is almost always unnecessary to peptonize the milk, but this may be done if the patient does not mind the taste. This gives 1,600 calories a day. Unless she has a distaste for it, she should take some olive oil; she is much more likely to be able to do this if the very best quality is given, that, namely, which comes from the first expression; if necessary, it may be flavoured with a little black coffee; the patient may have it as often as she cares for it in the day, provided it is not nauseating. Its advantages are that it is a food, bland and unirritating. It restrains the secretion of acid and it is a mild aperient. One fluid ounce contains 560 calories. The total daily caloric value of our diet is now about 1,700 calories, which is enough for a patient lying in bed; she does not lose weight on it. Ulcers cannot be expected to heal if the body is under-nourished. Any dietary which leads to considerable loss of weight is wrong. In former days, when rectal feeding was common for gastric ulcer, many patients failed to get well because they were starved severely; it is impossible to prevent this on rectal feeding.

The chief drug should be an alkali to counteract the gastric acidity; sodium bicarbonate is frequently chosen with excellent results, but if it leads to distension lime water or magnesia may be substituted. Not less than 25 grains of bicarbonate three times a day should be given. Tincture of belladonna checks the secretion of acid and excess of gastric movement. A good mixture is sodium bicarbonate gr. 25, tincture of belladonna m 10, bismuth subcarbonate gr. 15, tragacanth powder q.s., with chloroform water to make

one fluid ounce; this may be given thrice daily. If an aperient is required, magnesia is the best. This treatment can generally be easily borne for six weeks; it is very simple. I have used it for many years, and patients do well on it. Several authors change the diet as the case progresses. There is no need for this; we cannot see how the cure is proceeding, and we are more likely to cure and prevent relapses in proportion as we are thorough in treatment. Many diets published are complicated; the simpler they are, the more likely are directions to be obeyed.

Cases are much too often treated in a half-hearted manner. If the doctor decides that the patient ought to be treated as though she had a gastric ulcer, let her be so treated, and not just sent away with a bottle of medicine. We have no means of estimating, in the class we are now considering, how much the stomach was ulcerated when the patient was first seen, nor how the cure is progressing, therefore I would suggest certainly not less than four weeks on the treatment—I always try to get six. Perhaps some of the mildest cases are thus treated too long, but it is better to do this than to treat some for too short a time, so that they fail to get well or relapse. The time spent in bed must vary with the case: four weeks may be desirable, but often it is well, earlier than this, to get the patient up in a chair for part of the day, out of doors if possible, provided she is warm. A little massage to the limbs helps to keep her in good condition. When the period on this diet has come to an end she may by degrees return to normal food, but this should always be plain, without much meat, and she should always masticate thoroughly. Unless perforation takes place these patients do not require the aid of a surgeon.

Although, thanks to *x* rays, chronic ulcer, or ulcer B, is much more easy of diagnosis, the treatment is more difficult and uncertain. We badly want, as I have already said, a series of cases with *x*-ray pictures before and after medical treatment. Should it be decided to try this we employ the same as that just given, but it will have to be more prolonged; it must last for many weeks.

This class of ulcer is more common in hospital than in private practice. Hospital patients cannot afford to be away from work long; surgery is easier in hospital than in private practice, a cure from surgery is quicker and probably more certain than that from medical treatment, and nowadays the risk to life from surgical treatment is slight; therefore surgical treatment, whilst often desirable anyhow, is especially so for hospital patients. Medical treatment should only be undertaken for those in whom the *x* rays do not indicate an ulcer—for they certainly fail to reveal it sometimes—or show that it is small. Its progress should be watched by *x*-ray examinations, say every six weeks; if it is not getting less the surgeon should be consulted, as he should if relapse has occurred after medical treatment, or this fails to benefit the symptoms.

In trying to decide when to adopt surgical treatment we must remember that even if the ulcer heals, pyloric stenosis, hour-glass stomach, or perigastric adhesions may follow, and all require surgery. Perforation, subphrenic abscess, fatal haemorrhage, or transition from a simple to a carcinomatous ulcer may occur at any time before complete healing, and when this has taken place dyspepsia may remain.

We shall hear from surgeons the best way of dealing with these ulcers. Gastro-enterostomy is by no means a universal panacea; it often fails to do any good. On the other hand, in suitable cases—for example, pyloric obstruction—the benefit is so great that if the patient is one for whom this operation is indicated it had better be done at once without waiting for medical treatment.

Perforation requires instant surgery. Every hour's delay greatly increases the risk of death. However severe the haemorrhage in the class of acute ulcer or ulcer A, no attempt should be made to stop it by surgery. Experience has shown that this increases very considerably the risk of death, for patients blanched by bleeding bear operations badly, and further, as the bleeding is often from many points, the surgeon cannot stop it; it can almost always be controlled by nature, aided by medical help. If in the second class haemorrhage is severe, operation should not be undertaken until the patient has recovered from its effects; then the question should be discussed whether it would be wiser for the surgeon to interfere with a view of preventing a second bleeding; certainly after the second he should operate to prevent a third. Even the rare event of death during a profuse haemorrhage should not make the physician blame himself for failing to ask surgical aid; had the patient been operated upon during the bleeding the chances of recovery would have been lessened.

The medical treatment for bleeding from a gastric ulcer of any sort is absolute rest in bed in a quiet room without visitors and enough morphine subcutaneously to keep the sufferer gently under its influence. If the symptoms show that the bleeding has been excessive, 2 grains of calcium chloride dissolved in water should be injected intramuscularly night and morning; this dose definitely increases the coagulability of the blood. No food or drink should be given by the mouth or otherwise for twenty-four hours. At the end of this time, dextrose in solution in tap water, 525 grains to the pint, may be given by the rectum, half a pint two or three times a day. But it is unlikely that all this can be retained and absorbed, so that the patient hardly gets more than 800 calories a day this way. Although this is by far the best rectal feed it is a starvation diet on which it is improbable that ulcers will heal; therefore, in two or three days at latest some milk in hourly or half-hourly feeds should be given by the mouth, and, as rapidly as possible, the patient should be got upon the diet already mentioned as suitable for ulcer of the stomach. I strongly suspect that, in the past, very many cases have been lost from too prolonged rectal feeding and consequent starvation; it must be remembered that these patients are often already under-nourished from severe dyspepsia at the time the bleeding occurs. Transfusion may be of great help, but, in the nature of things, it is unfortunately often impossible to carry it out at short notice.

Drugs given by the mouth are of very little value in arresting gastric bleeding; sometimes it has appeared that a drachm of liquor ferri perchloridi, with an equal amount of glycerin to facilitate swallowing, has stopped the haemorrhage, but as it is so nasty and may cause vomiting I have given up ordering it. During recovery iron is useful, but it should not be given as an acid preparation, nor in a form likely to cause indigestion. Reduced iron, the old-fashioned dialysed iron, or the saccharated carbonate may be prescribed. Levico water in doses of two fluid ounces suits some and the minute quantity of arsenic in it may be helpful.

The treatment of other results of gastric ulcer, such as organic hour-glass stomach, pyloric obstruction, subphrenic abscess, and perigastric adhesions, is entirely surgical. Of these, the treatment of the last is likely to be disappointing, for, although in some cases surgery is brilliantly successful, in others either the adhesions are such that the surgeon cannot give much relief or the patient, usually a woman, has become from long painful suffering such an introspective invalid that her symptoms continue after operation, and she wanders from operation to operation, each of which she welcomes. We should, before opening the abdomen of a woman for chronic disease, remember that a certain number drift into this very unsatisfactory class. I remember one in whom operation led to complete relief of her pain due to adhesions so long as she was in the hospital; when she returned to sympathetic friends the pain came back; we readmitted her to hospital and it soon disappeared. It is not a bad aphorism to say that we should think twice as long before opening the abdomen of a woman for chronic disease as we should before opening that of a man.

Opinions differ widely as to the frequency of syphilitic gastric ulcer; in this country it is regarded as very rare. If in a case of gastric ulcer there is evidence of syphilis, anti-syphilitic treatment should be given, but surgery may be required for some of the results of the ulcer, such as hour-glass contraction. Whether the treatment by intramuscular injection of calcium salts combined with giving parathyroid by the mouth, as suggested by Vines, will prove of benefit time will show. Here also it will be well to watch the progress by occasional *x*-ray examination of the ulcer.

To sum up. Statistics as to the result of medical treatment are at present valueless. Unless it perforates, the acute ulcer or ulcer A requires only medical treatment. Very many of the other class will derive most benefit from surgery, but ulcers should not be operated on when bleeding.

REFERENCES.

¹ *Lancet*, November 3rd, 1906. ² *Ibid.*, January 14th, 1922.

THE Academy of Military Medicine at Vienna has been converted into a museum for the history of medicine under the direction of Professor Max Neuburger.

THE fourth Italian Congress of Radiology will take place in Bologna on May 9th, 10th, 11th, 1922, at the Orthopaedic Institute Rizzoli, under the presidency of Professor Aristide Busi. In connexion with the Congress an exhibition of radiological apparatus, both Italian and foreign, will be held. Further information may be obtained from the Secretary of the Congress, Dr. Alberto Possati, Villa Verde, Bologna.