susceptible to the influence of digitalis has been produced. Whether that is brought about by the direct influence of the drug modifying the stimulus or acting upon the muscle cells, or indirectly through the vagus, is not clear, but is a subject which awaits further investigation.

It is rare to get this form of idio-ventricular rhythm so persistent. In most of my cases it was transitory. In one other case I got a similar effect, but my opportunities were not sufficient for full observation. Dr. John Parkinson tells me that in one of his cases where this rhythm was persistent digitalis, though given in full doses, was without effect.

The Output of the Ventricle in Partial Heart-Block.

Further evidence of the difference between the two kinds of stimuli is found in what happens in partial heart-block. Here the bundle is damaged, so that the fibres are reduced in number, and the stimulus has a difficulty in passing from auricle to ventricle. However extreme the damage may be, and however weak the stimulus which manages to get through, the resultant contraction of the ventricle is always full and complete. In Fig. 12, where there is, as a rule, but one contraction of the ventricle to two contractions of the auricle, the radial beat is large and effective. In one instance the stimulus of the second auricular beat after some difficulty (as shown by the wide *a-c* interval) did get through to the ventricle, and here an efficient beat follows.

The inference from these facts is that a stimulus weakened in passing through the conducting system, on reaching the ventricle, gives rise to an effective ventricular systole, in contrast to the stimulus arising in the contracting system, as in auricular fibrillation, where the stimulus may not provoke a full systole.

The Different Origins of the Stimuli.

The normal stimulus which arises in the sino-auricular node and passes by the auriculo-ventricular node and bundle to the ventricle is a stimulus of a peculiar kind which arises in cells evolved for this particular purpose. The stimulus which arises in auricular fibrillation is produced by cells which were not evolved for this purpose, the stimulus being merely incidental to the contraction of the muscle cells. It is not likely that the same character of stimulus would be produced by cells evolved for the special purpose and by cells evolved for a totally different purpose.

The stimulus which arises in the conducting system apart from the sino-auricular node—that is, in the auricle and auriculo-ventricular node and bundle—is of the same quality as that which arises in the sino-auricular node. We can therefore recognize two kinds of stimuli: (1) a stimulus A, which arises in the conducting system, and (2) a stimulus B, which arises in the contracting system.

Is the Difference in the Stimuli one of Quality or one of Strength?

Although it may seem that the difference between the two kinds of stimuli is one of strength, yet there are certain reasons which suggest that the difference is one of quality. We have seen, for instance, the peculiar effect of the stimulus on the contraction of the ventricle in auricular fibrillation. This peculiarity is more likely to be due to the quality of the stimulus than to its strength, for the presumably weak stimulus which reaches the ventricle in partial heart-block gives rise to an effective ventricular systole.

The suggestion that the quality of the stimulus is different when it arises in the Purkinje fibres in the ventricle from that when it arises in the muscle cells of the ventricle may account for some curious differences in other abnormal rhythms, as in auricular flutter, extra-systoles, and the idio-ventricular rhythm.

This presentation of the subject is of very real importance for the practice of medicine. The pulse rate and the character of the pulse are among the most instructive signs the doctor uses in estimating a patient's condition. The nature of pulse variations, even with the normal rhythm, is

still imperfectly understood, and, as a consequence, only a small part of the information which they are capable of yielding is available. In the foregoing description some light is thrown upon the variations in rate and character which are due to abnormal rhythms, and we get a little nearer to understanding the action of digitalis, in that a factor hitherto unsuspected has been revealed. This aspect of the subject raises other questions, such as the nature of the stimulus in other abnormal rhythms—auricular flutter, the extra-systole, the idio-ventricular rhythm, etc. It also involves the question of stimulus conduction by nerves and the Purkinje cells. Can they convey and propagate from cell to cell stimuli of different qualities? The muscle cell to muscle cell stimulation in auricular fibrillation raises the question whether such intercellular stimulation occurs in other organs.

Summary of the Difference in the Heart's Contraction arising in the Conducting and Contracting Systems.

1. The stimulus for the normal rhythm of the heart arises in tissues specially evolved for the purpose. The stimulus in auricular fibrillation arises in tissue specially evolved for contraction, the stimulus production being merely an incident in the contraction.

2. There is a remarkable difference between the rhythms which arise in the conducting and contracting systems of the heart, due to the difference in the stimuli. These differences are shown especially in the response of the ventricle. The ventricular systole in the normal rhythm, as well as in those rhythms arising in any part of the conducting system, is complete and effective.

3. The ventricular systole, when the stimulus arises in the muscle cells of the auricle, is often partial and ineffective, and when it arises in the muscle cells of the ventricle it is always ineffective.

4. When the contraction starts in the muscle cells of the auricle, as in auricular fibrillation, the stimulus which arises is much more sensitive to agents which favour or retard its passage to the ventricle.

5. The effect of digitalis is strikingly different, it having a far greater effect on auricular fibrillation and flutter and on the idio-ventricular rhythm than on any rhythm arising in the conducting system.

REFERENCES.

- ¹ BRITISH MEDICAL JOURNAL, March and April, 1905. ² Gibson, A. G.: On the Primitive Muscle Tissue of the Human Heart, BRITISH MEDICAL JOURNAL, 1909, Vol. 1, p. 149. ³ Holmes, A. Hillyard: The Auriculo-Ventricular Bundle in Mammals, *Journal of Anatomy*, Vol. 55, pt. 4, p. 269.

THE VALUE OF THE COMPLEMENT FIXATION TEST IN THE EXCLUSION OF ACTIVE PULMONARY TUBERCULOSIS.

BY

A. LISLE PUNCH, M.B.LOND., M.R.C.P.LOND.,

ASSISTANT PHYSICIAN, BROMPTON HOSPITAL FOR CONSUMPTION AND DISEASES OF THE CHEST; PHYSICIAN TO OUT-PATIENTS, ROYAL NORTHERN HOSPITAL;

AND

A. HOPE GOSSE, M.D.CANTAB., M.R.C.P.LOND.,

PHYSICIAN, BROMPTON HOSPITAL; PHYSICIAN TO OUT-PATIENTS, ST. MARY'S HOSPITAL.

A METHOD of performing the complement fixation test as a means of diagnosis in pulmonary tuberculosis has been published¹ by one of us (A. L. P.). By this method a positive result has been obtained in at least 98 per cent. of cases of definite pulmonary tuberculosis, while 140 cases selected as non-tuberculous controls gave a negative result in 137.

In all, the test has been performed in approximately 1,100 pulmonary cases or controls. Of these some 260 were definite cases of pulmonary tuberculosis, the criterion of a definite case being the undoubted presence of tubercle bacilli in the sputum. All save five of these gave a positive result. Of the five which gave a negative result three subsequently gave a positive. We do not propose here to enter into a discussion

FIG. 12.—Simultaneous tracings of the movements in the neck (*a*, *c*) and of the radial pulse. There are two auricular waves (*a*) to one carotid (*c*) and radial beat, except in one instance where two successive *a* waves are followed by two *c* waves, and the distance between *a* and *c* is much increased in the last of these two beats. In this case the impulse from the auricle is hindered in passing to the ventricle, so that only the impulse from every second beat of the auricle reaches the ventricle except in one instance when the passage of the stimulus is delayed, as shown by the wide *a-c* interval. It is to be noted that the resulting radial beat is large and full, as compared with the radial beats in auricular fibrillation in Fig. 7.

of the possible explanation of these negative results, but we may mention in passing that we have formed the opinion that a negative result may occasionally be obtained in very acute cases of pulmonary tuberculosis due to the shortness of the interval between the invasion of the body by the tubercle bacillus and the onset of symptoms. Except in these rare cases the results of the test would suggest that a positive reaction is obtained by the time the infection gives rise to symptoms.

Some 140 presumably healthy people or patients suffering from non-tuberculous diseases were taken as negative controls. Three gave a positive result. These three cases have been considered in a previous publication. It is sufficient to mention here that in one definite evidence of enlarged, presumably tuberculous, glands in the neck was subsequently discovered; in another, on more careful examination, it was impossible definitely to exclude a tuberculous lesion in the lungs; while in the third no evidence of tuberculosis could be discovered. Of the remaining 700 cases some 220 gave a positive result and some 480 gave a negative. These were doubtful cases clinically, so that no definite conclusion could be drawn from them. All that could be said at the time without later investigation was that there seemed to be a reasonable correspondence between the clinical condition on the one hand and the negative or positive result on the other.

In many of the cases which gave a positive result the accuracy of the test has been proved by the finding at a later date of tubercle bacilli for the first time in the sputum, but it is not our intention to deal in this paper with the complement fixation positive cases.

The proof that cases giving a negative result are free from active pulmonary tuberculosis is a more difficult matter. It is only by the lapse of time that the practical proof of this contention can be firmly established.

It would be a very difficult matter to follow up all 480 of these cases, nor is it necessary in order to estimate the value of the test to attempt it. What is true of 50 unselected cases can be taken as approximately true of 500. Some 250 of these negative cases were at the time examined clinically by one or the other of us. Many of them have been examined by us at a later date. We only know of two cases of our own in which the test proved to be unreliable by the finding at a later date of tubercle bacilli in the sputum. Of the remaining 220 cases under our colleagues or other medical men, we know of three in which the test proved to be wrong by the finding of tubercle bacilli in the sputum. These three cases, with the two just mentioned, are the five cases referred to above. One was Sir John Broadbent's case and two were Professor Langmead's cases. If there are any other cases amongst these 480 in which the test has been proved to be unreliable by the finding of tubercle bacilli in the sputum we are unaware of them; and we may mention that we have been able to follow their course in a large number. We exclude a series of 50 cases which have been dealt with fully in a previous publication.²

The ultimate practical proof of the entire freedom from active pulmonary tuberculosis is the lapse of time, and we have taken an unselected series of fifty of our own cases in which the test was performed in 1920 with a negative result, and where possible repeated it from ten to twenty months later. A similar result was obtained in all except one. These cases were all patients at the Brompton Hospital in whom for various reasons pulmonary tuberculosis had been suspected. As the result of the first test we formed the opinion, and acted on it, that there was no active pulmonary tuberculosis. When the test was repeated from ten to twenty months later the greatest care was taken to have all bloods numbered by a third person, and included in series with known tuberculous bloods and unknown bloods, and the clinical condition handed to him. Thus if there should have been any doubt as to unconscious bias in the reading at the first test the numbering of the bloods at the second and the similar results amply prove the absence of bias at either.

Notes of Cases.

The clinical notes, taken before the tests were done, have been extracted, and as given below must serve as the evidence that we were dealing, in all the twenty cases,* with suspicious cases of pulmonary tuberculosis.

* The authors originally forwarded fifty cases, reported as those here printed. At our request they have consented, in order to save space, to the reproduction of the first twenty only. The reports of all the fifty cases presented a great similarity, and the twenty here given are a fair sample.

No. 1.—E. D. Haemoptysis, a teaspoonful on several occasions in previous three months. Loss of weight; weight 6 st. 2 lb. Sputum negative. Complement fixation test negative on May 31st, 1920. *Re-examination*: Feels better; slight cough and sputum; no crepitations; weight 7 st. No haemoptysis lately. Complement fixation test negative on April 22nd, 1921.

No. 2.—J. F. Complaints of cough, sputum, night sweats; no crepitations; weight 9 st. 10 lb. Sputum said to have been positive once in 1919; negative since. Complement fixation test negative on June 4th, 1920. *Re-examination*: "Feeling fine." No cough, no sputum, no night sweats, no haemoptysis; no crepitations; weight 9 st. 10 lb. Complement fixation test negative on April 30th, 1921.

No. 3.—T. Haemoptysis streaks three weeks before; cough, sputum, and slight night sweats. Sputum three times negative. Dullness and crepitations at right base. Complement fixation test negative on June 16th, 1920. *Re-examination*: Cough, sputum, and dyspnoea, especially on exertion. Signs as before. Sputum negative. Diagnosis, fibrosis of right lung. Complement fixation test negative on May 6th, 1921.

No. 4.—I. B. Sent up for diagnosis. Pulmonary tuberculosis suspected. Pain in right chest; wasting. Pleurisy in February, 1920. No crepitations. Complement fixation test negative on June 18th, 1920. *Re-examination*: Feels quite fit; no symptoms. Complement fixation test negative on May 26th, 1921.

No. 5.—R. G. Cough; sputum copious; dyspnoea. Temperature 100.4°. Weight 9 st. 8 lb. Physical signs of cavity at right apex. Sputum repeatedly negative. Complement fixation test negative on June 18th, 1920. *Re-examination*: Symptoms and signs as before; weight 10 st. 2 lb. Sputum still negative. Diagnosis, apical bronchiectasis. Complement fixation test negative on May 20th, 1921.

No. 6.—R. Cough; pain in right chest; pleurisy right base. Father, sister, and brother all stated to have died of pulmonary tuberculosis in previous three years. No sputum available. Complement fixation test negative on June 2nd, 1920. *Re-examination*: No symptoms now; no crepitations. Complement fixation test negative on April 30th, 1921.

No. 7.—W. O. E. Haemoptysis half a pint in November, 1919. Cough; loss of weight. Been to a sanatorium. Colour in sputum fortnight before. No crepitations. Sputum negative. Weight 8 st. 8 lb. Complement fixation test negative on June 14th, 1920. *Re-examination*: Some cough and sputum; negative for tubercle bacilli. No further haemoptysis; no crepitations. Weight 8 st. 5 lb. Complement fixation test negative on May 1st, 1921.

No. 8.—J. J. T. Cough; pain in left chest. In military hospital 1916. "T.B. lung." Weight 9 st. 1 lb. Clubbing of fingers. Physical signs of unilateral fibrosis of left lung. Complement fixation test negative on October 30th, 1920. *Re-examination*: Physical signs as before. No tubercle bacilli in sputum. Weight 9 st. Complement fixation test negative on May 4th, 1921.

No. 9.—F. F. Cough; sputum. Temperature 99.6°; night sweats. Sputum negative. No crepitations. Complement fixation test negative on June 24th, 1920. *Re-examination*: Cough and morning sputum; no crepitations. No sputum sent for examination. Diagnosis, chronic bronchitis. Complement fixation test negative on May 16th, 1921.

No. 10.—H. V. Cough; sputum; coarse râles over lungs. Weight 9 st. 1 lb. Sputum stated to have been positive once—in July, 1919. Frequent attacks of asthma. Complement fixation test negative on May 26th, 1920. *Re-examination*: Has been under treatment since for asthma only. Chest dry on examination. Weight 9 st. 6½ lb. Sputum negative. Complement fixation test negative on May 26th, 1921.

No. 11.—A. K. No cough. Patient stated she had coughed up one pint of blood a month before. Complement fixation test negative on June 29th, 1920. *Re-examination*: While an in-patient was diagnosed as gastric ulcer. No chest symptoms. Complement fixation test negative on May 28th, 1921.

No. 12.—J. H. War pensioner. Cough; slight sputum. Tubercle bacilli reported once only in sputum, December, 1919. Sputum negative since. No crepitations; weight 8 st. 7 lb. Complement fixation test negative on August 5th, 1920. *Re-examination*: Feels well; no cough; no sputum. Weight 8 st. 7 lb. Complement fixation test negative on June 1st, 1921.

No. 13.—W. E. W. Weakness; cough. Streaks of haemoptysis two years before. Coarse râles in both lungs. Weight 8 st. 13 lb. Sputum negative. Complement fixation test negative on December 1st, 1920. *Re-examination*: Cough and sputum present; no tubercle bacilli in sputum. Coarse râles in chest. Weight 9 st. 7 lb. Complement fixation test negative on June 1st, 1921, and again on November 15th, 1921.

No. 14.—E. S. Cough; sputum. Haemoptysis in streaks in past three weeks. Temperature 99.4°. No crepitations. Complement fixation test negative on July 8th, 1920. *Re-examination*: No more haemoptysis; no crepitations. Complement fixation test negative on June 2nd, 1921.

No. 15.—E. B. Cough; sputum. Haemoptysis (slight) three weeks before. Temperature 99°; pulse 104. No crepitations. The complement fixation test was negative on August 19th, 1920. *Re-examination*: Patient replied fourteen months later: "I am glad to say I am keeping in the best of health, and have done so for a long time past now; had I felt in any way bad I should have come up before."

No. 16.—L., aged 20. Pain in chest; cough. Temperature 101° each night for three weeks. Signs of pleural effusion; confirmed by exploratory needle. Complement fixation test negative on November 2nd, 1920. *Post-mortem* examination six weeks after above test showed malignant growth in left lung and effusion into pleural cavity. Primary growth in ovary.

No. 17.—G. H., war pensioner. Cough; sputum; pain in chest. Stated he coughed up four pints of blood in April, 1920. Weight 8 st. 3 lb. Sputum negative. No crepitations. Complement fixation test negative on December 1st, 1920. *Re-examination*: Pain in abdomen after food. No crepitations. Weight 8 st. 4 lb. Sputum negative. Complement fixation test negative on October 5th, 1921.

No. 18.—S. Cough; sputum; pain in back. Debility six months. Sputum negative four times. No crepitations. Complement fixation test negative on July 14th, 1920. *Re-examination*: No fresh symptoms. Has been at work since last seen. Complement fixation test weakly positive on June 4th, 1921. Examination of sputum seven months later positive. Brother died of pulmonary tuberculosis in October, 1921. Contact. Complement fixation test strongly positive on February 11th, 1922.

No. 19.—E. W. Cough; sputum. Haemoptysis (streaks) in May, 1920. Loss of weight; feels ill. Temperature 99.6°. Weight 8 st. 5 lb. Sputum negative. Complement fixation test negative on September 27th, 1920. *Re-examination*: Feels well; no cough. Weight 8 st. 5 lb. Temperature 98°. No crepitations. Complement fixation test negative on November 8th, 1921.

No. 20.—T. P. Cough; sputum; dyspnoea. Weight 8 st. 4 lb. No crepitations. Sputum negative. Complement fixation test negative on September 24th, 1920. *Re-examination*: Feels better; still has dyspnoea. Weight 9 st. 3 lb. No crepitations. Complement fixation test negative on April 22nd, 1921.

Comments.

A study of the above cases will show that the clinical condition at the time of the first test justified a strong suspicion of pulmonary tuberculosis. In all of them some combination of cough, sputum, fatigue, loss of weight, pyrexia, haemoptysis, and pain in the chest was present—in fact all the common symptoms of pulmonary tuberculosis. The frequency of haemoptysis is a striking feature.

A study of the re-examination notes will show an absence of any further evidence of the disease some ten to twenty months later. The too frequent absence of sputum reports when the patients were re-examined is due to the absence of any sputum. In no case did apical crepitations appear at the later examination, and almost all cases were examined by post-graduates as well as the writers. No. 18 had a weak positive complement fixation test at the re-examination, and the sputum was found positive seven months after the positive complement fixation test; but he had been a "contact" till four months after the second test was taken when his brother died, so the infection may have occurred between the first and second tests. Many patients had regained their normal health, some had improved, and others were unchanged. Two of the fifty cases died, and *post-mortem* examinations showed malignant growths of the lung or pleura as the cause of death. After the lapse of ten to twenty months or even longer forty-nine had not developed any definite evidence of pulmonary tuberculosis, nor, with a few exceptions, were they even under treatment.

It is the absence of the development of any more definite evidence of tuberculous infection in all but one of these suspicious cases in the course of ten to twenty months afterwards that would appear to us to offer striking evidence in favour of the accuracy of the test and to demonstrate its value in the exclusion of active pulmonary tuberculosis.

We wish to thank the Lady Almoner of the Brompton Hospital for her valuable help in tracing the patients.

The laboratory work has been done in the Institute of Pathology and Research, St. Mary's Hospital, under a grant from the Medical Research Council.

REFERENCES.

¹ Punch, A. Lisle, *Lancet*, vol. ii, 1920, p. 647. ² *Ibid.*, vol. ii, 1921, p. 497.

At a meeting of the Manchester Pathological Society on February 8th, with Mr. Howson Ray in the chair, Dr. J. A. Murray, Director of the Cancer Research Fund, demonstrated the pathological anatomy of experimentally induced squamous carcinoma of the skin of the mouse following repeated and long-continued application of coal tar. Lantern slides were shown illustrating the growth of the tumours, their metastases and dissemination, and the invasion of the subjacent tissues. Reference was made to the application of autoplasmic implantation of tumour fragments to bring out the powers of growth of the altered epithelium when removed from the direct irritating influence of the tar and of bacterial infection. By this method of artificial metastasis it was possible to demonstrate the malignancy of growths at a very early period. The relative advantages were discussed of other successful methods of inducing cancerous changes.

D

EVIDENCE OF PANCREATIC DISORDER IN RICKETS.*

BY

E. C. DODDS, M.R.C.S., L.R.C.P.,

CHEMICAL PATHOLOGIST AND LECTURER IN BIOCHEMISTRY, MIDDLESEX HOSPITAL; PATHOLOGIST TO THE ROYAL NATIONAL ORTHOPAEDIC HOSPITAL.

(From the Bland-Sutton Institute of Pathology.)

MANY observers have been working on the factors concerned in the production of rickets, and recently special prominence has been given to the view that the problem is intimately bound up with the question of diet, more especially with the amount and quality of the fats. It is urged also that carbohydrates play an important part, but as to the explanation of these two facts there is still considerable difference of opinion.

The well-known vitamin deficiency theory ascribes the production of rickets to a lack of the vitamin A, which is fat-soluble; but with regard to the carbohydrate part of the problem, no observer seems to have given a satisfactory explanation. In this communication the importance of both fats and carbohydrates is taken into account, but the conclusions to be drawn from the investigations made point in a different direction. As a result of the examination of cases of acute rickets in children, distinct evidence was found of pancreatic disorder, as determined by an increase (1) in urinary diastase, and (2) in faecal fat content.

1. The Diastatic Power of the Urine.

By means of its diastatic enzyme normal urine is able to hydrolyse starch solution to a mixture of products which give no colour with iodine solution. Wohlgemuth expressed this property of urine quantitatively by means of "units," the number of units of diastase being given by the number of cubic centimetres of 0.1 per cent. starch solution digested by 1 c.cm. of urine.

Method.

The usual method is to put up a series of test tubes, each being filled as follows:

Test tubes:	1.	2.	3.	4.	...	12.
Starch 0.1%	2 c.cm.	2 c.cm.	2 c.cm.	2 c.cm.	...	2 c.cm.
Urine ...	0.3 c.cm.	0.2 c.cm.	0.1 c.cm.	0.09 c.cm.	...	0.01 c.cm.
Saline...	0.7 c.cm.	0.8 c.cm.	0.9 c.cm.	1 c.cm.	...	1 c.cm.

The starch solution is put in first, then the urine, and the saline is run in last, in order to wash the first two down to the bottom of the tube.

The tubes are incubated in a water bath at 37° C. for half an hour, are then cooled under tap water, and enough N/50 iodine solution to produce a permanent colour is added to each tube. The tube immediately preceding one with a mauve tint is taken as that in which the 2 c.cm. of starch solution is just digested by the urine. Suppose 0.1 c.cm. of urine digests 2 c.cm. of starch solution, then 1 c.cm. of urine would digest 20 c.cm. of starch solution—that is, there would be 20 units of diastase present.

Towards the end of this series of diastase reactions, I modified the original method so as to take into account the hydrogen-ion concentration of the urine. The diastatic enzyme works best at a hydrogen-ion concentration of pH 6.1 (Michaelis), and since the pH of the urine varies normally between pH 4.5 to pH 7 it is obvious that unless the reaction of the urine is pH 6.1, the diastatic power will be decreased. The newer technique is to bring the urine's hydrogen-ion concentration to approximately pH 6.1 by diluting it with a buffer solution of phosphates and then determining the diastase as in the ordinary method. A description of this method will be given elsewhere, but it should be pointed out that it can only increase a low reading of the ordinary method, and can never decrease it.

The normal number of diastatic units in urine lies between 6 and 30. Any increase is said to indicate pancreatitis, with the possible exception of cases of eclampsia, in which the diastatic power is said to be high.

In Table I are given the figures from a series of cases of acute rickets; all are high enough to warrant a diagnosis of pancreatitis. The mean of 17 cases is 154 diastatic units.

These results were controlled by the examination of a series of normal children of about the same age as the rickety cases; as will be seen from Table III, the results were all low. As some observers maintain that there is a general infection in rickets, series of septic cases showing no rickets, and with and without fever, were examined (Table III). It was also thought advisable to make control observations on a series of cases where there was bone erosion (Table III).

* A report to the Medical Research Council.