A Lecture to Graduates

ON

THREE YEARS OF PELVIC SURGERY.

DELIVERED AT THE MANCHESTER ROYAL INFIRMARY,

BY

W. E. FOTHERGILL, M.A., M.D., C.M.,

PROFESSOR OF OBSTETRICS AND GYNAECOLOGY, VICTORIA UNIVERSITY
OF MANCHESTER.

DR. C. P. BRETNALL, Resident Surgical Officer to the St. Mary's Hospitals, Manchester, and Dr. W. E. Powell, Senior House-Surgeon, Special Departments, Manchester Royal Infirmary, have been kind enough to enumerate and analyse the cases on which I have operated personally at the St. Mary's Hospitals and at the Manchester Royal Infirmary during the last three years. Their figures deal with 1,354 consecutive operations, and show the relative frequency with which common pelvic lesions occur, and indicate the measure of success which may be expected when these lesions are dealt with by surgical methods. The figures also bring out certain debatable opinions and illustrate certain points of view which form interesting subjects for discussion. I thank my two colleagues for the time and care they have bestowed on this analysis, which I propose to bring before you with brief comments on the methods used and the results secured.

Pelvic Infections.

During the three years there were 82 cases of pelvic infection. Of these, 5 were operated upon during the acute phase by the vaginal route, drainage being secured through the floor of the pouch of Douglas. In the other 77 cases the abdomen was opened after waiting until the pulse and temperature had been normal for some time. In 35 of these 77 cases the body of the uterus was removed together with both appendages, while in the remaining 42 cases the injured organs alone were removed, the uterus, one ovary, and one tube being conserved. Drainage by the lower angle of the abdominal incision was used in 13 of the 77 cases. There were no deaths amongst these 82 cases of severe pelvic infection, nor were there any unpleasant sequelae, such as the formation of faecal fistulae. This success I attribute to two factors. First, the results of infection have a natural tendency to recovery. They are not progressive diseases like new growths. Inflammation is not a disease, but a curative process. If judiciously helped by the surgeon it will generally save the patient, though it may be at the cost of her reproductive function. Secondly, in these cases the abdomen was only opened after the acute inflammation was over, the vaginal route being chosen whenever it was really necessary to let out pus during the continuance of constitutional disturbance. Some of you may be horrified to hear that the uterus and both appendages were removed in 35 out of 77 cases. But a uterus without ovaries is no use, and, if left isolated in the middle of the pelvis, with its blood supply impaired, it often becomes a source of trouble. It is best to remove everything except the cervix unless it is possible to leave a fairly sound uterus, one fairly sound ovary, and one fairly sound tube—in short, a working set of reproductive organs. The 35 patients that have suffered the radical operation have their menopause behind them, and will be healthy and comfortable as to the pelvis for the rest of their lives. As to the 42 other patients, some of them may certainly have children; but, on the other hand, some of them are sure to have recurrences of infective trouble with disability, discomfort, and further surgical intervention.

Pelvic Growths.

In 200 cases "fibroids" were the leading feature. One tumour proved to be a sarcoma and one an adenomyoma. Cancer of the cervix complicated one case and cancer of the body another; while small ovarian cysts were present in 13 cases. Nine of the myomata were cervical in position, many of them were degenerated, and several were infected, one patient having general peritonitis on admission. One sloughing myoma was removed in small pieces by the vaginal route, the abdomen being opened in the other 199 cases. Panhysterectomy was done 10 times, supravaginal hysterectomy 180 times, myomectomy 7 times; while in two cases the growths could not be removed. In these 200 cases there were six deaths:

1. General peritonitis on admission; drainage only.
2. Ill after five years' bleeding; abdomen reopened for partial obstruction; no cause found.
3. Fibroid in broad ligament; shock; died the same day.
4. Extreme anaemia; recovered from operation; died some weeks later from anaemia.
5. Embolism tenth day.
6. Uterus contained a dead ovum; pelvic sinuses thrombosed; embolism third day.

Panhysterectomy was done only when the myoma was cervical and in the case complicated by cancer of the cervix. I regard the occurrence of cervical cancer after supravaginal hysterectomy as a mere coincidence. If the cervix should be removed because a woman has a myoma, surely several other parts of her body which are prone to cancerous change should also be taken for the same reason. In operating for myoma I generally follow the guiding principle of leaving nothing but the cervix, unless it is possible to leave a working set of reproductive organs. It is difficult to leave ovaries with their blood supply unimpaired. There is evidence that ovaries left *in situ* do not persist more than a couple of years, and that they often become the site of pathological changes.¹ A woman can only have one menopause, and I prefer that she should get it over just after her operation rather than two years or so later. The surgical menopause is regarded with undue apprehension, and J. W. Bride's² investigations show that there is nothing solid to be gained by leaving ovaries when the uterus is removed.

The present series contains only 25 cases of cancer of the cervix, as during the last three years I have been referring a great many of these cases to Dr. A. Burrows of the Manchester Radium Institute. Of the 25 cases 12 were inoperable and received palliative treatment by curette and cautery. The other 13 cases were dealt with by abdominal total hysterectomy. There were no deaths and no unpleasant sequelae. The cases were carefully selected, and we hope that some of them will pass the test of freedom from recurrence for five years. There were 23 cases of cancer of the body. All of them were attacked by the abdominal route, and satisfactory total hysterectomy was completed in 21 cases. The prognosis is five times as good as in cancer of the cervix. There were three deaths:

1. From exhaustion, fourth day.
2. From paralytic ileus, third day.
3. Radium put in parametria at time of incomplete operation; healing, but died from exhaustion on the eighteenth day.

There were 14 patients who were found on abdominal section to be the subjects of ovarian cancer, primary so far as could be ascertained. Most of these patients had a quantity of free fluid in the abdominal cavity (hydroperitoneum). Removal of the growths was attempted in each case, but was found to be impossible in five instances. Of these 14 patients, 5 died of exhaustion before leaving hospital. It may be asked if operation in these cases is justified. There are three reasons for it. First, the removal of a malignant ovary often gives considerable relief during the remainder of the patient's life. Second, in certain cases, in spite of a laboratory diagnosis of malignancy, there is no recurrence at all. Third, without opening the abdomen it is often impossible to make a complete diagnosis. For a uterine myoma, a fibroma of the ovary, or an ovarian cyst, if its blood supply is impaired, may bring about the appearance of free fluid in the peritoneal cavity. Cases with hydroperitoneum, therefore, should not be tapped, but should be submitted to exploratory incision. If this is always done a considerable percentage of the patients can be saved.

Simple cystic and solid growths of the ovary and broad ligament amounted to 114 cases. One of these had general peritonitis on admission and was treated by drainage only. The growths alone were removed in 97 cases, both appendages together with the body of the uterus being removed in 16 cases. Here, as in other conditions, the rule was followed of leaving nothing but the cervix, unless a working set of reproductive organs could be retained. There were four deaths after the 114 operations:

1. Dermoid cyst and fibroid; adhesions; abdomen reopened fourth day for signs of obstruction; no cause found; died the same day.
2. Patient aged 80; admitted for intestinal obstruction; healed well after removal of large cyst; died twenty-third day, intestinal paresis.
3. Two solid ovarians; bronchopneumonia.
4. Suppurating ovarian with peritonitis on admission; drainage only.

Other Pelvic Disorders.

There were 59 cases of retroversion considered to require operative treatment, and dealt with by means of Webster's sling. I find this to be the most satisfactory of the 87 operations which have been devised in recent years for the correction of retroversion. Many of the cases were complicated by the results of old pelvic infection, and in several dyspareunia due to prolapsed ovaries was the leading indication for treatment.

Uncontrollable bleeding at the menopause was treated by supravaginal hysterectomy in 10 cases, which, I am given to understand, could have been dealt with successfully by means of radium. There were also 13 cases of ectopic gestation requiring salpingectomy, and remarkable only because they included no example of rupture with haemorrhage demanding "urgency" operation.

Further, there were 23 unclassified abdominal sections whose consideration is outside our present purpose, interesting as many of them were. They included an inverted uterus, a double uterus, a sequestered ovarian adenoma implanted on the rectum, diverticulitis, a renal sarcoma adherent to an inflammatory mass in the pelvis, a cyst of the common bile duct, desmoid growths of the abdominal wall, various cases of hydroperitoneum of obscure origin, and various cases of tuberculous peritonitis involving the genital organs. There was no case of primary genital tuberculosis.

Review of Abdominal Operations.

Thus there were 545 abdominal sections in all, and the deaths were 18 in all—that is, 3.3 per cent. It is clear that the mortality could be kept lower than this by selecting the cases; but if operation is undertaken whenever it is thought that life may be prolonged or that suffering may be relieved, a number of post-operative deaths must occur. A careful study of the notes of the 18 patients who died brings out a striking fact which is not always kept in mind. Only three of the eighteen were in ordinary good health when they came for treatment. The other fifteen had been injured by prolonged bleeding, by infections and degenerations, or were the subjects of malignant disease. The three healthy persons died from shock, embolism, and pneumonia respectively. In the other fifteen, pathological factors had been at work before treatment was sought. The lesson is that the removal of lumps from the pelvis and abdomen is like making a will—it should be done in time of health. Three deaths in 545 is $5\frac{1}{2}$ in 1,000, and we know that four deaths occur in 1,000 cases of labour. Thus, when a woman is in good health, she can have a simple uterine or ovarian tumour removed with little more risk and with much less discomfort than that involved in having a baby. But with delay up go the risks. No pelvic lump can be guaranteed to remain in a healthy condition; and if the patient declines to part with one in time of health the responsibility should rest on her and not on her medical adviser.

Vaginal Operations.

Passing on to the cases in which vaginal operations were done, there were 406 cases of genital prolapse. These were treated by methods which I have sufficiently described elsewhere, and all left hospital soundly healed. F. H. Lacey's recent investigation³ of my 1914-15-16 cases of prolapse shows that permanent cure was obtained in 97 per cent. of them. There were also 112 cases of rupture of the perineum and recto-vaginal fistula, who were all discharged cured. The urinary fistulae numbered 16 and were cured with one exception, which requires further treatment. The whole vulva was excised seven times for cancer and once for leucoplakia, and these operations, with those previously mentioned, make up 560 classified vaginal operations.

There were also 249 smaller vaginal interventions, including cases of haematocolpos, septate vagina, unruptured hymen, senile endometritis, pyometra, cysts of the vulva, Bartholinitis, caruncles, polypi, sterility, spasmodic dysmenorrhoea, cervical lacerations, hypertrophied clitoris, retained products of conception, and various diagnostic explorations of the uterine cavity. There was no death after any of the 809 vaginal operations; which, added to the 545 abdominal sections, give the total of 1,354 consecutive operations in 1919, 1920, and 1921.

REFERENCES.

¹ Polak: *Amer. Journ. Obstet.*, 1918, pp. 199 and 349. ² Bride: *Journ. Obstet. and Gyn. British Empire*, 1922, vol. xxix, No. 1. ³ Lacey, *ibid.*, 1921, vol. xxviii, No. 2, p. 251.

SOME COMMON DEFECTS OF DIET AND THEIR PATHOLOGICAL SIGNIFICANCE.

ABSTRACT OF THE OLIVER-SHARPEY LECTURES DELIVERED BEFORE THE ROYAL COLLEGE OF PHYSICIANS,

BY

E. MELLANBY, M.D.,

PROFESSOR OF PHARMACOLOGY, UNIVERSITY OF SHEFFIELD.

LECTURE II.

IN his second Oliver-Sharpey lecture, delivered on May 4th, Professor E. Mellanby began by reminding his audience of the conclusions to which his first lecture had pointed. He had stated that foodstuffs could be divided into two groups. The larger group, which included the major part of the average diet, was practically devoid of sodium, calcium, chlorine, and two vitamins—namely, the fat-soluble A and the antiscorbutic factor. The reason why this was the general diet was, because modern urban communities lived at a distance from the areas where food was produced, and therefore preference had to be given to foods for popular consumption which were easily transportable and which had good keeping properties. The economic situation made it necessary also that foods should be cheap and abundantly produced. He had also pointed out that taste was an uncertain factor in deciding the relative value of foods, and did not prevent people from eating the wrong thing. Certain foods which seemed to be good, such as cereals, upon which children flourished and grew fat, might, under certain conditions of ill balance in other respects, give a deceptive impression of being more nutritious and satisfying than they really were. He had then proceeded to consider, what would be the main theme of this second lecture, the factors in diet which were responsible for the calcification of teeth and bones.

Professor Mellanby went on to say that in all his investigations upon deficiency diseases he had been impressed by the fact that as soon as there was any interference with a natural food product it became less suitable for food and lost some of its best properties. He took as an example the case of milk: if milk was allowed to go sour and acid caseinogen to be precipitated the calcium was lost, so that acid caseinogen used in a diet already deficient in fat-soluble vitamin made the resulting rickets much worse.

Calcium and Fats.

Even if butter were added to a diet a certain additional quantity of calcium must be added to obtain the maximum physiological value. Butter was less satisfactory from the point of view of rickets prevention than the milk from which the butter was made, probably because, in the course of separation, the substance lost some of its calcium. Butter therefore was not as good a food as one would expect, and the results could not be improved in the case of an ill-balanced diet by giving larger and larger quantities of butter. The more butter that was given the worse under these circumstances did the animal become, and the animal might be made very ill by being given much butter without a certain amount of calcium to balance the diet.

This fact had been worked out by McCollum, of Johns Hopkins University, who had tested on rats the value of a diet as affecting growth, fertility, and the power of the mother to rear her offspring. McCollum had shown that the conditions could not be made better merely by increasing the fat. He had added first of all a mere trace (0.1 per cent.) of calcium carbonate, and the rats did better in growth, though the second generation of young rats did not thrive at all. But on doubling the percentage of calcium carbonate, not only was the growth of the adult rats much improved, but the second generation also began to flourish. Cod-liver oil had the power of utilizing to the full every bit of calcium present in the diet, and with 2 per cent. of cod-liver oil added to a diet in which there was 0.1 per cent. of calcium carbonate the condition of the animal was much better than with 8, 10, or even 20 per cent. of butter, with the same minimum of calcium, and no cod-liver oil. If, however, the amount of calcium was raised to 0.5 per cent. butter under those conditions became as good as cod-liver oil. The cod-liver oil could use the minutest quantities of calcium, which the butter could not. McCollum's explanation of these facts was rather hesitating, but the simplest view to take was that cod-liver oil contained a large amount of the fat-soluble vitamin and butter only a small amount.

The lecturer then passed on to speak of the work of Zilva and Miura in the biochemical department of the Lister