CANCER OF THE BREAST.

TREATMENT OF THE PROEMIAL BREAST.*

BY

SIR G. LENTHAL CHEATLE, K.C.B., C.V.O., F.R.C.S.,
SURGEON AND LECTURER IN SURGERY, KING'S COLLEGE HOSPITAL.

[With Special Plate.]

No surgeon of experience can be satisfied with the results obtained by the present surgical treatment of cancer when it arises primarily in the breast.

The object of this paper is to direct attention to the existence of a potentially dangerous state of the breast, and to indicate the advisability of removal of a breast in this state rather than wait for the evidence of disease for which surgical measures have been proved to be so unsatisfactory. The consideration of the following facts explains why surgical measures are so often doomed to failure when primary cancer is clinically recognizable as such in the breast.

First. It is an aphorism that cancer of the breast must have developed considerably before it has attracted attention to its presence. By the time it has made itself manifest it too often has spread beyond the limit of the most extensive and skilful operation.

Second. The dilemma is really worse than this bald statement makes it appear, for the patient is usually the first to discover the indications of a tumour. Unfortunately her next step too often depends upon the relative importance she attaches to their warnings. When she regards them as relatively unimportant there will be delay in receiving the treatment which should be adopted at the earliest possible stage.

Third. While all are familiar with examples of complete surgical successes in cancer of the breast, we also know that the number of complete surgical failures is too great to justify an attitude of passive contentment with the present treatment. Moreover, the successes claimed within a few years of the operations have often proved illusive. Operative measures upon cancer of the breast have been followed by complete quiescence for ten and twelve years. Success assumed on a post-operative limit of quiescence for five years or so must be regarded as too optimistic.

Fourth. In some breasts which have been excised because they were considered to be in the proemial state I have found I was wrong and that they contained tumours. Upon careful microscopical examination of whole sections of separate breasts I discovered in them tumours of the following kinds: (1) simple papillomata, (2) papillomata the innocence of which is exceedingly doubtful, (3) malignant papillomata, and (4) one with a small cancer which was infiltrating fat outside the gland. In addition to my own patients I must refer to two others. Mr. Sampson Handley kindly sent me a breast for microscopical examination. I had seen the case clinically, and then I regarded it in the proemial state. In one of the large ducts I found a papilloma the benign nature of which is open to considerable doubt. My colleague, Mr. Burghard, who has given me great encouragement by showing his interest in my work, also kindly sent me a breast which I should have described upon clinical examination as also being in the proemial state. It was only by very careful microscopical examination of whole sections that we discovered a small cancer within the breast and infiltrating the fat outside the gland, also an apparently simple papilloma in a duct in another part of the gland. In all these patients clinical signs of tumours were absent.

Taking into consideration the whole position, it would be of great surgical importance could a line of treatment be indicated that would deal adequately with breasts in which cancer or papillomata were likely to occur, and not, as we do now, devote our attention to those breasts in which these diseases are definitely present. An absolute indication of this line of treatment would be of basic importance, and its establishment as a regular surgical procedure would be a great step in the direction of safety.

I hope to prove that there is a condition of the breast which is a prelude to those particular pathological changes to which I have alluded. I name it the "proemial breast."

* The term "proemial breast" is employed to indicate a condition of the breast which makes it a prelude to later developments of simple papillomata and malignant papillomata and other forms of cancer. It is regarded as more accurate than the terms "precancerous," "precursory," etc., which might otherwise be employed.

I am so fully convinced that I have already begun treating breasts on this assumption. Careful microscopical examination revealed in some of them the early stages of the diseases I had hoped to be in time to prevent. I take these facts as offering remarkable evidence in my favour and supporting my belief that there is such a thing as a proemial breast.

Before I describe the characteristic features of the proemial breast it is essential to clear the ground in one important respect. It is a fallacy to maintain that every breast that appears nodular on palpation is in a pathological state. Provided there be not too great a layer of superjacent fat most adult female breasts are multinodular on palpation. The multiple nodules are mainly prominent parts of mammary ducts, and can be recognized as such by careful observation. Palpation should be continued in the normal directions of the ducts, of which the "nodule" is only a part, bearing in mind that a duct becomes smaller in calibre the nearer it is to the periphery of the gland. The examination is conducted best by supporting the pendent breast with one hand and palpating it with the fingers of the other. Also it must be borne in mind that the vascular turgescence of puberty, menstruation, and pregnancy increases the size of the glandular structures and may render them painful. It is possible that physiological turgescence may occur from other causes. Therefore I attach importance to the facts that in all such conditions the whole of the gland, and not only a part of it, would be taking part in the disturbance, and that the multiple nodules would be of the same size when corresponding parts of the same breast were compared. To make sure that the condition be physiological it is essential to examine the breast when the cause of the turgescence has passed.

Ducts are more accessible to palpation than the acini because they are more rigid in structure and towards the nipple more superficial in position.

THE PROEMIAL BREAST.

The proemial breast is worthy of due recognition in future pathological and clinical considerations for two reasons: (1) It is in a state that makes it a prelude to further pathological changes of a highly important character—mainly simple papillomata, malignant papillomata, and other forms of cancer. (2) Its clinical recognition and adequate treatment occasionally enables the surgeon to recognize (microscopically) the presence of one or more of these changes, before any clinical sign of their existence is obtainable by any other means—that is to say, when the proemial breast has merged into those profoundly important stages to which it has acted as a prelude. When once established the condition of the proemial breast is permanent. To describe next its clinical and pathological appearances, and its treatment:

Clinical.

The patient, who has generally reached the age of from 40 to 45 years, complains of pains in the breast. Occasionally it is so severe and has lasted so long that in spite of all kinds of treatment she can bear it no longer, and insists on having the breast removed. Generally the pain is not so severe and is of an aching character which is not always affected by menstruation. On questioning the patient, it is found that the pain began sometimes as early as 30 years of age—more often it commenced somewhat later; also that it may have lasted some six months or one, two, three, or more years. The pain is continuous. On examination both breasts may appear multinodular. But while the nodules of the unaffected breast correspond to the signs I have described as indicating a normal breast, the nodules of the diseased breast are swollen, painful and tender on palpation, and are not of the same size in corresponding parts of the same gland. Sometimes a nodule at the periphery is bigger than one situated nearer the nipple. Careful palpation reveals the fact that the swollen nodules are mainly composed of prominent parts of tortuous ducts. The state of affairs may be limited to a segment of the breast, or two segments, one of which may be on the opposite side of the same breast, of which the intervening parts are normal. Or the whole gland may be affected by the same disturbance. There may be an intermittent or continuous discharge of serum from the nipple. A thin layer of coagulated serum may be seen on the surface of the nipple. The discharge is always small in amount.

In the later stages of this complaint the pain disappears and cysts are present in varying number. Massage, rather severe in its application, may induce a thick white discharge from the nipples and constantly from the same duct orifices.

The fluid contains desquamated epithelial cells. It is necessary to caution surgeons not to massage the breast to elicit this information, since it must be borne in mind that it is possible that a breast diagnosed as being in the proemial state may contain a small undiagnosable cancer, and there would be risk of disseminating cancer cells under these circumstances.

Pathology.

In the early stages there is to be observed a more or less diffuse desquamative hyperplasia of the epithelium in one duct, or two or more ducts, and also in some of the acini which are in direct connexion with them. The desquamative hyperplasia is most obvious in the smaller ducts (see Fig. 1) and acini. Ducts may be affected alone and acini may also show epithelial activity where none exists in the corresponding duct. The consequence of an abnormal collection of their contents leads to distension of the ducts and acini and accounts for the immediate results of this process—namely, the formation of cysts and stagnation of their contents (see Fig. 2).^{*} Small cysts in the process of formation contain the active desquamative hyperplasia of epithelium. The large cysts are lined by a degenerated atrophied epithelium which at parts has been shed.

The connective tissue around the affected glandular elements undergoes progressive secondary change and the connective tissue cells in parts begin to multiply (see Fig. 1). Also small accumulations of lymphocytes may occur at isolated and infrequent intervals, and can be seen immediately outside some parts of the ducts and acini. In the later stages these accumulations may be more pronounced and occupy longer tracts of ducts (see Fig. 2). I repeat my belief that the proemial breasts act as menaces by being preludes to and by encouraging grave changes within them, and this contention receives support from the fact that in what I estimated and diagnosed as being only proemial breasts I also found intracystic growths of simple papillomata, of malignant papillomata, and other forms of early cancer.

Treatment.

By making particular allusion to multiple small nodules occurring in normal breasts I have at once excluded many glands that hitherto have led to misconception in many minds, and it should not lead to the assumption that both glands are in the proemial state in the same individual, although it may be so.

From what I have said concerning the proemial breast it must be concluded that I regard it with supreme importance and distrust. In saying this I do not suggest that at its best it is certainly doomed to lead to grave pathological changes, but I hold that it is in a condition which is entirely favourable for their later genesis, while, on the other hand, in alarming frequency such changes have unexpectedly been found already to be present. I look upon it and treat it in precisely the same way and for the same reasons that compel surgeons to remove an appendix that has been inflamed. The proemial breast bears exactly the same relationship to cancer and papillomata of the breast as the proemial appendix does to general peritonitis. Cancer of the breast and general peritonitis are just as far beyond surgical control now as they always have been. General peritonitis is prevented by early surgical measures. My endeavour is to bring the preventive treatment of cancer of the breast similarly into line. Preventive surgery in relation to cancer elsewhere has been adopted. Take, for instance, Sir Henry Butlin's and Sir A. Mayo-Robson's work on leucoplakia of the tongue, and the latter's work and that of Sir Berkeley Moynihan on the gall bladder and on gastric ulcer. Consciously and unconsciously surgeons are acting on the same principle when they remove early those prostate glands which are interfering with the normal passage of urine.

My treatment of the proemial breast, in whatever stage it exists, is to remove it in the following way: I make one long curved transverse incision below the nipple, with its convexity downwards. Its length is equal to the limits of the breast; the outer end of the incision is continued into the axilla. Upper and lower flaps of skin are dissected from the limits of the breast and the axilla is opened. The nipple is reflected with the upper flap; it is not removed, care being taken to separate it from the parts immediately beneath it. In so doing it is wise to remove any white fluid that may appear at its base or from the surface from which it has been cut;

it might be infective. The breast and fascia over the pectoralis major are dissected away *en bloc*, with the lymphatic glands attached. It is essential that the removed gland should be subjected to a most careful microscopical examination for fear the breast might have passed from the proemial stage to changes of gravest issue. Should the presence of such a disaster be discovered, it is necessary then to remove the skin formerly covering the breast, pectoralis major and minor, and more complete dissection around the higher parts in relation to the axillary vein.

My hope and belief is that in the majority of proemial breasts cancer, at all events, will not be found to be present. If, however, it should be discovered, it will be a great advantage to be thus warned of the necessity of a complete operation at a much earlier date than would otherwise have been possible.

I publish the following reports of four breasts presumed by me to be in the proemial state before operation: and the microscopical appearances I found in three of them, which proved I was wrong in that presumption.

CASE I.

Female, aged 45 years, single. She is waiting for operation.

Left breast: Small gland, nipple not retracted, no discharge from nipple. In an upper and outer segment of breast there is a fluctuating swelling, not adherent to the skin, with defined margins; situated immediately above it and covering its upper margin is a longitudinal scar made at an operation performed elsewhere five years ago for the excision of a cyst. The rest of the gland appears normal. There are no enlarged glands in axilla.

Right breast: Small gland, nipple not retracted, no discharge from nipple. In an upper and outer segment of the breast there is a fluctuating swelling, not adherent to skin, with defined margins. Immediately surrounding it are multiple nodules which can be distinguished as swollen, tender, and painful parts of tortuous ducts. The nodules are irregular in size and larger than normal. There are no enlarged lymphatic glands in axilla.

The left breast probably contains a large cyst and possibly several small ones. The right breast is in a more serious state; the precise condition can only be revealed by careful microscopical examination of whole sections of this breast. I will report later what is discovered on that examination.

CASE II.

Female, aged 43, married, one child 8 years old, which she nursed.

Right breast: Nipple not retracted, no discharge from nipple; the gland is multinodular in all parts, most marked in segments on outer and inner areas of breast. The nodules are of irregular size, larger than normal, and are painful and tender on palpation; menstruation makes no difference to pain. The pain, which has been borne for eight months, has not been improved by rest, hot fomentations, and arm carried in sling. There were no enlarged glands in axilla.

The left breast was normal.

The right breast was removed. Fig. 3 represents the state of affairs discovered upon microscopical examination of large sections made from this breast. D is the situation of the nipple, and E E E is the direction of ducts towards it. A normal duct is to be seen at C C C. A duct, A, contains a large multiradicular papilloma, apparently benign. Another duct, B, contains the beginnings of multiradicular papillomata at three separate parts. In another part of this breast, and having no connexion with parts connected with the distribution of ducts A and B, there existed a small fibroadenoma.

CASE III.

Female, aged 58 years, married, no children.

Left breast: Nipple not retracted, no discharge from nipple. Multiple fluctuating swellings in breast of various sizes, the largest one inch in diameter. The margins of all were defined. The skin was not attached to them. She had noticed the swellings several months ago. The breast had been painful, but was not painful upon a clinical examination. There were no enlarged lymphatic glands in axilla.

The right breast appeared normal on clinical examination, and no complaints were made of it.

The left breast was removed and whole sections of the breast were cut. Many cysts were discovered of duct and acinous origins, but in a segment of its outer margin was a small cancer in the gland, and which was infiltrating the fat outside it (see Fig. 4). The whole lesion was about the size of a sixpence. In another part of the gland there was a duct containing a multiradicular papilloma.

This patient is considering whether she will take advice and have a more complete operation performed.

CASE IV.

Female, aged 42, married, two children, which she nursed.

Right breast: No retraction of nipple; no discharge from nipple. Contained in an upper and outer segment a large fluctuating swelling. Its surface was not adherent to skin. The breast was nodular throughout; some nodules were painful and irregular in size, and many were distinguishable as parts of tortuous ducts. There were no enlarged glands in axilla.

The left breast appeared normal.

The right breast was removed, and whole sections of the breast were cut for microscopical examination. I found that the

^{*} Further Contributions to the Study of Cysts and Papillomata of the Breast, by Sir Lenthal Cheate, *The British Journal of Surgery*, vol. ix, No. 34, 1921.

SIR G. LENTHAL CHEATLE: CANCER OF THE BREAST.

FIG. 1.—Transverse section of small duct. At five or six different parts the epithelial lining is undergoing a desquamative hyperplasia, and by this process the duct is being distended. The connective tissue cells are proliferating at B and at other parts. There are no lymphocytes present in this section.

FIG. 2.—Reproduction of a duct, A, towards its termination in branches marked B. The lumens of all are enlarged and full of stagnant desquamated cells. The connective tissue outside them is dense and contains a limited number of lymphocytes, which are in immediate juxtaposition to the outside walls of the duct and its branches. It is interesting and instructive to note that the acini leading into the ducts at C1 and C2 and elsewhere are not in the least distended, a fact that suggests the cause of distension of the ducts is not due to obstruction higher up in their courses. $\times 30$.

fluctuating swelling contained besides its fluid contents a large papillomatous malignant growth. The epithelial cells lining the papillomatous parts of its structure are columnar.

In the main parts of the tumour the epithelial cells are rigidly confined within what presumably is the cyst wall; but at isolated intervals the epithelial cells can be seen most definitely passing through the wall and invading the surrounding connective tissue. There was a blood clot in this cyst. In other parts of this breast in the neighbourhood of the tumour can be seen ducts the epithelial linings of which are in a marked state of desquamative hyperplasia, very similar to that seen in Fig. 1. Upon the condition of things being discovered by microscopical examination the patient was advised to undergo a complete operation.

The sections have been cut too recently to enable me to get them drawn for reproduction in this article. At a future date I will publish drawings made from them. The sections are at the disposal of all who care to see them.

SUTURE OF RUPTURED LEFT DOME OF DIAPHRAGM.

BY

C. JENNINGS MARSHALL, M.S., M.D., F.R.C.S.,

ASSISTANT SURGEON, KING'S COLLEGE HOSPITAL.

With Radiographic Commentary by ROBERT KNOX, M.D.,
Radiologist to the Hospital.

[WITH SPECIAL PLATE.]

A YOUTH of 19 was admitted to hospital for injuries sustained by being thrown from his bicycle. While riding, he was thrown by a skid in front of a motor bus; he fell with the right knee undermost and then lost consciousness for a while. Unfortunately, no further details of the accident were available. At any rate, it was certain the bus passed over no part of his body. I saw the patient after attempts had been made to reduce a supposed dislocation of the knee.

There was severe shock on admission; temperature 97°, pulse rate 120, respirations 20 to 24. The right knee presented a fracture separation of the femoral epiphysis with a longish piece of diaphysis attached on the inner side. The lower end of the diaphysis presented on the outer side of the joint subjacent to the skin, which was whole. This particular displacement of fragments was probably accounted for by the manipulations.

Pain was complained of in the chest at the left costal margin about the eighth and ninth cartilages. There had been no dyspnoea, no sputum, no nausea or vomiting. The pain disappeared at the end of four days, and there was no symptom of trouble at all.

The skin showed no sign of trauma. The left chest moved but little, and was less expanded than the right. Slight cardiac pulsations were seen on the right side of the sternum; the right border of the heart was made out nearly two inches to the right of this; the left border was not determined.

Over the front of the left chest was tympany, extending up to the second rib and out to mid-axilla, where it began to yield to what became at the back slight relative dullness. Auscultation over the tympanitic area showed distant breath sounds at the upper and outer margins, none at the inner and lower part. Elsewhere the chest gave signs of partial collapse of the lung. At the left margin of the sternum there was heard at intervals a loud, intensely metallic clanking, synchronous with the heart beat. After a fortnight borborygmi were heard in the chest; they were infrequent, but had great intensity even as high as the apex of the axilla. Against the possibility of their being transmitted one noted: (1) their maximum intensity was thoracic, (2) their loudness, and (3) apparent origin very near the stethoscope. On a few occasions it was possible, by a sudden dipping of the finger-tips in the epigastrium, to cause a splash to be heard in the thorax. Diaphragmatic hernia was diagnosed on these grounds clinically.

On the fifth day the femur was plated under spinal analgesia. As the lower fragment offered nothing on the outer side but a very short cancellous strip, the incision was made on the inner aspect. Anatomical reposition was effected. Movement of the joint was begun a week later, and the other forms of physiotherapy added as soon as the scar permitted.

The condition of the lower limb had for this time made a proper x-ray examination of the chest impossible. The patient was screened in the supine position. The cardiac displacement was evident; the left pleural "cavity" was poorly illuminated compared with the right, and the diaphragm on that side was some one and a half inches lower. Bismuth gruel swallowed by the patient passed rapidly, to lie as a mass 3 inches by 2 inches transversely at the level of the fourth left costal cartilage.

It was thought that operative interference was indicated, though in bed the patient had no symptom, alimentary, respiratory, or cardiac. It is probable, however, that the condition of affairs on exertion might cause considerable embarrassment; further, the possibility of internal strangulation had to be borne in mind.

Intratracheal ether with the patient in the lateral position was given, the table being angled opposite the lower costal margin to "open out" the chest towards the operator. Five inches each of the eighth and ninth ribs were resected in the axilla. The pleura was opened at the upper part of the lower periosteal tube, for its

whole length. The in-sucking of air was very slight and there was no respiratory disturbance (as, indeed, was to be expected, nearly the maximum mediastinal displacement already being present). The first structure encountered was the omentum at the greater curve of the stomach. This organ, distended to the size of a coco-nut, lay uppermost; below it was the transverse colon, and on the inner side were many coils of small intestine. The left edge of the liver just projected into the pleural cavity. The spleen was not seen.

The rent in the diaphragm began through the oesophageal opening and extended outwards and forwards for three inches, skirting the back of the trefoil. Except at the oesophageal end of the anterior limb of the tear, omentum was adherent to the edges (on the abdominal aspect chiefly). The lesser curve of the stomach appeared only at its middle, while three-fourths of the greater was seen. The cardia and the lower end of the gullet were not displaced or loosened. The gap in the diaphragm was a little over an inch at its widest; the edges were strongly resistant to a greater increase than half an inch in this direction. It seemed that strangulation was a very distinct possibility. The viscera were reduced and retained by a moist pack. At first this seemed almost impossible of accomplishment, but when the head was elevated by adjusting the table, and a supporting sandbag removed from the front of the abdomen, it became a matter of the utmost simplicity. The edges of the rent were freshened, and omental adhesions separated. Four mattress sutures were used to draw the inner edge a quarter of an inch under the outer, whose loose edge was then attached by a close row of interrupted sutures to the upper surface of the inner edge. A widely imbricated line of union was thus secured. There was but little tension to overcome in getting the edges together.

The parietes were then closed, the pressure of intratracheal insufflation being considerably increased. It was observed that the lung inflated rapidly, particularly the lower lobe, whose margin at the completion of the parietal suture extended below the line of incision and actually touched the chest wall.

Convalescence was interrupted only by bronchitis for three days. The absorption of the remaining air was gradual, H.A.B. becoming normal in six weeks.

The patient was seen again eight months afterwards. There were no symptoms of any kind. He indulges in cricket and other sport, cycles and does heavy work. The range of knee movement was 70° to 180°.

To suspect such a condition is usually to diagnose it. Symptoms offered no help, except that the complete absence from the start of dyspnoea seemed rather striking. The mechanical condition was very similar to that in pneumothorax—lung collapse, mediastinal displacement, air in the pleural cavity, albeit enclosed in bowel.

The loudness and superficial character of the rare borborygmi were nearly conclusive. The succussion splash induced by a jerk in the epigastrium may be worthy of note. The fact that the heart does not return to its place is suggestive if one has a case suspected as pneumothorax under observation for some time. A positive bismuth meal result is conclusive, as the stomach must almost inevitably be the viscus first herniated. It is conceivable that a part of the fundus might be so gripped as to prevent the entry of the bismuth. Obviously the supine position is to be adopted. The rare herniation of intestine alone—for example, colon—could be excluded or detected only by frequent examination.

With regard to the operation itself, the whole procedure was emphatically easy. While resection of ribs might have been avoided by a powerful retractor, I doubt whether the access would have been so comfortable. A longer incision would have been needed, and the window into the chest would have had ends not so wide as the middle, which here seemed a helpful feature. One thing is certain: approach from the abdomen would have been far more difficult, reduction would have been more difficult, and a deliberate imbrication of the rent could not have been done. I should like to emphasize the very striking help of posture in effecting reduction. It was noted how tough is the diaphragm—one felt satisfied that no sutures could possibly tear out.

While one did not fear any sudden change of intrathoracic tension, insufflation ether seems to be far the best for this purpose; the anaesthesia given by Dr. Richards was entirely uneventful, and at the end provided the most gratifying feature of the whole procedure—the rapidly expanding lung after its two months' collapse. The upper lobe did not expand anything like as much as the lower, possibly as this received the direct blast from the trachea.

Comments on Radiographic Appearances.

The radiograph, taken on the first visit (September 4th, 1920) to the x-ray department, shows an interesting condition on the left side of the thorax. The outline of the diaphragm is irregular and very ill defined. Above it, as high as the second intercostal space, is a translucent area which is limited in the mid-line by the spine and laterally by the ribs and intercostal spaces. Its upper margin is well defined