sterilized at the time Caesarean section is performed, or should she be allowed to run the risk of becoming pregnant again? So recently as ten years ago it was widely held that the majority of cases should be sterilized at the time of the first Caesarean section. At the present day the general view is in complete opposition to this, and it is held to be rarely justifiable to sterilize a patient except in such circumstances as we have already considered, where the condition found at operation renders this course necessary in order to obtain a successful result—for example, hysterectomy for fibroids. Now that Caesarean section is so frequently performed for conditions other than obstruction in the pelvis, obviously there is no reason why many of the mothers should not have any number of subsequent labours—and a number do so. In cases of pelvic contraction, however, the view generally held is, "once a Caesarean always a Caesarean," and there are many cases on record of women undergoing Caesarean section four, five, or even more times, quite successfully. Unfortunately it is a fact that in a certain number of cases the uterine scar is not strong enough to withstand the strain of a subsequent pregnancy, and rupture of the uterus has been found to occur far too frequently. The direct cause of the weakness of the scar is sepsis at the time of the original operation.

Another fact to be noted is the relatively high rate of sterility following the operation. This, I think, is due not so much to the operation itself as to the fear that it engenders in the minds of the patients and their husbands and consequent avoidance of pregnancy.

In conclusion, I should like to emphasize, first, the supreme importance of educating ourselves and—a far more difficult matter—our patients and their mothers and mothers-in-law in the absolute necessity of keeping a careful watch on all women throughout the period of pregnancy; it is the only means of avoiding occasional disaster. Secondly, the urgent need for the provision of maternity beds in all large towns and country districts. Possibly since the introduction of the panel system and the almost universal employment of midwives in maternity cases in industrial districts we may be able to view these questions in a more detached spirit than heretofore. In any case the remedy lies first and foremost in our own hands; if we make it our business to insist that beds for maternity cases are provided, I feel convinced that the time will not be far distant when women will clamour to be admitted to them.

LITERATURE.

Ga'abin and Blacker: *Practice of Midwifery*, seventh edition, 1910. Herman: *Difficult Labour*, fifth edition, 1910. BRITISH MEDICAL JOURNAL, Reports of Section of Obstetrics and Gynaecology, October 1st, 1921. Ibid., December 10th, 1921. *Proc. Roy. Soc. Med.*, vol. xiv, I, November, 1920.

CAESAREAN SECTION.*

BY

SAMUEL J. CAMERON, M.B., F.R.F.P. AND S.GLAS.,

ASSISTANT TO THE REGIUS PROFESSOR OF MIDWIFERY, GLASGOW UNIVERSITY; OBSTETRIC SURGEON AND GYNAECOLOGIST, GLASGOW MATERNITY AND WOMEN'S HOSPITAL.

CAESAREAN SECTION is one of the most spectacular but also one of the easiest of abdominal operations to perform if the patient has not undergone the operation on a previous occasion. With practice this operation can be accomplished rapidly; with an efficient staff of nurses and assistants I have performed the entire operation in seven minutes. A general practitioner with an average surgical training may safely perform it in cases of obstruction due to contracted pelvis, provided the case comes under observation early in labour and that he can secure thorough sterilization of instruments, suture materials, and gauze. On reviewing my more recent results I find that I have lost one patient in seventy-nine consecutive cases of Caesarean section. The only case which died succumbed to an ordinary bronchopneumonia ten days after operation. Ether alone had been administered in this case, and I would strongly urge that for purposes of anaesthesia chloroform should be used in preference to ether in the case of rachitic patients, as they are extremely susceptible to certain pulmonary disorders; a large number of them suffer from bronchitis and emphysema.

Within recent years the indications for the operation have been extended, and at the present time the obstetrician will be most liable to be called on to employ it in one of the following conditions:

1. In all cases of contracted pelvis where the obstruction prevents effectively the foetal head from engaging at the brim. In studying the causes of operative failure in my earlier cases of contracted pelvis I found that the chief danger was sepsis. At that time I performed the operation, even if forceps had been employed unsuccessfully by the practitioner, or if the membranes had been ruptured for many hours. As a rule the patients recovered, but a few died, and others had a prolonged convalescence due to septic infection. Accordingly I determined to operate only in cases which were probably free from contamination, and therefore I performed craniotomy in most cases which had been examined vaginally before admission to hospital, and in all of these cases if the membranes had been ruptured for longer than twelve hours.

2. Neoplasms in the uterus, ovaries, or pelvic bones sometimes cause such a degree of obstruction that Caesarean section has to be performed. As a general rule, cystic tumours of the ovary can be removed through the abdominal wound without incising the uterine wall, and whenever this is feasible it should be the operation of choice. In order to minimize the strain to which the abdominal wound is subjected during labour forceps should be employed when the second stage is reached. On the other hand, we can seldom remove a fibromatous tumour which has obstructed labour without first performing Caesarean section; nevertheless it is possible when a subperitoneal tumour is pedunculated, and in one instance of this kind I was able with my hand in the vagina to push the tumour upwards over the foetal head, and by so doing operative interference was avoided at the time. When a patient with a fibroma does not wish to be sterilized, occasionally we are able to perform myomectomy immediately after delivery has been effected by Caesarean section. As regards carcinoma of the cervix, the radical operation should be practised as soon as the case comes under observation if there be a reasonable chance of success; the child's life is of secondary importance. If the disease be too advanced to permit of extirpation, pregnancy should be allowed to proceed, and at full time Caesarean section should be followed by supravaginal hysterectomy, as the risk of sepsis is thereby diminished.

3. Excessive cicatricial contraction of the cervix and vagina may so obstruct labour that the lives of mother and child can be best safeguarded by Caesarean section. As a rule this difficulty follows an operation where the cervix has been amputated and the vaginal walls have been extensively denuded and sutured for the cure of uterine prolapse.

4. The operation is valuable in certain cases of accidental haemorrhage and placenta praevia, but in placenta praevia I recommend it mainly in cases where the patient is elderly and the child is alive. It seems to me that the operation should not be recommended in such cases if the patient be young, as she may be left with a weak uterine scar which may give way during a subsequent pregnancy. It is also a justifiable procedure in any case where the placenta is central and the os is not dilated. Many lives have been saved in cases of severe accidental haemorrhage by means of this operation. If the cervix be closed and the bleeding profuse it appears to be the most certain method of controlling the haemorrhage. Theoretically the proper procedure would be to remove the uterus without incising it, as the patient can ill afford any additional loss of blood, and by adopting this plan the main vessels supplying the organ are at once clamped. In practice I find, however, that evacuation of the contents shortens the operation, as the surgeon is not hampered by a bulky organ, and, after all, the loss of blood is relatively small. In cases of accidental haemorrhage I usually perform hysterectomy immediately after removal of the foetus and placenta, but this procedure is not always necessary, and cases may make an uninterrupted recovery after the uterine wound has been sutured in the usual way.

5. There is a tendency at present to advocate Caesarean section in cases of eclampsia, but as far as my observations go much better results can be obtained by less drastic measures which leave the uterus intact. I refer to induction of labour in combination with medicinal and other measures.

6. Vento-fixation and vaginal fixation of the uterus may disturb the position of the organ to such an extent that delivery by the natural passage is impossible, and in such circumstances Caesarean section is indicated. I have had personal experience of three cases of this character.

7. In rare instances this operation is performed in cases where labour is obstructed by a retraction ring which has formed in advance of the presenting part.

* Abstract of a lecture delivered to the Medico-Chirurgical Society, Queen Margaret's College, Glasgow.

Time of Operation.

The best time for operating on the patient is during the first stage of labour before the membranes have ruptured, as cervical dilatation ensures free drainage. In most instances, to suit my own convenience, I operate before labour begins. The patient is prepared in the usual way for an abdominal operation, and if a vaginal examination has been made recently a few antiseptic douches should be administered.

Site of the Abdominal and Uterine Wounds.

For many years I have operated on my gynaecological cases either through an incision in the rectus sheath or through a transverse wound. More recently I have adopted the incision through the rectus sheath in cases of Caesarean section, owing to the fact that a weak wound is liable to be obtained by a middle-line incision, as the abdominal wall in this situation is attenuated by the large gravid uterus. It is a matter of importance as to whether the right or left side is chosen. In my opinion the incision ought always to be made on the right side, as the uterus usually lies towards the right, and therefore the left margin of the organ approaches the middle line of the abdomen. Over a year ago I was performing the operation in the country, where the lighting equipment was miserable, and the uterus was exposed through an incision in the left rectus sheath. As soon as the uterine incision was made profuse haemorrhage occurred from the lower angle of the wound, and it was found that some large veins in the left broad ligament had been severed; great difficulty was experienced in arresting the haemorrhage. The left border of the uterus in this case was situated near the middle line, and if the incision had been made on the right side of the abdomen the above unpleasant complication would have been avoided. Two knives are always used by me—one for the abdominal and the other for the uterine wall—and I regard this as an important detail in my technique as it prevents infection of the uterine tissues if the skin be not sterile.

Every surgeon who has had a large experience in dealing with cases of repeated Caesarean section has probably on many occasions been hampered by numerous and dense adhesions in the region of the former scar or scars. The attachment of the uterus to the abdominal wall can often be diagnosed before the abdomen is opened, owing to movements communicated to the uterus being accompanied by dragging on the abdominal wall. If attached in the usual way the operation in these circumstances may be dangerously prolonged, and I have known a patient in the practice of an accomplished obstetric surgeon succumb from shock following a tedious operation of this description. In dealing with these cases I strongly advocate the evacuation of the uterine contents through transverse incisions in the abdominal and uterine walls. By so doing the surgeon can avoid the labyrinth of adhesions, and as the uterus is usually firmly fixed to the parietal peritoneum the organ will remain there in a state of ventro-suspension. Sometimes this attachment of the uterus to the parietes favours incomplete retraction, and on several occasions I have observed fairly profuse *post-partum* haemorrhage in such cases. As a rule I make the transverse incision a short distance above the upper extremity of the old scar; this will usually be above the level of the umbilicus and near the fundus of the uterus. On opening the abdomen the free portion of uterine wall near the fundus is severed transversely, and the child is extracted.

Treatment of the Uterus.

Before opening the uterus I insert four large swabs, so that two are situated laterally and the other two at the upper and lower angles of the abdominal wound. The swabs are not removed until the wound in the uterine wall has been closed. After the uterus has been opened the operator should feel for a leg and drag the limb through the wound. If he discover that he has seized a hand he should at once release it and find a leg, as the child should be delivered as a breech. As the limb is often greasy, the nurse who is attending to the dressings should keep a swab in readiness to pass to the operator to prevent his hand from slipping during the extraction. Immediately the child is delivered the uterus should be drawn through the abdominal wound on to the abdominal wall. There the organ should be turned inside out, so that the placenta and membranes can be stripped from their attachment. For this purpose a large pledget of gauze will be found most useful. In most instances the membranes in the region of the cervix are the most difficult to detach, and it is highly desirable that they should be entirely removed, as

even a small portion overlying the os may prevent the lochia from escaping for several days.

When the extraction of the placenta and membranes has been accomplished the inner wall is thrust back into position in order that the uterine wound may be closed. The method which I now adopt is to begin by inserting three interrupted sutures of silk; one suture is placed in the middle of the wound and the other two midway between the first suture and the upper and lower angles of the wound. Each suture passes through the entire depth of the uterine wall, with the exception of the inner layer. These sutures are firmly secured, and the remainder of the wound is closed with similar sutures of catgut. Finally a continuous suture of catgut is used throughout the entire length of the wound. The three sutures of silk are a safeguard in case the catgut becomes absorbed too rapidly.

Closure of the Abdominal Wound and After-treatment.

As soon as the wound in the uterus is closed the four large swabs are withdrawn and the abdominal incision is closed in layers. For many years I have used anchor sutures to support the wound and to keep the small gauze dressing in position. No other covering is applied to the wound. Before the patient leaves the operating table the surgeon should observe whether blood is escaping from the vagina or not; if it is not, he should grasp the uterus and compress it through the abdominal wall. Dilatation of the cervix may sometimes be necessary to remove an obstructing portion of membrane.

In my surgical practice I have for many years allowed my patients every liberty of movement as soon as they become conscious after the operation. They are encouraged to lie on their side and to sit upright in bed in Fowler's position. By so doing drainage is promoted, flatulence is diminished, convalescence is shortened, and the liability to thrombosis is lessened.

ECLAMPSIA: EVOLUTION AS A CAUSATIVE FACTOR.

By S. E. KARK, M.B., CH.B. EDIN.,
CAPE TOWN.

BEFORE proceeding in detail with the subject-matter of this paper I must briefly summarize a paper dealing with the general causation of the toxæmias of pregnancy read by me at the recent South African Medical Congress held at Cape Town (October, 1921).

1. A normal pregnancy is one in which all the changes that take place incidental to that condition are necessary for the purpose of gestation—that is, all the changes brought about must be functional.

2. As with almost all pregnancies there are concomitant functionless changes apart from frequent pathological-like manifestations, it is evident that almost all pregnancies are not quite normal.

3. To explain this anomaly in nature—that such an important physiological process as gestation should almost always be abnormal—I advanced the view that in the process of evolution of the various vital functions of the body the placental form of reproduction was evolved last and comparatively late. While all the other vital functions are now mature and harmoniously correlated to each other, gestation is still in some degree developmentally immature, hence its many physiological imperfections. In consequence of this tardiness in the process of evolution of the placental form of reproduction, the pregnant state approaches in some degree primitive conditions; gestation is, as it were, a physiologic atavism. This is evidenced by the following facts:

A. That in the pregnant woman the secondary feminine characteristics often become less marked; she approaches the masculine type in many respects—that is a recession towards the primitive state, in which secondary sex characteristics were not as yet markedly differentiated. This metamorphosis towards masculinity is shown by the frequent occurrence in pregnancy of:

- (a) Overgrowth of hair on the body, especially on the face.
- (b) Coarsening of the features.
- (c) Thickening of the skin.
- (d) Hypertrophy of some of the bones.
- (e) Enlargement of the hands and feet.
- (f) The stability of calcium metabolism.

B. Another result of this physiologic atavism of pregnancy, and one fraught with even more important and far-reaching