Case.	Clinical.	Radiographic.	Fract. Test Meal.	Verified.	Case.	Clinical.	Radiographic.	Fract. Test Meal.	Verified.
6	Gastric ulcer	? Duodenal ulcer	Type 2	Pyloric ulcer.	72	Gastric ulcer	Gastric ulcer	Type 2	Large ulcer, lesser curvature.
13	" "	? Duodenal ulcer	Type 5	Appendicitis.	77	?	Delay in large intestines	Type 5	Appendicitis.
17	" "	Delay in emptying, otherwise normal	Type 2	Ulcer on lesser curvature.	78	?	Nil abnormal	Type 2	Nil at operation.
18	Recurrent gastric ulcer	Gastro-enterostomy working	Type 6	Perigastric adhesions only.	80	Gastric ulcer	Nil abnormal	Type 2	Ulcer near cardia.
20	Gastric ulcer	Two indentations on greater curvature	Type 2	Ulcer on lesser curvature.	82	?	Hour-glass stomach or ulcer	Type 1	Perigastric adhesions.
22	?	Nil abnormal	Type 5	Nil found at operation.	83	?	? Pyloric ulcer	Type 2	Distended duodenum (? Wilkie).
27	?	Nil abnormal	Type 4	Pyloric cicatrix.	85	?	Nil found: ? duodenal obstruction	Type 2 and 5	Pyloric ulcer; appendicitis.
28	Gastric ulcer	? Pyloric stenosis	Type 4	Pyloric stenosis.	92	Gall stones	? Colitis	Type 5	Enteroptosis.
30	Duodenal ulcer	Small hypertonic stomach	Type 3	Duodenal ulcer.	103	Gastric ulcer	Hour-glass stomach	Type 1	Cardiac ulcer.
32	Gastric ulcer	Hour-glass stomach	Type 2	Two ulcers on lesser curvature.	106	" "	No record	Type 5	Pyloric ulcer.
35	" "	? Gastric ulcer	Type 6	Minute ulcer near pylorus.	115	Pyloric spasm	Pyloric stenosis	Type 1	Nil found.
37	" "	? Duodenal ulcer	Type 2	Carcinoma on lesser curvature.	116	Gastric ulcer	? Pyloric ulcer	Type 3	Duodenal ulcer.
38	" "	Pyloric ulcer	Type 4 and 5	Pyloric ulcer.	117	" "	Nil abnormal	Type 1	Mobile kidneys.
45	" "	Nil abnormal	Type 1	Nil abnormal.	118	" "	Hypertonic stomach ? duodenal irritation	Type 6	Mobile kidneys.
46	?	? Duodenal ulcer	Type 1	Nil abnormal.	119	Carcinoma of stomach	Fundal new growth	Type 4	Simple pyloric stenosis.
56	Carcinoma of stomach	Nil abnormal in stomach	Type 6	Gall stones.	123	Gall stones	? Pyloric stenosis	Type 1	Gall stones.
57	Gastric ulcer	Ulcer middle of greater curvature	Type 1	Small ulcer of fundus.	141	? Nil	? Duodenal ulcer	Type 5	Duodenal ulcer.
58	" "	Duodenal ulcer	Type 3	Duodenal ulcer.	142	?	? Pyloric stenosis; ? duodenal ulcer	Type 4	Simple pyloric stenosis.
64	" "	? Pyloric ulcer	Type 1	Gastric ulcer.	151	?	? Duodenal ulcer	Type 2	Ulcer near pylorus.
67	?	? Prepyloric ulceration	Type 6	Mobile kidneys.	152	Pyloric ulcer	? Duodenal irritation	Type 4	Pyloric stenosis.

Curve Type 5 (Fig. 5).

This is the "climbing" curve or curve of tardive hyperacidity (Hawk), due presumably to deficient relaxation or spasm of the pylorus of a non-continuous type; this picture has also been seen in what can only be called idiopathic hyperchlorhydria, since no spasm or cause for spasm could be recognized. A great majority of these curves were, in my own series, associated with either small pyloric ulcers (leading presumably to spasm), or with subpyloric conditions, such as appendicitis, where perhaps pyloric spasm was a reflex event similar to that observed in the experimental intestinal injuries of Cannon and Murphy. Removal of the subpyloric condition, which was postulated as a cause of the "reflex hyperchlorhydria" expressed by this type of curve, has, in my own series, been followed by an alteration approximating to Type 1.

Curves of this type were obtained in six cases which came to operation, and which exhibited lesions varying from a small pyloric ulcer to an inflamed appendix.

FIG. 5.—Curve Type 5.
Case 81. FIG. 6.—Curve Type 6.
Case 108.

Curve Type 6 (Fig. 6).

The so-called achylic curve, which may mean deficient secretion or early and complete neutralization, has personally been obtained in a variety of conditions which appear unrelated. These included normal cases and cases of gastric ulcer, gastric carcinoma, gall stones, movable kidneys, gastroptosis after gastro-enterostomy, and, in six instances, of exophthalmic goitre. There does not, therefore, seem to be any great diagnostic value in this type of curve, since it can be obtained in so great a number of conditions; on the other hand, it can serve as an indicator of the efficiency of the stoma when a gastro-enterostomy has been made.

From a general consideration of these curves valuable help in diagnosis would seem to be given, particularly by Types 2, 4, and 5. A comparison of this and other evidence with the actual finding as discovered by operation will, however, best illustrate this statement. The accompanying table shows that the fractional test meal method is worthy of comparison with the recognized method of opaque meals and x rays. Forty cases have been taken at random; in each the clinical, radiographic, fractional test meal, and verified findings have been set out.

In concluding this note, I would wish to express my indebtedness to my colleagues at the David Lewis Northern Hospital for the provision of some of the clinical material.

Conclusions.

1. The acidity curves plotted from fractional test meals are of some value in obtaining knowledge of the factors which modify the acidity.

2. Alterations in these factors bear some sort of relation to gastro-intestinal lesions; hence the acidity curves may throw some light on these lesions.

3. These curves seem divisible into at least six types of varying diagnostic value.

4. By comparison with other methods the fractional test meal is at least admissible as an aid to diagnosis.

REFERENCES.

- ¹ J. A. Ryle: *Lancet*, September 4th, 1910. ² A. F. Hurst: *BRITISH MEDICAL JOURNAL*, April 24th, 1920; Bolton: *Ulcer of the Stomach*, 1913, Part III, chap. II. ³ Boldyreff: *Transactions 11th Congress of Physicians*, St. Petersburg, 1909; *Quart. Journ. Exper. Phys.*, 8, 1, 1914. ⁴ T. Izod Bennett and J. F. Venables: *BRITISH MEDICAL JOURNAL*, October 30th, 1920. ⁵ Spencer, Meyer, Rebhuss, and Hawk: *Amer. Journ. Phys.*, March, 1916. ⁶ Charles Bolton and Gordon W. Goodhart: *Lancet*, March 4th, 1922. ⁷ Spencer, Meyer, Rebhuss, and Hawk: *Ibid.* ⁸ Schütle: *Zett. f. klin. Med.*, 1895, vol. xxviii. ⁹ J. A. Ryle: *Ibid.*

TREATMENT OF HUMAN HOOKWORM INFECTION
WITH CARBON TETRACHLORIDE.

BY

LUCIUS NICHOLLS, M.D., B.A. CANTAB.,

DIRECTOR OF THE BACTERIOLOGICAL AND PASTEUR INSTITUTES, CEYLON;

AND

G. G. HAMPTON, M.D.,

STATE DIRECTOR OF THE INTERNATIONAL HEALTH BOARD, ROCKEFELLER FOUNDATION.

In the *Journal of the American Medical Association* of November 19th, 1921, Dr. Maurice C. Hall, of the United States Bureau of Animal Industry, published a paper on the use of carbon tetrachloride for the removal of worms in animals. He stated that the results were satisfactory, and that almost all worms were removed by one dose. He recommended that 0.3 c.cm. should be given for every kilogram of weight of the animal. He tested the drug on monkeys, and gave two to five times this amount, and reported that no untoward symptoms ensued. Dr. Hall himself took in the morning 3 c.cm. of carbon tetrachloride, and afterwards did his usual day's work and was not inconvenienced in any way. He suggested that the drug would be valuable for the treatment of hookworm disease in man.

Several drugs are used at present for the treatment of this infestation, but all of them have disadvantages. Thymol and beta-naphthol, except in large doses, are ineffective, and are

difficult to administer to children, and occasionally deaths occur after their use.^{1 2}

Chenopodium has been used very largely in recent years, especially where extensive campaigns have been started among heavily infected populations in various parts of the tropics. It has these disadvantages:

1. It is objectionable to the patients.
2. It is frequently followed by depression, and it is necessary for the patient to remain at rest and away from work on the day that the drug is taken.
3. One day's treatment is ineffective with the majority of patients.
4. It is impossible to standardize chenopodium, which contains a large number of chemical bodies of unknown composition, and one sample of the drug occasionally is found to be more potent than another.
5. It is dangerous to administer to very young or very sick children.

Carbon tetrachloride, CCl_4 , can be prepared cheaply and in a high degree of purity. It is a colourless liquid with a sweetish, pungent taste, and it produces a slight sensation of burning when it is taken into the mouth. Its boiling point is 76.74°C . and its specific gravity is 1.63195. It is soluble in alcohol and ether, and will dissolve a large number of organic substances, especially those of a fatty nature. Previously it has been used in the practice of medicine as an anaesthetic and as an inhalation for hay fever. It was much used at one time as a dry shampoo, but it fell into disrepute because a woman, aged 29, died suddenly during its use for this purpose. Dr. B. H. Spilsbury stated at the inquest that the deceased was at the time of her death in a condition of status lymphaticus.³ The preparation used by barbers would not be the chemically pure drug; and the commercial preparation contains two very poisonous substances—carbon bisulphide and carbonyl chloride, and as much as 2 per cent. of the former may be present.

Most fat solvents, such as chloroform, tetrachlorethane, or carbon tetrachloride, if they are inhaled for long periods cause subacute hepatitis, which may be followed by advanced fatty degeneration of the liver.⁴ But these effects do not occur after one or two doses of these drugs.

We decided to test this drug especially with a view to the treatment of school children and young children who are admitted to hospital with various diseases and who are also heavily parasitized by hookworms. One of us in association with Dr. Leach was the first to use the drug in Ceylon, and fourteen persons were treated for hookworm disease with doses ranging from 3 to 10 c.cm. The patients took the drug well and did not show the dislike to it that most patients show when chenopodium is taken. They did their usual day's work and complained of no untoward symptoms.

A murderer under sentence of death volunteered to take the drug, and 6 c.cm. were given one hour after the midday meal, and the same day he passed four worms (*Ascaris lumbricoides*); thirteen days later this dose was repeated at 7 a.m. before any food had been taken. The patient stated that at 11.45 he felt a little giddy and sleepy, but this soon passed. The execution took place a week after the second dose, and a *post-mortem* examination was done one hour after death. No hookworms or ascarids were found in the intestinal tract. Sections of his organs were examined microscopically, but showed no signs of degeneration of any kind. Therefore it is apparent that in single doses carbon tetrachloride is a safe remedy.

Next it was tried on the students of an agricultural school, who were from 18 to 25 years of age. A preliminary examination of samples of their faeces showed that all of them were infested with hookworms. No purgative was given on the preceding night. They all received a dose of 3 c.cm. of the drug at 6.30 a.m. before any food had been taken; no purgative was given after the drug was administered. They were instructed to carry on their usual mode of life, and restrictions were not placed upon their work, play, or diet. Not one of them was inconvenienced by the drug, and they all continued their day's work in the gardens or at the school house. They were supplied with "chatties," which were numbered, and they were instructed to pass their stools into these vessels. In this way all stools passed for three days after treatment were collected and carefully searched for worms.

A total of 719 hookworms, or an average of 36 for each student, were recovered. Besides these 13 *Ascaris lumbricoides*

TABLE I.

No.	Age.	Haemoglobin by Taquist.	Examination of Faeces before Treatment.			No. Stools passed on Day of Treatment.	Worms present in the Stools.									Total Worms for 3 Days.			Examination of Faeces 10 Days after Treatment.		
			Hookworms.	Ascaris.	Trichuris.		1st Day.			2nd Day.			3rd Day.			Hookworms.	Ascaris.	Trichuris.	Hookworms.	Ascaris.	Trichuris.
							Hookworms.	Ascaris.	Trichuris.	Hookworms.	Ascaris.	Trichuris.	Hookworms.	Ascaris.	Trichuris.						
1	21	80	+	-	+	0				2						2			(-)	-	+
2	20	80	+	-	-	2	1						11			12			(-)	-	+
3	21	65	+	+	+	0				116	1		45			161	1		(-)	+	+
4	20	70	+	+	+	1	0			38			0			38			(-)	+	+
5	20	80	+	+	+	1	0			34			8			42			(-)	+	+
6	22	80	+	+	+	2	1			37	1		7			45	1		(-)	+	+
7	19	75	+	+	+	1	0	1		83			3			86	1		(-)	-	+
8	18	70	+	+	-	1	0			1			8			9			(-)	+	+
9	21	80	+	+	-	2	9			19			5			33			(+)	+	-
10	21	75	+	+	+	1	0			19	2		10	4		29	6		(-)	-	+
11	18	75	+	+	+	1	0			2			12			14			(-)	+	+
12	18	70	+	-	+	3	0			3			2			5			(-)	+	+
13	21	80	+	+	+	1	0			33	4		11			44	4		(-)	+	+
14	21	70	+	-	+	0				0			26			26			(-)	+	+
15	25	70	+	-	+	1	0			1			7			8			(-)	-	-
16	19	80	+	-	+	1	0			14			4			18			(-)	+	+
17	22	70	+	+	-	1	0			9			24			33			(-)	+	+
18	23	70	+	-	-	1	0			96			12			108			(+)	-	+
19	22	60	+	-	+	2	0			0						0			(-)	+	+
20	23	65	+	-	-	0				0			6			6			(-)	-	-
Total			...	...	...	22	11			507	8		231	4		719	13				

+ = Plain smear positive for ova. (+) = Centrifuged deposit positive for ova. (-) = Centrifuge and salt flotation method negative for ova.

were passed. Microscopical examination of faecal specimens ten days after treatment showed ova of hookworms in only two specimens, which indicates that 90 per cent. were cured. Table I is a summary of this work. It will be noted that two-thirds of the worms were passed on the second day after the treatment.

Next the drug was tried on sixty-four male and female students whose ages were from 7 to 17 years. Samples of their stools were examined microscopically for the ova of worms: sixty were positive for hookworms and four were negative.

The method of examination adopted was to make smears from the stool, and if ova were seen this was signified opposite the number of the student by a *plus* sign; if no ova were seen a portion of the stool was stirred up in water and centrifuged; the deposit was smeared on a slide and examined, and if it was positive it was signified by a *plus* sign in brackets thus (+). If by this method it was negative, a remaining portion of the centrifuged deposit was stirred up in concentrated salt solution, for the high specific gravity of this causes the ova to float to the surface; when a slide is brought into contact with this surface the ova will adhere to it. This is known as the "salt flotation method," and when stools contain so few ova that this method must be used to reveal them the result is signified by a *plus* sign in brackets and an asterisk thus (+)*. When they are negative by this method it is signified by a *minus* sign in brackets (-).

A dose of from 1 to 3 c.cm. of carbon tetrachloride was given to each student at 6.40 a.m., before any food had been taken. The drug was given with a small quantity of water. There was no frowning or other evidences that the children objected to the drug, and an hour afterwards they took their regular morning meal. One twelve-years-old boy vomited when he had finished his meal, but the manager informed us that this child often vomited. Each child was furnished with a "chatty," on which was a number corresponding to a number on a tag tied securely to the child's wrist. A dispenser was present during the time this experiment was being carried out, and when a child wanted to use its "chatty" the dispenser compared the numbers, so as to prevent mistakes in the use of the vessels. The manageress assisted with the girls in the same manner. Two children failed to go to stool on the day the treatment was taken; the others had from one to seven stools each, with an average number for the sixty-four children of 2.93 stools; which indicates that the drug is a mild aperient. A dose of magnesium sulphate was given on the second morning after treatment to save time in collecting the stools. This purgative was taken before the early morning meal and all the stools passed that day were saved.

Four hundred and six stools were collected during the two days, which is an average of 6.34 stools from each patient. Each day a small quantity of formalin was poured into these vessels to preserve the stools. These were taken to the laboratory, where each collection of faeces was washed thoroughly on a sieve of fine wire gauze (forty strands to the inch); all that would not pass through the mesh was washed out on to a large full-plate photographic dish and searched for worms. It was washed again and placed in small quantities at a time in Petri dishes, and examined first upon a white and then upon a black background, because hookworms are dark when they are engorged with blood and preserved in formalin, and show up well upon a white plate, whereas those that contain no blood are discovered more easily on a black background.

A total of 2,776 hookworms, or an average of 43.3, were recovered from the 64 collections of stools. But of these 5 were negative. Besides these, 260 ascarids were obtained from 32; 15 trichuris from 6; and 4,945 *Oxyuris vermicularis* from 58 students. The discharge of trichuris and oxyuris indicates that the drug acts throughout the intestinal tract, for we have not previously seen trichuris expelled after the use of other anthelmintic drugs, or *Oxyuris vermicularis* passed in such large numbers.

Ten days after the treatment samples of the stools of 54 of the students were examined for ova (unfortunately samples of the other 10 were not available as the students had gone away on vacation); 2 of these were positive for hookworms by the simple smear method, and 4 were positive by the combined centrifuge and salt flotation method, which is a percentage of 88 probable cures.

The drug was given to a number of young children who had previously been treated with chenopodium on one or more occasions; the amount of chenopodium which was given

to children of 4 years of age was 4 minims divided into two doses and taken early in the morning with an hour's interval between each dose. This is as much as may be administered safely to children of this age. Table II shows the result—namely, that carbon tetrachloride will remove all or a large number of hookworms which remain after treatment with chenopodium.

TABLE II.

No.	Age.	Sex.	Previous Treatment with Chenopodium.	Dose of Carb. Tetrachloride.	Worms Expelled.			Examination of Faeces after Treatment.		
					Hookworms.	Ascaris.	Trichuris.	Hookworms.	Ascaris.	Trichuris.
1	4	M.	Once, 10 days previously	10 minims	9	1	0	-	+	+
2	4	F.	Twice, 10 and 28 days previously	10 "	52	15	0	+	+	+
3	3	F.	Twice, 15 and 32 days previously	10 "	35	1	0	-	+	+
4	6	M.	Twice, 23 and 34 days previously	10 "	52	3	1	(+)*	+	+

(+)* = Centrifuge and salt flotation method positive for ova.

The drug was given to a number of children who had pyrexia from various causes, and were anaemic and emaciated; larger doses were used than with the children shown in Table II; the results were excellent and are shown for a few of the cases in Table III. None of the children were in any way inconvenienced by the drug.

TABLE III.

No.	Age.	Sex.	Disease.	Dose of Carb. Tetrachloride.	Temperature at Time of Treatment.	Worms Expelled.			Examination of Faeces after Treatment.		
						Hookworms.	Ascaris.	Trichuris.	Hookworms.	Ascaris.	Trichuris.
1	4	M.	Enteric	15 minims	102°	54	0	0	(-)	+	+
2	9	F.	Malaria	25 "	100.3°	167	4	4	(-)	+	+
3	5	F.	Continuous P.U.O.	15 "	101.2°	6	0	0	(-)	+	+
4	6	M.	Continuous P.U.O.	15 "	102.1°	31	0	0	+	+	+
5	4	F.	Irregular P.U.O.	15 "	100°	48	2	0	(-)	+	+

Conclusions.

1. Carbon tetrachloride is an efficient anthelmintic remedy for hookworms.
2. The drug may be administered safely in doses of 10 to 20 minims to children of 3 and 4 years of age, even when they are seriously ill from various causes.
3. It aids the expulsion of *Ascaris lumbricoides* if it is followed by a purgative, but it is not as effective as chenopodium in killing this worm.
4. The drug does not seriously deteriorate on keeping. Many children were given doses of carbon tetrachloride which had been stored in the laboratory for three years.
5. It is more valuable than chenopodium for campaigns against hookworm disease because—

- (a) Patients do not object to its taste.
- (b) It is not necessary to precede or follow the administration of it by a purge.
- (c) It is more efficient than chenopodium and has not the depressing effects of that drug.
- (d) It is much cheaper than any other drug that has been used.
- (e) It can be prepared in a high degree of purity, and a chemically pure preparation should always be used.
- (f) The person who is being treated can do his usual day's work.

Children when one year of age may be given 10 minims of carbon tetrachloride with safety, and this dose should be increased by 2 minims for each year of apparent age. Thus a child of ten would receive 28 minims, a youth of sixteen

40 minims ($2\frac{1}{2}$ c.cm.), and an adult dose should be from 50 to 80 minims ($3\frac{1}{2}$ to 5 c.cm.), according to the size of the patient.

Finally, chenopodium is soluble in carbon tetrachloride, and a mixture of one part of the former in four of the latter should prove to be efficient for the expulsion of *Ascaris lumbricoides*.

Our thanks are due to Dr. W. P. Jacocks of the Rockefeller Foundation and Dr. Catherine Anderson of the Lady Havelock Hospital for much kindly assistance.

REFERENCES.

- ¹ *Journ. Amer. Med. Assoc.*, May 29th, 1920, vol. lxxiv, p. 1503. ² U.S. Public Health Reports, January 17th, 1913, vol. xxviii, No. 3, pp. 119-124. ³ *BRITISH MEDICAL JOURNAL*, July 24th, 1909, p. 243. ⁴ *Ibid.*, May 17th, 1919, p. 605.

SOME OBSERVATIONS ON THE MECHANISM OF DRUG ACTION:

A STUDY IN PHARMACO-DYNAMICS.

BY

O. C. M. DAVIS, M.D., D.Sc.,

LECTURER IN MATERIA MEDICA IN THE UNIVERSITY OF BRISTOL;
PHYSICIAN TO OUT-PATIENTS, BRISTOL ROYAL HOSPITAL
FOR SICK CHILDREN AND WOMEN.

It has long been recognized that certain chemical compounds have very little effect upon the animal organism, whereas others produce a marked reaction when administered under similar conditions. The object of the present paper is to bring together some scattered observations which help to throw some light on such facts as these, and to explain the mechanism of pharmacological action. The subject will be dealt with under several headings, and the first one is naturally a consideration of the question of

Toxic Properties in Relation to Atomic Weights.

About the year 1815 Prout, who was a physician, published a paper in the *Annals of Philosophy* pointing out that in many cases the atomic weights of the elements were approximately whole numbers, and deducing from this that hydrogen was the primary matter from which all elements are formed by various condensations. The observations of Prout and the work of Newlands, Lothar Meyer, Mendelejeff, and others led to the well-known periodic classification of the elements.

This classification led to much valuable research and to many important discoveries, but apparently little has been written from the point of view of pharmacological relationships. A careful consideration of the periodic table reveals the fact that there are many relationships between the pharmacological reactivities of the elements (and their compounds) and their periodic grouping. A few examples will illustrate this point; if we compare the elements of the magnesium group we notice that the tendency is towards increase of toxicity with increasing atomic weight, the greatest toxicity being associated with mercury compounds. The oxygen group is also of interest, passing from a gaseous element to a typical solid non-metal, sulphur, thence to selenium and tellurium, with certain metallic characteristics. The hydrogen compounds of this series illustrate well the increase of toxicity with increase of molecular magnitude, starting with the innocuous substance water, and passing to the extremely toxic telluretted hydrogen.

A very valuable research was carried out by Michet¹ in 1881 on the relative toxicities of a number of metals. He experimented upon fishes with solutions of metallic chlorides, diluting the solutions so that the fishes could survive forty-eight hours in the diluted solutions. He applied the term "limit of toxicity" to the weight of metal present in the form of chloride which would be contained in 1 litre of such a solution. In his original paper he deplores the fact that no relationships between toxicity and atomic weight were observed. Unfortunately this is a case where an excellent research was largely thrown away from a scientific point of view by misinterpretation of the actual findings. Had he tabulated his results in weights proportional to atomic weights many relationships would have been revealed, but he was presumably ignorant of the nature of "molecular solutions." One example shows this fairly well.

	Michet's Values.	Michet's Values Recalculated and Multiplied to give whole Numbers.	Atomic Weight.
Calcium	2.4	600	40.0
Strontium	2.2	250	87.6
Barium	0.78	56	137.4

It is clear from this table that the toxicity of an element belonging to the same periodic family increases with increase in atomic weight, but, needless to say, such a law is subject to variations. It is highly improbable that such periodic relationships as have been referred to are accidental, and with increasing knowledge more light may be thrown on this interesting branch of pharmacology.

Influence of Solubility.

One of the factors long recognized as influencing pharmacological activity is the solubility of the drug concerned. There are so many well-known examples of this that it is quite unnecessary to refer to them in detail; it is sufficient to point out that in practical pharmaceutical chemistry various methods are used to increase the solubility of insoluble compounds; thus slightly soluble alkaloids are converted into more soluble alkaloidal salts, and in the realm of synthetic organic chemistry various groups are introduced into the molecule with the object of increasing solubility.

It has also been recognized that certain constituents of animal organisms exert a selective action in dissolving drugs introduced into the organism, and thus certain tissues are specially affected.²

Vapour Pressure.

Its vapour pressure is a factor of the greatest importance in its effect on the reactivity of the drug. For instance, chloroform, ether, and other volatile anaesthetics would be useless for the purpose if they did not possess a high vapour pressure. During the recent war the question of vapour pressure was of the utmost importance in connexion with gas warfare. It is terrible to think of the calamitous possibilities of easily prepared toxic substances possessing a high vapour pressure.

Relationships between Chemical and Pharmacological Reactivity.

The increased attention given to the synthesis of organic compounds during the middle and later years of last century has enabled chemists to determine the structure of chemical compounds and in collaboration with pharmacologists to demonstrate relationships between the chemical and physiological reactivity of drugs.

Some years ago³ the rates of hydrolysis of a considerable number of anilides were investigated by the writer. These compounds, of which acetanilide (antifebrin) and phenacetin are common examples, owe their activity in part to the fact that they are hydrolysed slowly within the organism with liberation of the parent base according to the general equation—

It was shown that there were definite relationships⁴ between velocity of decomposition and degree of toxicity as previously determined by various pharmacologists.

Chemical Affinity.

In a further research⁵ on isomeric substances having similar pharmacological action, it was shown that in certain cases chemical affinity and toxicity were quantitatively proportional. Such observations have more than a theoretical importance; for example, this difference in chemical affinity doubtless explains the differences observed by Breil and Nierenstein when carrying out experiments upon trypanosomes with sodium salts of the isomeric amino-phenyl-arsenic acids. These observers found that the para compound, also known as atoxyl, is very reactive, the meta compound less so, and the ortho compound inactive.⁶

The relationship between the compounds is shown by the following formulae:

Another example of varying pharmacological activity is seen in the case of the isomeric hydroxy-benzoic acids, the ortho compound, better known as salicylic acid, being the only one possessing marked reactivity.

"Affinity Constants."

Ortho-hydroxy-benzoic acid	...	0.102
Meta-hydroxy-benzoic acid	...	0.00367
Para-hydroxy-benzoic acid	...	0.00286