has not only been seen but explored by opening, it is impossible to be certain that disease is absent. In the early stages of gall-bladder disease cure by surgery is sure and safe; in the presence of such complications as stone in the common duct and acute infections the risk becomes greater, and malignancy may be the termination of chronic simple gall-bladder disease.

When the gall bladder is diseased its removal is indicated—in two stages, if life or the integrity of the common duct is threatened if done in one. Drainage alone will not certainly cure, nor will it prevent the later onset of carcinoma in gall bladders believed, on external examination, to be healthy at the time of drainage, after removal of calculi. I have known it to arise from one to four years after drainage of what was considered a sufficiently healthy gall bladder to leave after the removal of calculi.

One of the most recent advances in abdominal surgery is that dealing with disease by splenectomy. Although we know that removal of the spleen is followed by restoration to health in suitable cases, until we are fully cognizant of its functions we cannot understand the part it plays in disease and why removal in certain cases is efficacious. That long life was possible after removal of the spleen in man we owe to clinical observation after splenectomy for injury. Interference next followed in such conditions as wandering spleen with torsion of the pedicle or the rare tumours. More recently, diseases of the spleen associated with blood changes have been successfully treated; I allude to such groups as splenic anaemia, Banti's disease, and splenomegalic jaundice. In the former, restoration to health occurs, and so far as we can ascertain no evil effects follow; we can justifiably speak of cure. It may be otherwise when operation is carried out in the later stage of the disease when cirrhosis of the liver has supervened. With this complication, when haemorrhages have taken place before operation, although apparent restoration to health is the rule, yet death from haematemesis at a later date is not unknown, in spite of the fact that splenectomy has been suggested as the treatment for certain cases of haematemesis in the absence of gastric or duodenal ulcer, splenomegaly, or recognizable hepatic cirrhosis. Our limitation here is due to neglect of treatment in the early stages of the disease.

Of the cause of the disease and the splenic enlargement, whether due to thrombosis—and if so, the infecting agent and its organ—we are in ignorance. Why removal of the spleen alters the blood picture is also beyond our ken. We are sadly limited in our dealings with diseases associated with splenic enlargement in not knowing their cause.

I know of no condition in which recovery is so dramatic as in splenomegalic haemolytic jaundice. In the familial type discoloration may have persisted from the early years of life, and yet within forty-eight hours of removal of the spleen it has disappeared. The fragility of the red cells, part of the disease, may be unaffected by splenectomy. As in other chronic abdominal diseases, general exploration has to be carried out, particular attention being paid to the gall bladder and ducts, as these cases are often complicated by pigment calculi.

Splenectomy in pernicious anaemia and in leukaemia is experimental, and until more evidence is forthcoming we should stay our hand.

Surgery has many legitimate triumphs in the abdomen. Let us see to it that it is not brought into disrepute by operations for trivial complaints, and that removal of the appendix or gall bladder, or fixations of the colon, do not become common procedures in every discomfort in this region.

PAIN IN GENITO-URINARY AFFECTIONS:

ITS VARIATIONS AND THEIR INTERPRETATION.

BY

W. K. IRWIN, M.D., F.R.C.S. EDIN.,

HONORARY CASUALTY OUT-PATIENT SURGEON, ST. PAUL'S HOSPITAL
FOR GENITO-URINARY DISEASES.

THE variations of pain in genito-urinary affections are so numerous and often so misleading that their interpretation may be difficult. It is hoped that the following notes will facilitate the early recognition of the causes at work.

Pain arising in this system frequently shows itself in parts other than the seat of the lesion, and it may be well at the outset to emphasize the distinction between (1) *site pain*, which is felt in or over the lesion, and (2) *referred pain*,

which through abnormal perception by the senses is assigned to the region supplied by an anastomosing nerve trunk or to the distribution of an irritated nerve.

Renal and Ureteric Regions.

Affections of the kidney are comparatively painless if the ureters remain patent. *Renal calculi* may give rise to three varieties of pain: (1) *site pain*, (2) *referred pain*, (3) *renal colic*. The *site pain*, which is more prominent during the diurnal hours of activity, is a dull ache, and is felt in the front of the loin, in the posterior renal angle behind, or in both places. The *referred pain* may be assigned to the healthy kidney (reno-renal reflex), testicle or labium, bladder, urethra, thigh, knee, calf, inside of foot, sole or heel, and may be of a burning nature. In *renal colic* the pain, which may be described as "tearing or gripping," is characterized by agonizing paroxysms radiating down the line of the ureter and along the urethra, or referred to the external abdominal ring and into the testicle, causing contraction of the cremaster muscle.

Oxaluria may cause unilateral pain similar to that of renal calculus. Such pain as there may be in *tuberculous kidney* is a dull lumbar ache described as toothache-like, though tuberculous debris in a ureter may cause *ureteric colic*. *Movable kidney* is characterized by a dragging ache, aggravated by movement, but the kinking of a ureter in a Dietl's crisis may produce pain resembling the colic caused by a calculus. In *kidney tumour* pain, which is not increased by movement, may result from increased intrarenal tension, from ureteral obstruction by blood clot, or from pressure on surrounding structures. *Ureteral calculi* usually give rise to symptoms akin to those of renal calculi, the pain being due to the distension and spasmodic contraction of the kidney pelvis and of the ureter above the stone rather than to the irritation produced by the stone itself. Pain of a similar character may be caused by an aberrant blood vessel, by stricture or kinking of the ureter, or by its becoming blocked by tuberculous debris, blood clot, or a villous growth of the kidney pelvis.

Vesical and Urethral Regions.

Apart from acute blocking of the ureter genito-urinary pain as a rule has its source in the bladder, prostate, or prostatic urethra. Pain originating from the upper half or more of the bladder is usually felt above the pubis, but pain from the ureteric orifices and adjacent part of the bladder wall is sometimes referred to the corresponding inguinal ring or along the corresponding side of the penis. Perineal pain suggests disease of the vesical base or neck, of the prostate, prostatic or membranous urethra, or of some structure intimately associated with one of them, constant chronic pain being in most cases due to prostatitis, or to carcinoma or calculus of the prostate, and pain increased at the end of micturition to calculus, inflammation or ulceration of the bladder; but in any of these conditions the pain may be referred to the end of the penis. Urethral pain if before micturition is usually caused by obstruction from impacted calculi, etc.; if only during micturition, by inflammation or ulceration of the urethra.

Pain due to *vesical calculus* occurs principally at the end of micturition, and may be described as burning and stinging or sharp and cutting. It is usually felt at the neck of the bladder and in the urethra slightly behind the external meatus, but may be felt in the perineum and at the root of the penis. In children it is often very severe, the patient screaming and pulling at his foreskin, especially at the end of micturition; but in old patients if the stone lies in a post-prostatic pouch, so as not to come in contact with the vesical neck, little if any pain may be felt. The pain may gradually decrease owing to the calculus becoming larger and smoother by the deposit of phosphates around it. The effect of rest upon pain due to vesical calculus is so great as to be an important diagnostic sign.

Such pain as is caused by *calculi of the prostate* is usually felt only during and after micturition, but there may be a constant ache increased by defaecation and unconnected with micturition, while in *calculi of a seminal vesicle* painful defaecation and seminal ejaculation may occur. *Acute cystitis* produces scalding pain along the urethra during micturition, while at the end of the act, or if the patient tries to hold his water, there may be pain in the region of the vesical neck. In tuberculous cases severe pain as a rule indicates ulceration of the trigone or the onset of mixed infection. The features of pain in *acute urethritis* and *prostatitis* are too familiar to need any description. *Acute spermatoecystitis* is associated

with deep pain in the perineum, pain on defaecation, and perhaps pain in the sacrum, testicle, or hip-joint. In *vesical tumour* there may be pain due to cystitis, blood clot blocking the urethra, ureteric obstruction, or pressure upon nerves. The pain may at first be unilateral, pointing to the growth being on the corresponding side of the bladder. In *carcinoma of the prostate* the pain, which is unconnected with micturition or movement, is usually a constant dull ache, but is sometimes severe and lancinating. It may be referred to the suprapubic region, urethra, perineum, hip-joints, thighs, testicles, etc., and is liable to be mistaken for sciatica.

Patients may complain of agonizing pain in the bladder, perineum, etc., and of other symptoms for which no adequate explanation can be found; but before diagnosing the condition as neurotic careful search must be made for an organic cause, and commencing spinal disease must not be forgotten.

TWO CASES OF INTRACRANIAL COMPLICATION OF ACUTE EAR DISEASE.

BY

PETER MACDONALD, M.D.,

SENIOR HONORARY SURGEON, EAR, NOSE AND THROAT DEPARTMENT,
YORK COUNTY HOSPITAL.

AND

J. ACOMB, M.B., B.S.,

ASSISTANT HONORARY SURGEON, EAR, NOSE AND THROAT DEPARTMENT,
YORK COUNTY HOSPITAL.

THE two cases described below of intracranial complication of acute otitis media were in-patients at the York County Hospital, one under the nominal care of one of us, the other of the other. The treatment, however, operative and other, as well as the process of diagnosis, was the joint work of both of us, so that they may be regarded as patients common to us jointly. They both present unusual features which seem to us to make them worth recording.

E. L., a man aged 36, was admitted on January 19th, 1922. Ten days previously he had had "influenza," and about a week previously, about January 12th, he became deaf in the right ear, and this ear began to discharge. A day or two later he became deaf in the left ear; but this ear did not discharge. There was a doubtful history of a rigor on January 18th, the day before admission.

On admission, both mastoid processes were somewhat tender; the patient was very deaf in both ears. The right drum membrane was acutely inflamed, and there was discharge from the tympanum. The left meatus was full of wax, and on removal the membrane was found somewhat inflamed, but not apparently bulging. The temperature was about 100°. There was some very slight swelling and redness of the right optic disc, and a doubtful swelling of the left disc. The patient was somewhat dull and lethargic, but did not appear very ill.

The right mastoid process was operated on the same evening. The mastoid cells were inflamed, and many of them contained granulations; but there was very little, if any, free pus. The lateral sinus was laid bare, but appeared to be healthy. The left drum membrane was freely incised.

Next day the conditions seemed to have improved, and there was but little change for a week, the temperature fluctuating between 98.4° and 100°. There was little, if any, discharge from the left ear. The general condition improved; there was less drowsiness and lethargy. The swelling of the optic discs, however, increased, and on January 23rd a haemorrhage was noticed in the neighbourhood of the left disc.

On the evening of January 26th he had a rigor, and the temperature rose to 105.8°. A diagnosis of thrombosis of the lateral sinus, which had formerly been suspected, was now definitely made, and it was decided to explore the sinus. The right sinus was suspected, there being but small evidence of any serious trouble on the left side. There was, in fact, little evidence, except deafness, of any trouble on the left side. The right sinus was accordingly laid bare at greater length. It looked healthy, but a needle was inserted into the sinus and blood aspirated from it. Blood came away freely, and it was evident that the sinus was not thrombosed, at least above the level of the jugular bulb. It was thought that the vein might be thrombosed at the bulb, and so the internal jugular was exposed in the neck, but it was quite evident that there was no thrombosis.

In view of the presence of the optic neuritis we decided, although there was no other evidence of brain abscess, to explore the temporo-sphenoidal lobe. The tegmen antri was removed, and an area of dura exposed. There was no extradural abscess. The dura appeared to be bulging. We opened it and passed a sinus forceps in several directions into the temporo-sphenoidal lobe, but no pus was found.

We were nonplussed. We had found nothing to account for the condition, and after a hurried consultation we decided to explore the left mastoid and left lateral sinus. The mastoid process was accordingly rapidly opened. The mastoid cells were inflamed and many of them filled with granulations, and on laying the sinus bare it did not look healthy. A syringe needle was inserted into the

sinus, and no blood could be aspirated from it. The jugular vein was rapidly exposed in the neck and tied above the common facial, and the clot curetted out of the sinus until blood flowed freely from its upper end. The petrosals were evidently not thrombosed, so that the upper portion of the jugular vein had to be kept tied, and we were unable to wash through from the sinus. The temperature next day was normal. It rose on January 28th, and remained between 99° and 100.5° until February 9th. It is probable that this was due to pus in the upper portion of the jugular vein and bulb into which the septic thrombus broke down.

The pathologist's report showed that the blood aspirated from the right sinus was sterile, and a culture from the thrombus on the left side contained streptococci.

The condition of the patient continued to improve, except that the optic neuritis persisted, until February 20th, when he became drowsy and lethargic, and further operation was arranged for on February 21st. But on dressing the right ear on the morning of that day a copious discharge of pus was found in the right mastoid wound, apparently from the region of the exposed dura of the temporo-sphenoidal lobe, although the wound had so closed up that the source could not be precisely determined. From this time the patient made an uninterrupted recovery, and the optic neuritis rapidly disappeared. He was discharged on April 22nd, with both wounds healed, and hearing in both ears was good. He has been seen recently, and is in good health.

The course of events can now, in retrospect, be fairly well established. An infection of both middle ears was contracted, and while there was no violent inflammation of either mastoid process, there was an extension of the inflammatory process on the right side into the temporo-sphenoidal lobe, setting up a slow abscess there, and on the left side to the lateral sinus, setting up a septic thrombosis. By opening up the right temporo-sphenoidal lobe we allowed of an ultimate escape for the pus there, although we did not actually strike the abscess, which was undoubtedly present. On the other hand, had we struck the abscess at the time of the operation on January 26th it is quite likely that we might have been satisfied that we had found enough to account for the symptoms, and might not have attacked the thrombosis of the left sinus until too late. Had the abscess not evacuated itself spontaneously on February 21st we should most likely have attacked the left temporo-sphenoidal lobe before going back again to the right in view of the presence of the neighbouring septic condition in the right sinus. As it was, we did just enough to allow of recovery, and freely admit that this was largely a matter of good fortune in what was a puzzling case.

The second was a case of middle-ear disease, complicated by infection of the mastoid antrum and cells, and sinus thrombosis. The special feature was the extent of the thrombosis and the comparative mildness of the symptoms.

H. H., male aged 19, had a history of earache from the age of 6 years with intermittent discharge from the right ear, especially after he had "contracted a cold."

He was admitted to the hospital on November 19th, 1921, suffering from a scanty discharge from the right ear, and was feeling ill and shivery. The temperature was 102°, pulse rate 84, respiration rate 20. The right tympanic membrane was incised and a discharge of serum and blood came away. The following day he felt better, but later he had a rigor, with a temperature of 105° falling, with profuse perspiration, to 99.5°. We decided to open up the mastoid and explore the sinus. The mastoid cells and antrum were found to contain pus and granulations. The sinus was pulsating freely and appeared normal. It was explored and found to contain thin pus. The internal jugular was exposed in the anterior triangle; it had become practically converted into a bag containing fluid pus. The vein was divided, but it was found impossible to reach the lower limit of the clot although the vein was followed down to the level of the sterno-clavicular articulation. The sinus was followed backwards and to the bulb; it was opened and as much of the clot and thin pus removed as possible. Neither end of the vein was tied.

Bacteriological examination of a specimen taken at the time of operation was reported as a pure culture of streptococci. Intravenous injections of polyvalent antistreptococcal serum were given. Three days after operation streptococci were reported as being present in a blood specimen taken. An autogenous vaccine was prepared and given.

The progress of the case was uneventful, and on February 1st, 1922, the patient was discharged, his condition, according to our notes, being as follows: "Looking very fit. No dressing needed for mastoid or scar in the neck."

The remarkable features appeared to us to be the mildness of the symptoms considering the extent of the disease, the absence of pain over the classical sites, the extent of the thrombosis, the absence of pyaemic complications despite the presence of fluid pus in the jugular vein and of streptococci in the general blood system, and the impossibility of removing the whole of the infected material.

Our thanks are due to our house-surgeon, Dr. Janet Nicol, for her supervision and notes of these cases.