a slight modification by fixing the tube on the cannula first, and then pushing the trocar through it. I found that when the trocar was withdrawn the slit in the elastic tubing immediately closed, and prevented anything passing through. A further experience led me to devise the instrument represented in the annexed sketches, which I have employed both in enterostomy of the small intestine and in inguinal colotomy when it was necessary to immediately open the bowel.

I have now used it sufficiently frequently to speak of its utility. After its insertion in the bowel, an ordinary antiseptic dressing is applied to the wound, and the cannula is held in position by strapping applied over the dressing. In two or three days it may be removed, as union between the parietal and visceral peritoneum will then be sufficiently firm to prevent fouling of the peritoneal cavity, but if it be thought desirable it can be retained *in situ* for several days longer, although after the third day a little moisture will escape by the side of the cannula.

In the BRITISH MEDICAL JOURNAL for July 18th, 1891, is a paper by Mr. Paul, in which he describes his method of performing inguinal colotomy in cases where it is necessary to immediately open the bowel, and in a letter to the BRITISH MEDICAL JOURNAL for September 26th, 1891, he states: "So far as I am aware no other safe method has been suggested for this purpose." Mr. Paul's suggestion of dividing the colon, inserting a glass drainage tube into its upper aperture and closing and returning within the abdomen the lower end, after Madelung's method in lumbar colotomy, has to my mind the disadvantages—first, that it involves an extensive operation in a patient ill fitted to bear it; secondly, the resection and enterorrhaphy, if required later in order to close the aperture, would involve both a serious and difficult operation; and thirdly, if the lower bowel be completely closed by the disease, and foul secretion should collect in the *cul-de-sac*, it may ultimately burst into the peritoneum, producing fatal peritonitis. That this last accident is not imaginary I have distinct proof, in that it occurred in two cases within a month where Madelung's operation had been performed by one of my colleagues.

Inguinal colotomy performed either in the way suggested by Mr. Herbert Allingham, or after the method of Mr. Harrison Cripps, and immediately opened by the method here shown, need occupy no longer than from fifteen to twenty minutes, and I have myself performed enterostomy by this means on one occasion in ten minutes, cocaine being the anæsthetic, the patient being too ill to bear general anæsthesia.

SIGMOIDOSTOMY SIMPLIFIED.

By H. A. REEVES, F.R.C.S.Ed.

Surgeon to the Hospital for Women, and to the Royal Orthopædic Hospital; Senior Assistant Surgeon to the London Hospital.

DURING the past few months I have had occasion to perform sigmoidostomy on three patients, at the Hospital for Women, for cancer of the rectum, and this I have done by an unusual, if not absolutely novel, method. Though I have not heard or read of an exactly similar proceeding, it may be that I have been anticipated, but as to this I am indifferent, my object being to attract attention to an easy plan and one applicable to nearly all cases. In some cases of obstruction where the colon is much distended it may be difficult to get out a loop of the sigmoid and, moreover, it may be necessary to open at once, but even in such cases by carefully evacuating some of the liquid faeces through a trocar the bowel tension may be sufficiently relieved to permit of withdrawal.

The first case, a woman, aged about 36, was under the care of my colleague, Dr. R. T. Smith, who asked me during the early months of the year to see the patient on account of inability to empty her rectum. A sarcomatous mass arose from the left sacro-iliac synchondrosis, the sigmoid was pressed upon, but its lumen was not completely obstructed, so that faeces passed into the rectum, but not being evacuated collected there. Atony of the rectum seemed to be caused by interference with the propagation of the peristaltic intestinal wave through the sigmoid. At this time I thought sigmoid-

ostomy not urgently needed, and directed that the rectum should be kept evacuated and laxatives given, and watched the patient. In about a month, the tumour having increased rapidly, and pain being great, and also as obstruction was increasing, I operated in the following manner.

An incision through skin, muscles, and peritoneum is to be made in the position I recommended when introducing the inguinal or iliac operation some eleven years ago. The peritoneal opening should be just large enough to allow a loop of sigmoid to rest in it without constriction. The sigmoid is pulled out, and when the descending colon is taut, or nearly so, a pressure forceps is pushed through the mesentery about a quarter of an inch from its attachment to the bowel, and a straight piece of elastic catheter, No. 10 or 12, with the stylet inside, previously cut four inches long and thoroughly cleaned, is caught in the forceps and drawn through. This is supported on the outside of the abdominal wall at each end, by a small pad of lint; a piece of green protective smeared with carbolic oil, 1 in 60, should be applied over the intestinal loop after the skin and muscle incision has been closed. A thick roll of gauze is packed around the bowel and a layer of cotton wool placed over the protective so as to prevent undue pressure when the binder was applied. It is well to give a quarter or one-third grain of morphine immediately after the operation to check anæsthetic vomiting, or to let a nurse sit by the patient and gently press over the site of operation during vomiting, but if care be taken that the peritoneal opening be of the right size, there is no fear of the small intestine being protruded and nipped by the side of the sigmoid. However, if in any case vomiting be prolonged it is safest to explore the incision rather than be beguiled into taking the symptoms as due to prolonged anæsthetic vomiting.

At the end of a week in this case I opened the bowel longitudinally and stitched the gut edges to the skin, but have not done this in later operations. Care must be taken not to form a twist in the gut, as, apart from some risk of obstruction, faeces will pass from the lower (which is really the upper) opening, and, unless the condition be recognised, may lead to the erroneous view that there is regurgitation from the rectum. I have lately used polished vulcanite rods, which are dipped in carbolic oil, instead of the catheter.¹ They are pointed at one end to pass through the mesentery, or bevelled off as in the adjoining figure. Should the perito-

Diagram of sigmoid loop at the time of passing the rod. 1, Site of incision in bowel; 2, mesentery; 3, rod. The figure to the right shows the bowel opened. 1, Upper opening, not represented large enough; 2, lower opening; 3, rod and spur of mesenteric side of bowel over it.

neum be tough a nick may be made over the end of the rod, and, before passing it, the operator should see that no vessel is in the way. If necessary, the bowel can be opened at the end of the second day, but, unless there be urgency, I prefer to wait till the third or fourth, and I leave the rod in place two or three days longer, as it appears to form a better spur the longer it is left.

The subsequent behaviour of the bowel is interesting. In this case—and this seems to be the rule—the loop be-

¹ These are made by Messrs. Mayer and Meltzer.

came congested, and its serous coat took on a pink and then a red velvety aspect, and became granular. The coats of the bowel also became thickened, and this could be distinctly recognised at the time of opening the gut when a transverse piece was taken between the finger and thumb prior to opening with scissors. In two of the cases the enlarged vessels of the bowel wall had to be tied when the gut was opened. After withdrawal of the rod the bowel is gradually drawn back by the mesentery and by its own weight as far as it is permitted by the adhesions formed between its mesentery and serous coat and the parietal peritoneum, or rather the muscles and skin, for the parietal peritoneum retracts, and is not in contact with it to any large extent. In from ten days to a fortnight no one would know what kind of operation had been done. Personally I think the result is neater, and that the feel of the spur would allow me to state the method of the operation. The simplicity of the proceeding is such that the dexterous surgeon may execute it in five minutes, and I have done it in seven in a thin woman in whom there was no delay in tying vessels and whose sigmoid mesentery was neither too short nor too long, but, if the latter condition exist, time is spent in finding the bowel, and I have had one such case by this method and two or three by the plan of opening the bowel at once, which I originally adopted. This patient recovered well from the operation with the slight exception of retention of urine, which I have not noted in other cases. She subsequently went into a workhouse, but before leaving hospital a growth on her sternum was noticed, and I heard that she died about three months after with large sarcomatous growths on her skull and elsewhere.

For notes of the following cases I am indebted to Mr. Ernest Brock, House-Physician to the Hospital for Women:

Mrs. H. P., aged 42, admitted September 8th, complaining of pain in the back. Pain in the abdomen, with frequent desire to go to stool, had come on eighteen months earlier. The motions were jelly-like, and she had lost flesh. About 4½ inches up the rectum the finger met with a hard, more or less nodular, irregular mass, almost blocking up the lumen of the bowel. It was very painful, but there was no hæmorrhage on examination. As it felt pedunculated and movable an attempt was made to pull it down under chloroform, and pass an *écraseur* round its base. As this could not be done, consent for colotomy or posterior excision was awaited. The general condition of the patient was fairly good. The temperature and urine were normal. On October 8th sigmoidostomy was performed, as obstruction was threatening. An incision was made just above and internal to the anterior superior spine, and extending downwards and inwards for about 2½ inches. The peritoneum was cut through on a director, and the bowel easily reached and drawn out. No "slack" could be obtained. The bowel was found to be full of rather hard motion, and an attempt was made to squeeze this into the rectum, but without success. The mesentery was then perforated with pressure forceps, and a gum elastic catheter No. 10 was drawn through and held the bowel in position on the abdominal wall. The upper and lower ends of the incision were then stitched up with silk. The whole was then covered with oiled protective and wool. On October 12th the bowel was opened with scissors in a line parallel with the gut. Flatus readily escaped. On October 14th the catheter and stitches were removed. On October 17th the bowels had acted well after a single enema. No feces or matter had been passed by the rectum. In a few days the patient left hospital, having recovered from the operation.

Mrs. E. K., aged 35, was admitted September 28th, complaining of "pressure" in the back passage, with a feeling of weakness. "Six months ago she fell down, striking her fundament." Ever since then she had had more or less pain in defecation; for the last two or three months she had had almost continuous slimy discharge from the rectum, and until ten days ago she had passed blood with almost every motion. She had no actual pain, but great sense of pressure. She had been losing flesh for twelve months. She was fairly well nourished, slightly anæmic, but otherwise looked in good health. About an inch within the anus the finger met with a very rugged indurated mass, extending about four-fifths round the gut, the free part being anterior; it extended upwards for about 1½ inch, the finger readily feeling healthy mucous membrane beyond. There was no enlargement of inguinal glands. The temperature and urine were normal.

Posterior excision was proposed and its risks pointed out. The patient preferred colotomy, as the nature of the growth was not made known to her. On October 1st an incision was made in the left inguinal region, about 2½ inches long, commencing just above and internal to the anterior superior spine, and extending downwards and forwards. The layers of the abdominal wall were cut through, and the peritoneum opened on a director, and the bowel easily reached; the latter was drawn out of the wound as far as possible, but no "slack" could be obtained; the mesentery was then perforated with a pressure forceps, and a gum elastic catheter drawn through to support the bowel. The upper and lower ends of the abdominal incision were then stitched up with silk, and the whole covered with a piece of protective and wool. Hæmorrhage was only trifling.

On October 8th the bowel was opened with scissors in a line parallel with the gut. Flatus readily escaped. The catheter and stitches were removed. The bowel protruded from abdominal incision about 1 inch.

By October 15th the bowel had fallen back to some extent. The artificial opening acted well almost every day, and no discharge had occurred from the rectum. The patient was discharged a week later.

The second case is of interest to me, because at the first

examination I was inclined to think that it was one of those remarkable coincidences which we meet with now and again in hospital and private practice. Shortly before this patient's admission, and within a week of each other, two women were admitted with broadly attached pedunculated growths high up the rectum, on the left side, which bulged externally or near the anal orifice during defæcation, and thus caused some obstruction and gave rise to much glairy mucous discharge. From their appearance I took them to be broken-down adenomatous growths; but microscopic examination by Dr. Dalton revealed their malignant nature. At the operation they were easily pulled outside the anus, but there was a long invagination of the upper part of the rectum and lower part of the sigmoid, so that I did not put a snare round the base of the growths but removed them with scissors, scraped their points of attachment, then cauterised, and finally applied a matico plug. They recovered rapidly.

The efficacy and extreme simplicity of the operation described, and also of that I adopted when first drawing professional attention to the advantages of inguinal colotomy, must be my excuse for saying that I entirely fail to appreciate the various modifications which have been introduced. What need can there be for prolonged and serious proceedings, such as cutting away some inches of colon, or of lengthening the operation by stitching the gut to the parietal peritoneum in various ways, or of dividing the bowel and invaginating the lower end according to Madelung? The double-barrel stitching and subsequent excision of part of the bowel as done by Maydl is a lengthy proceeding, and the method of Finc, of Geneva, recently revived, of opening the transverse colon is going unnecessarily high up the bowel. All the advantages claimed for these proceedings can be obtained by the simple plan herein recommended, and all their dangers and drawbacks obviated. I would urge surgeons to remember the fact that in opening the colon for malignant disease not amenable to a more radical proceeding, such as excision, the object is to relieve pain and prolong life, whether done for or without obstruction, and that any operation of a palliative kind, such as colotomy, should perforce be of a comparatively innocent nature. The simplification of operative measures should, in the interests of patients, be the aim of the surgeon, and experience abundantly proves that the simpler an operation is the better are its results.

Since writing the above I have operated on a fourth case in a man at the London Hospital, who recovered rapidly.

MEMORANDA: MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

A CASE OF PERFORATING ULCER OF THE DUODENUM.

THE following case, in conjunction with the two cases related by Mr. C. B. Lockwood, may be interesting to some.

E. J., aged 25 years, married, with one child, had always had very good health, with the exception of having had slight attacks of indigestion, about which, however, she had never had medical advice. On May 16th, 1890 (when away from home acting temporarily as housemaid), after tea, consisting chiefly of new bread and butter, she had an attack of severe abdominal pain, and felt sick, but the attack passed away in a little time, and on May 17th she had a little pain after food, but continued her work, and on May 18th drove home a distance of three miles, and walked about all day and back in the evening to her situation. On May 19th, after tea, the pain became very severe, and she vomited a little food, not mixed with blood, and nearly fainted; she was helped to bed, when she continued in great pain all night, vomiting several times some bilious fluid, which was not stercoraceous. The bowels had acted once naturally, on May 18th; and she had ceased to menstruate the previous Saturday week, having been regular up to that time. When seen for the first time on May 20th, at 10 A.M., she was lying in bed with sunken eyes and a very curious expression, and her knees and thighs flexed, com-