drainage tube, and as a rule healing by first intention is readily obtained. I have performed this operation repeatedly during the last seven or eight years, with comparative ease and with relief to the patient for the time being; but I must also confess that in by far the majority of the cases the pain recurred at the end of a year or two, necessitating those further measures which I shall detail in my next lecture.

[This lecture was illustrated by diagrams representing the position of the different divisions of the nerve, as given by Mac Cormac and others, and also by drawings of the different operative measures alluded to.]

FURTHER OBSERVATIONS ON QUARANTINE.¹

By HENRI MONOD,

Directeur de l'Assistance et de l'Hygiène Publiques, Ministère de l'Intérieur, République Française.

WITH your permission I will leave on one side matters which are non-essential. That the illustrious Sir John Simon condemned quarantine in 1866 is a matter of small importance, since one of the official delegates of Great Britain proposed its recognition by the Conference at Vienna. Was quarantine abolished at British ports in 1866? The answer is in the negative, since Sir John Simon protested against it, and one of his arguments might have been that that very year it had failed to prevent cholera from claiming nearly 14,000 victims.² An occasion for employing quarantine against cholera—at least, against cholera prevalent in Europe—can hardly be said to have again presented itself until 1884, and by that time sanitary work was already far advanced in Great Britain. I am not then very far wrong when I say that Great Britain defended itself against cholera by means of quarantine until the time came when she found a much more complete guarantee in “sanitation.”

What difference does it make whether cholera has or has not been brought to Europe by an English ship? Will it be denied that cholera is endemic in India? Will it be denied that it is transmissible, that the lethal germ can be transported by sea? Will it be maintained that, supposing the disease to break out on board an English ship, the medical service and disinfecting methods at sea are so well organised that the infection is certainly destroyed before reaching the Mediterranean? If the answer to all these questions must be in the negative, what does it matter whether the disease is carried from India to Europe in an English or in a Turkish vessel? It is possible that it may travel under the flag of a foreign nation, but it is not impossible that it may travel under the British flag.³ That is enough for the purpose of my argument.

It is unnecessary to revert to the excessive quarantine imposed by the British possessions of Malta, Cyprus, Gibraltar. I have not said that England was responsible. I have said that England would be in a better position for convincing foreign nations after she had succeeded in convincing her own subjects. Is not that a very reasonable proposition? Distance is a natural protection to England. The quarantine which protects her is the length of time which it takes to reach her ports, and it is interesting to observe that the British colonies in the Mediterranean, finding themselves under the same conditions as the Mediterranean nations, act,

¹ In continuation of a correspondence published in the *BRITISH MEDICAL JOURNAL* on October 10th, 1891, and in reply to an editorial article published on the same date.

² 13,553. Farr, *Vital Statistics*, p. 389.

³ I need not recall that in 1832 cholera reached France from England via Calais: that in 1865 it was an English ship, the *Sydney*, which carried it to Egypt, and that from there it exploded over Europe; that in 1891 it was an English ship, the *Deccan*, which introduced it into Camaran, from whence it spread to Jeddah and to Mecca.

with regard to quarantine, not only like them but worse than they.

Let us pass on to points of more practical importance. The progress of sanitary administration in India appears to you, it would seem, to be satisfactory, and you express regret that I have never glanced at the reports published regularly by that administration, since I could not but have been struck by the results attained, notably in the diminution of mortality. In truth, these documents deserve more than a glance.⁴ How, if I had not studied the last official report on the sanitary measures taken in India, could I know that “there is not room for much congratulation as to the improvement of the sanitary condition;”⁵ that “the localities are progressively attaining a more unhealthy state;”⁶ that “the foul state of certain districts defies all description”⁷? Where, if not in this report, could I have learnt that the vital statistics published in India, such as are quoted against me, deserve very little confidence, and that the diminution of mortality, when it occurs, “is due to imperfect registration rather than to an improvement in the public health”⁸? What other document could have given me with equal authority the number of deaths from cholera in India during recent years, and have enabled me to prove with equal certainty that sanitation in India had not as yet seriously influenced the mortality from cholera?⁹

Let it not be said that my quotations are picked, detached from their context, faithful to the letter but not to the spirit. The general impression to be gathered from these reports, which are precise and honest, is, in truth, that which my quotations give, and matters have not changed much in India since Dr. Furnell, in the *Indian Medical Gazette*, April, 1886, wrote that there was not “a hamlet, not a village, which was not supplied with water liable to the worst forms of pollution.”¹⁰ It was only the other day, at the International Congress in London, that Dr. Pringle said that in India “95 per cent. of the population live in the villages, and that the cisterns of these villages are also the water-closets.” Examine the descriptions given at this Congress in London by Mr. Ollivant of Bombay, by Dr. Holbein Hendley, by Mr. Baldwin Latham, and then, perhaps, you will be disposed to say with the last named: “The sanitary statistics of Bombay and Calcutta prove that it is not the climate which causes the terrible mortality of a portion of India, but the total absence of the simplest hygienic precautions.”¹¹

It will not be denied that cholera is one of those preventable diseases to which reference is made, nor that it is among those which can be most certainly controlled by sanitary measures. Even in India more than one example might be

⁴ I have requested the India Office to exchange its publications for those of the Direction de l'Hygiène Publique en France. It has been good enough to inform me, under date September 26th, 1891, that my “request is under official consideration.”

⁵ “Decline in fever mortality is not so marked as to justify much congratulation on the improvement of the public health.” Dr. Simpson, health officer of Calcutta. *Report on Sanitary Measures in India in 1888-89*, p. 122.

⁶ “Each year's delay in carrying out the constitution of broad streets brings with it greater difficulties and expenses; the crowding together of houses is becoming greater, the open spaces are being more encroached upon, the breathing space is gradually contracting, and the localities are progressively attaining a more unhealthy state.” Dr. Simpson (*ibid.*, p. 123).

⁷ “The most graphic description of the foul state of some districts will not give as good an idea of the truth as a quarter of an hour's visit.” Mr. E. C. K. Ollivant, Municipal Commissioner of Bombay (*ibid.*, p. 127). “Defies all description” is not too strong for this is what the Deputy Sanitary Commissioner of Madras writes as to the town of Tirupati: “There are no public latrines. Portions of back yards are set apart for private resort, holes being left in the walls so that pigs may enter and clean up. People defecate anywhere, and the whole town is pervaded with the odour of pig's dung and human excreta” (*ibid.*, 104).

⁸ “No less than forty-three of the forty-nine districts of the province (North-Western and Oudh) shared in this reduced mortality. As death registration is imperfectly conducted, it is impossible to decide how much of the reduction may fairly be attributed to better general health; probably the greater part must be owing to careless recording..... This decrease (in the Punjab), it is feared, must be more largely attributed to imperfect registration than to improved health conditions” (Sir B. Simpson, Sanitary Commissioner with the Government of India, *ibid.*, pp. 78, 80). Sir B. Simpson expresses himself in almost the same terms on the registration of deaths in each of the provinces of India.

⁹ The following are the numbers:—1878: 318,228 deaths from cholera; 1879: 269,336; 1880: 118,856; 1881: 161,712; 1882: 350,971; 1883: 248,860; 1884: 287,600; 1885: 385,928; 1886: 208,371; 1887: 488,788; 1888: 270,408 (*Report on Sanitary Measures in India in 1888-89*, p. 80).

¹⁰ I have not the English text; I quote from the translation in the *Revue d'Hygiène* (Août, Septembre, 1891).

found which would prove this. How is it, then, if the work of sanitation is carried on with so much success, that deaths from cholera continue to be counted by hundreds of thousands? How is it that there are no indications of any diminution in the terrible mortality? Let us compare the curve of mortality¹¹ from cholera in India with that of the general mortality¹¹ in England for the years 1878 to 1888. They are shown in the accompanying charts.

Chart showing number of deaths from cholera in British India (Report on Sanitary Measures in India in 1888-9, p. 80).

Ah! here, indeed, a glance is sufficient. From the comparison I will not conclude "sharply" (sic) that sanitary matters are attended to in England and neglected in India. I will conclude that much has been done for sanitation in England, but that in India not enough has been done to resist cholera successfully.

¹¹ (Note by translator) : M. Monod uses the word *mortalité* in both places

The second hope which I expressed in my letter, but to which no reference has been made, is that until cholera has been destroyed in its cradle the local administration should shut it out from the Indian ports and prevent its exportation. I admit that the Native Passenger Ships Act is an important advance. To have made it compulsory to have a medical service on board every ship carrying more than 100 pilgrims is a considerable advance (Section 26). It is something, also, to have given the Government the power to examine the pas-

Chart showing general death-rate of England (Fifty-second Report of the Registrar-General, p. lxxvi).

sengers as to their sanitary condition, and to remove those recognised to be suffering from an infectious disease. (Section 30). I will make only two observations. Why is this medical inspection permissive and not compulsory? If the Government can order, it can refrain from ordering. It will not order except when it can do nothing else; that is to

say, only when cholera has become epidemic in the port affected. Is this sufficient? Certainly not. Is it essential that cholera should be epidemic in Bombay in order that pilgrims embarking in Bombay should carry cholera with them? Numerous examples, of which one is quite recent,¹² prove the contrary. It would be an event of the first importance if it were decided that no pilgrim, under any circumstances, could embark without obtaining a sanitary passport. This is my first criticism. The second is as follows: Cholera can be transmitted not only by persons but also by things. The things likely to carry the infectious germs are especially those in common use. Now these things can be sterilised and rendered harmless without being damaged, by being placed in a steam-pressure stove; this has been proved scientifically by experiments made at the Exposition Universelle, in 1889. At the present time thirty-four of your large ships are provided with these stoves. To erect such apparatus in the principal ports of Hindostan would not cost much, and would be a great advantage. If no pilgrim could leave India without having been seen by a physician, if in every case the effects of pilgrims were passed through the disinfecting oven, there would be, in reality, every chance that they would no more carry cholera into the Red Sea.

I say "no more," and this brings me to my third proposition. In my previous communication I said that if, in spite of all, the fatal germ escapes, a grave responsibility is taken if opposition is made to measures designed to destroy it before it has sown death upon the European continent. The answer I receive is, There ought not to be quarantine under any name by which it is concealed; quarantine has never prevented the spread of anything. That the measures ought not to be quarantine I am quite willing to allow, and I have already let it be understood that there are better things to be done; medical inspection and disinfection, well performed, are, in my opinion, very much better guarantees than the old quarantine system. But to say that quarantine has never prevented anything is too absolute an assertion. How many facts can be invoked on the other side? I will quote two; in the first, the absence of quarantine measures was the cause of a terrible epidemic; in the second, evil appears to have been checked by a severe quarantine. In 1865 an English ship, the *Sydney*, arrived at Suez. Owing to an incorrect declaration by the captain who concealed the fact that deaths from cholera had occurred on board, she was allowed to pass. No quarantine. She arrived at Alexandria and disembarked passengers. Cholera broke out, increased, and caused 60,000 deaths in Egypt. In 1891 an English ship, the *Deccan*, arrived at Camaran with more than 1,200 pilgrims. Cholera was reported on board this ship. The pilgrims went to Jeddah. Cholera accompanied them and prevailed severely in Jeddah and Mecca. On August 8th, the ship *Adana* left the port of Jeddah for El Tor with 971 pilgrims. During the voyage 18 deaths occurred from cholera, 6 occurred during disembarkment at Jebel-Tor. Quarantine was imposed there. More than 10,600 pilgrims underwent a most rigorous quarantine. The scourge was stopped at the entrance to the Canal; Egypt and Europe were preserved from it. A very robust belief in the inefficacy of quarantine is needed to maintain that quarantine was useless in this case. Would it have been better to have allowed cholera to have freely entered the Mediterranean, and ought nothing to have been learnt from the lessons of 1865 and of 1884?

I fear that here again we shall not be in agreement, and that you will think that I am right only when I admire. I would regret this very much, for in order to destroy cholera it is necessary for us to have not only the active but the energetic co-operation of England, and a journal such as the BRITISH MEDICAL JOURNAL which has so much legitimate influence on public opinion could do much to obtain that assistance. I hope at least that it will be believed that in this discussion I am inspired by no other sentiment than a desire to ascertain the truth, and to serve the public weal.

¹² I allude to the epidemic which broke out in September on board the English ships, the *Marathon* and the *Redbreast*. As a general rule, of the victims of cholera in Bombay few are inhabitants of the city. "Of the total cases of cholera, 84.3 per cent. were strangers," Dr. T. S. Weir, health officer of Bombay. (*Report on Sanitary Measures in India in 1888-89*, p. 132.)

REMARKS ON AN OPERATION FOR ABSCESS OF THE LIVER.

By PATRICK MANSON, M.D., M.R.C.P., LL.D.

A good many years ago I was asked to see a man said to be suffering from intermittent fever. After examining him carefully and hearing his story, I concluded he was suffering not from intermittent fever but from abscess of the liver. In addition to other symptoms, there was distinct enlargement of that organ; but, owing to the absence of localised swelling or bulging or localised pain or tenderness, I could not say positively what was the exact situation of the pus, if pus were present. I determined to aspirate, partly with the view of clearing up diagnosis, and partly in the hope that if I found pus and emptied the abscess with the aspirator I might cure the patient. At the time of which I speak the aspirator was a new instrument and much was expected of it—cure of liver abscess among other things.

In the absence of definite guide from either swelling or pain or other localising sign, I elected to enter the needle at a point just below the ribs and about midway between the ensiform cartilage and the nipple line. The abscess was not reached until the needle had been driven in to its full extent—that is, about 4 inches. Some 10 ounces of pus were withdrawn. For a day or two the patient improved, but hectic recurring, I had to repeat the operation, and eventually the man died from exhaustion and hectic.

At the *post-mortem* examination an abscess of very moderate dimensions, with somewhat thick and uncollapsing walls, was found deep in the liver, well towards the upper and back part of the right lobe. The rest of the gland was healthy, and there seemed to be no reason, could this abscess have been opened and drained in an efficient manner, why the case should not have terminated in recovery.

Until recent years the operative treatment of abscess of the liver had proved so unsatisfactory that not a few physicians of experience—Martin and Budd, for example—deemed it right in all such cases to abstain altogether from recommending active surgical interference. They considered—and with a fair amount of statistical support—that a patient had a better chance of recovery if the abscess were left to itself than if it were opened with knife, trocar, or caustic; they considered that the chance of recovery by waiting for possible rupture of the abscess externally, or through the lungs, or into the alimentary canal, was distinctly greater than any offered by operation. Whether operated on or left alone, the mortality from suppurative hepatitis in pre-Listerian times was about the same, and was set down at something between 80 and 90 per cent. Since the introduction of aseptic methods, combined with a bolder system of treating the abdominal cavity, the position of Martin and Budd and their followers is no longer tenable. Surgery has reduced the mortality from this disease to something between 15 and 50 per cent., and it is now fully recognised that the rules which guide treatment of abscess elsewhere must be rigidly observed in treatment of abscess in the liver.

It is one thing, however, to recognise rules and principles, it is often quite another thing to apply and carry them out; and unfortunately, in the case of the liver, there are anatomical conditions which make the efficient application of these rules and principles peculiarly difficult. Suppose that the initial difficulty of diagnosis, often very great, has been successfully overcome, and suppose that the aspirator has demonstrated the presence of pus at the upper part of the right lobe of the liver, a very usual situation; and suppose the abscess is found to lie some three or four inches from the surface, how are we to treat it so as thoroughly to empty it and provide efficient drainage? The direct road to an abscess in this situation is through the thoracic wall, pleura, lung and diaphragm—an impracticable route in the early stage of liver