bers pouring out three drops carefully, but after that her hand shook, and she saw a quantity of the draught run into the measure glass. How much really ran into the glass she does not know, but subsequent examination of the bottle makes it unlikely that more than twenty drops in all had been poured out. However, believing that her calculation was approximately correct (that is, that she had measured off about ten drops), in which estimate it is probable she was fairly accurate, she added water to the quantity in the glass, and drank it off. The friends, hearing her moving about, returned to the bedroom, and learned what had occurred. Almost at once she began to talk nonsense as before, but now more rapidly and incoherently. After a time this behaviour grew so alarming that I was summoned.

I saw the patient at 12.45 A.M., about two hours after she took the last dose. She was, of course, still in bed. The pulse was small, 92 per minute. The pupils were dilated and stationary, not reacting to light. The respiration was sighing, and general depression was very marked. The face was not flushed, and, though the patient was delirious, she was not noisy, nor was there causeless laughter, as in poisoning by atropine.

By way of treatment I gave at once $\frac{1}{4}$ grain of morphine sulphate by means of the hypodermic syringe. At 1 A.M. the pupils had begun to react to light slightly, and the patient was then just asleep. At 2 A.M. the patient was sleeping quietly, but was easily roused. The pupils responded to light a little more markedly than they did an hour previously. Complaint was made of dryness in the throat.

Next day the patient was almost quite well again, but the pupils were still dilated, and the vision yet imperfect. During the next few days the recovery was rapid and complete, but fully three days elapsed before the pupils and vision had returned to their normal condition.

AN OUTBREAK OF ENTERIC FEVER APPARENTLY TRACED TO AN ANTECEDENT CASE AFTER AN INTERVAL OF TWELVE MONTHS.¹

By G. REID, M.D.,

Medical Officer of Health to the Staffordshire County Council.

THE following summary of the circumstances attending an outbreak of enteric fever is the outcome of an inquiry, conducted on behalf of the Staffordshire County Council, in which I was assisted by Dr. Hosegood, of Cannock, the medical officer of health of the district in which the cases occurred. The interesting feature of the outbreak, which occurred in the autumn of 1890, is the long interval that apparently elapsed between the original introduction of the specific contagion into the district and its second manifestation.

The cases in question—seven in number—occurred within a period of two months in various houses in different parts of the parish of Church Eaton, and, with one exception—the last case that occurred—all the patients were, in one way or another, in daily communication with a farm house (the Church Farm) in which Cases iv, v, and vi occurred. The following are the circumstances attending each case:

CASE I.—A child, living a quarter of a mile away from the village, in whom the first symptoms appeared about August 20th. The mother of this child was laundrywoman at the Church Farm, and went daily to the farm, taking the child with her. Both mother and child were in the habit of taking meals at the farm, and drinking the well water there.

CASE II.—A child living in a cottage near to the Church Farm, in whom the fever showed itself about the same time. For some months previously the well belonging to this cottage had been dry, and during that time the family had been in the habit of sending for water to the Church Farm well.

CASE III.—A young man, living a quarter of a mile away from the village, who worked at the Church Farm, and who habitually took his meals with him to the farm, and drank the well water there. This case occurred towards the end of September.

CASE IV.—A youth who lodged at the Church Farm and who was attacked a day or two after the preceding case.

CASE V.—A child, the grandson of the tenant of the Church Farm, who lived at the farm, and who was attacked immediately after the previous case.

CASE VI.—A young man who worked on the Church Farm and lived in the house. This case followed the previous one very closely.

CASE VII.—A child living in a cottage in the village, some distance from the Church Farm. This was the only case that occurred in which the contagion could not be traced directly to the farm.

The farmhouse in question was not directly connected with the drains, so that the cause of the fever could not be traced to within the house. The outside arrangements corresponded with those generally found in farm buildings. The well, a comparatively deep one, sunk in the gravel and built in loose stonework (not set in cement), was close to the back door; it was covered over and had a pump attached. Within eight yards was the proverbial privy, with the large, foul, leaking cesspit, and, close by, a pigsty, the drainage from which lay on the surface. The house slop water was thrown on to an open channel on the paved yard close to the pump, and discharged into a catchpit, about five or six yards off, into which the manure heap drained, and from which there was an overflow pipe communicating with a cesspool some distance off.

The premises adjoined the churchyard, which, however, at its nearest point was 30 yards distant from the well and on a lower level. With regard to the quality of the Church Farm well water, an analysis had recently been obtained and it was pronounced to be good and fit for domestic use. As, however, no safe chemical standard of purity can be laid down, in order to ascertain whether the quantity of organic matter present corresponded with that in adjoining well waters, I analysed three samples and found that there was an excess both of free and albuminoid ammonia and also of chlorine in the Church Farm well water as compared with the other two, additions which could only have come from an impure source, namely, soakage from the privy or other collections of filth on the surface.

So far, then, I think we have conclusively traced six of the seven cases to the well water at the Church Farm; for, although three of the cases occurred in the farmhouse itself, and it might therefore be said that two of them had resulted from carelessness in dealing with the discharges of the other, this is contraindicated by the fact that only a few days had elapsed between the appearance of the disease in the three patients.

From whence, then, was the specific contagion introduced into the well? It was said that some years previously cases of "low fever" had occurred from time to time in the village, but for four years, with one exception, no illness of that nature had been heard of. To this exception it would seem the cause of the outbreak in question is to be attributed. In the autumn of 1890, twelve months previous to the subsequent outbreak, two cases of enteric fever occurred, one (imported) in the Church Farmhouse, and the other in a family whose milk supply came from the farm. From this time until the occurrence of the cases just reported, no case of enteric fever had been heard of in the neighbourhood.

To summarise the facts just related, After an interval of four years, during which time no cases of enteric fever had been heard of in the neighbourhood, a case was imported into the village, and gave rise, in all probability, to a second case. Twelve months then elapsed, and seven cases occurred in five different houses within a period of two months, six of these, if not the seventh, being directly connected with the farmhouse in which the imported case occurred a year before.

Was the second outbreak connected with the first, and, if so, why the long interval between the two? Had the interval between the original case and the second outbreak been shorter, one could easily have understood the connection, considering the proximity of the well to the privy cesspit, into which, no doubt, the discharges from the first case had been thrown. As, however, twelve months elapsed between the two, if we are to look upon the second outbreak as the outcome of the first, the contagion in the interval must either have lain dormant, or it must have taken that time to travel from the cesspit to the well, which is hardly likely, seeing that the one was so close to the other, and that the daily consumption of water must have been considerable.

As regards the outbreak in question, I venture to suggest that its sudden occurrence was the result of renewed vitality and not of evolution; that, in fact, the specific virus remained for twelve months in a dormant state until conditions favourable to its development were present. In any case, the out-

¹ Read at a meeting of the Staffordshire Branch of the British Medical Association.

break is an instructive one, as showing how impossible it is to arrive at a conclusion regarding the safety of a water supply upon chemical evidence only.

HEPATOTOMY AS A TREATMENT FOR HEPATIC ABSCESS.

By FRANCIS T. HEUSTON, M.D., F.R.C.S.I.,

Surgeon to the Adelaide Hospital; Consulting Surgeon to the Coombe Hospital; Professor of Anatomy, Royal College of Surgeons.

THE very interesting and instructive papers on the surgical treatment of hepatic abscess, appearing in recent numbers of the *BRITISH MEDICAL JOURNAL*, induce me to bring forward the notes of a case which lately came under my observation, as I am strongly of the opinion that the fear expressed by most writers on this subject as to hæmorrhage from the liver, and the danger of pus escaping into the peritoneal cavity is much overrated, as shown by recent abdominal surgery.

I can readily understand that if a liver abscess be evacuated by an aspirator, pus may escape by the side of the needle into the peritoneum. Or that, as the sac fills again and the tension becomes great, the pus will follow the track of the needle, and having no point of exit be forced into the peritoneal cavity; but in cases where a free incision is made in the first instance and kept patent while pus continues to form, there can be little or no danger of the pus passing into the peritoneal cavity even when no adhesions exist between the parietal peritoneum and that covering the liver, as the normal contraction of the abdominal muscles is such as to hinder extravasation, as shown in the case of the second abscess I operated on.

A. P., aged 27 years, a clerk, booked with the Imperial British East African Company as an assistant, in December, 1890, and was sent to Mombaza, 140 miles north of Zanzibar. Soon after arrival he was attacked by remittent fever; and after a residence of four months, was sent to Aden to recruit his health. On the voyage, the weather being very hot, he slept without bed clothes, and on awaking in the morning felt chilly, and had a stitch-like pain in his right side; this, however, shortly disappeared, giving place to a heavy, dull, persistent pain in the right side. After a short stay in Aden he returned to Mombaza in much better health. About six weeks later the fever again attacked him, which necessitated his return home in September, 1890. On the voyage home he so far recovered his health that on his return he was appointed clerk on the Great Southern and Western Railway, which post he soon relinquished owing to a recurrence of his illness. I saw him on December 3rd, 1891, and took the following notes of his condition. The complexion was sallow, the pulse 112, regular and compressible, the tongue coated with thick white fur; he was losing flesh rapidly, and for the past three weeks vomited large quantities of a light green fluid, solid food caused pain in the epigastrium. The bowels were regular, with light coloured motions; he had a feeling of numbness and tingling in the left leg, corresponding to the cutaneous distribution of the musculo-cutaneous nerve; sensibility was diminished. His liver was greatly enlarged, giving rise to bulging out of the ribs and fullness in the epigastrium. He complained of a dull aching pain in the hepatic and right iliac regions, also of tenderness below the costal arch, chiefly in the epigastric region; the spleen was slightly enlarged; flatus passed frequently by the mouth and rectum; there was constant nausea and occasional mucous vomiting; the breath was very offensive. The urine was clear, high coloured, acid, of specific gravity 1021, and contained a trace of albumen and bile pigments, but no sugar; the chlorides were abundant.

The patient was kept under observation until December 8th. The temperature varied between 100° F. and 101° F. in the morning, and 102° F. to 103° F. in the evening; his liver increased rapidly in size, a distinct enlargement occurring in the epigastric region to the right of the median line.

Operation being determined on, an incision 3 inches in length was made parallel to the right costal arch in the epigastric region, commencing opposite the seventh costal cartilage, and the liver, which was pale in colour and not adherent to the abdominal wall, exposed. A small aspirator needle was passed into the liver, pus being found at once, a little oozing out beside the needle; 20 ounces of a chocolate coloured pus were at once removed by the aspirator. Tension being now relieved in the abscess, a free incision was made through the walls of the sac, and about three pints of pus removed, the opening in the liver being held by forceps in close contact with the wound in the abdominal wall. The sac was now washed out with boric acid solution, and sutured to the abdominal wall by two rows of gut, the first attaching the liver half an inch from the opening in the abscess sac to the parietal peritoneum and deep muscles, the second attaching the margin of the opening to the incision in the skin. Two large drainage tubes were introduced to keep the opening patent, and the whole dressed with zinc sulphate gauze.

The condition of the patient improved rapidly after the operation: the temperature varied from 99° F. in the morning to 101° F. in the evening; nourishment was freely taken. For the first few days there was a very free discharge of pus; the sac was washed out daily with boric acid solution, and the amount of discharge rapidly became less. On December 18th pain was again complained of in the epigastrium; his temperature rose to 102°, and by December 21st a slight tumour was observed in the

epigastrium to the left of the median line. His symptoms becoming aggravated, an aspirator needle was passed into the tumour on December 23rd and pus found. I now made an incision to the left of the median line, exposing the left lobe of the liver, which I found partially adherent to the abdominal parietes. On incising the liver I removed about a pint of pus, but I found that when I attempted to suture the liver to the abdominal wall the stitches tore through the liver substance. So a strip of zinc sulphate gauze was introduced, and pressure applied by a pad to the abdomen below the opening. At the time of this operation the abscess opened into previously had almost ceased to discharge pus, and was practically well. The abscess opened into now made an equally favourable and rapid recovery under daily dressings as before, but the patient's strength gradually failed, and he died from asthenia on December 31st, eight days after the second operation.

Post-mortem Examination.—The liver was adherent to the abdominal wall at the sites of the operations, completely closing off communications with the peritoneal cavity. It weighed 110 ounces; the abscess cavity operated on in the right lobe was represented by a fissure about two inches in depth and lined by a distinct membrane and contained no pus. The abscess cavity in the left lobe was capable of containing about 5j of fluid. Another large abscess was found in the posterior portion of each lobe filled with pus; those were entirely out of reach of operative interference. The spleen weighed 3xij; the kidneys were normal; there was no evidence of peritonitis.

PERINEAL DRAINAGE IN SUPRAPUBIC LITHOTOMY.

By H. HERBERT, F.R.C.S.ENG.,
Surgeon Indian Medical Service.

VERY little reference is made in British textbooks to this mode of drainage in suprapubic operations, still less to the differentiation of cases requiring it. And for this two reasons are perhaps accountable. First, it is really very seldom needed in civilised countries, where cases of stone usually come under treatment at an early stage. And, secondly, if the drainage by this route is not perfect, it is practically useless. Drainage by the urethra is too uncertain to be employed with any satisfaction. To be perfectly successful, a tube other than the ordinary lithotomy tube or soft catheter must be used. These are liable to blockage either from the falling down of the collapsed bladder wall or by mucus and pus. I found a silver lithotomy tube pierced with two rows of large closely set openings for the terminal $1\frac{1}{2}$ inch of its length sufficient for the purpose. The puncture in the perineum being made after the suprapubic wound with the left index finger in the prostatic urethra from above as a guide is one of the simplest of counter-openings, large enough only for the introduction of the tube, and, if kept accurately in the middle line, necessitating the loss of scarcely a drop of blood. True drainage is secured as distinguished from the overflow through the suprapubic wound, which, together with the cavity of the bladder, is kept perfectly clean, and thus absorption from a large raw surface is avoided.

The first time I was led to practise this drainage was purely accidental. The patient was an emaciated Arab, 30 to 40 years old, with a stone (since given to the Leeds Museum) over 10 ozs. in weight, largely phosphatic, with alkaline urine. Having tied the penis with a tape after the injection of only 4 ounces of fluid (instead of using the pressure of a column of fluid through a large catheter, as should always be done), the membranous urethra ruptured, and the penis, perineum, and scrotum rapidly filled up. This is an accident which I have not seen recorded elsewhere. The vesical walls were much hypertrophied. To relieve this, I made a median puncture at the base of the perineal swelling, extending into the urethra, so that it should be available for bladder drainage afterwards. But it necessarily wounded the bulb, and the consequent loss of blood must have contributed to the ultimate fatal result. For some days soft catheters were used for drainage, and the patient put on his back on an inclined plane. But it was not till the above-mentioned tube was used that we succeeded in keeping the large upper wound dry, and even then it was found advisable to insert a strip of gauze into it as an additional precaution. On the fifth evening the drainage was discontinued, as the one constant position was irksome to the patient. But next morning the usual drop in temperature was absent, and the patient distinctly not so well, so it was resumed till the ninth evening, when again an attempt was made to dispense with the tube. The patient's temperature had varied between 100° and 103° F., but his tongue was