abnormally sensitive to the vaccine, yet the clinical change for the better appears in a much shorter time, and we may therefore only conclude that the patient's general resistance does not coincide with his resistance at the focus of infection. This is demonstrated in a remarkable manner in cases of large carbuncles accompanied by grave toxic symptoms. The disappearance of the slate-blue colour round the lesion and its replacement by a bright red erythema is a striking change that occurs in a few hours. A possible explanation is that the first antibodies produced by the vaccine are all removed from the circulation as quickly as they are formed by some process of concentration at the site of infection. Whether the inflamed tissues in that area play an active or a passive part in this fixation of the antibodies is a matter of speculation at present.

The following cases are the first few treated with the vaccine, and nearly every one illustrates some point discussed above. The cases have not been "selected," but are a true consecutive series.

Case 1.—Aged 26. Septic finger due to running dirty scissors into it. Next day, small pustule, lymphangitis, and axillary adenitis. The finger was fomented and the lymphangitis disappeared, but swelling of the whole digit commenced. Serum only on incision. General lassitude. Vaccine given on seventh day. Next day the swelling was rapidly subsiding; there was distinct tenderness at the point of original injury. Pus visible that evening; on incision, a large quantity escaped and there was complete relief from pain. Next day, a clean granulating surface, which was completely healed on the fourth day after administration of vaccine.

Case 2.—Aged 11. Streptococcal peritonitis, apparently idiopathic. Abdomen opened, drain inserted: prognosis hopeless. Vaccine given twenty-four hours later, up to which time the patient was maniacal and apparently dying. Heroin failed to calm the patient. Six hours after vaccine was given the patient went quietly to sleep, temperature and pulse fell, the former several degrees, and the case made an uneventful recovery.

Case 3.—Pyæmia; purulent arthritis of right elbow with abscesses in axilla and groin. Temperature 100° to 102.2° , pulse 100 to 120. Vaccine given. The next day the temperature was still swinging, but on the down grade. A second dose was given four days later, and temperature and pulse settled finally on the tenth day.

Case 4.—Aged 11. Acute osteomyelitis of spine and infection of canal. Prognosis hopeless. Vaccine given on the sixth day at 11 a.m. That evening the temperature rose half a degree, but the next morning was down one degree; the pulse fell from 140 to 110. The patient was operated on that afternoon, and died early next morning.

Case 5.—Carbuncle of hand, involving half of palm, index finger, and back of hand; acute toxic symptoms and swelling of arm; tissues blue-grey in appearance. Prognosis very serious. Vaccine given; no reaction, but tissues had lost blue colour within twelve hours; the carbuncle was surrounded by deep scarlet erythema, hot to the touch; toxic symptoms much less. Not content with this a second dose was given next day, and caused an alarming reaction for a few hours; then all passed off. A large slough separated on the eighth day, and a clean granulating surface was left which healed rapidly.

Case 6.—Infant aged 8 months. Furunculosis and toxæmic absorption; sleeping badly. Vaccine given and child taken home. Next day there was great improvement; the child slept well from then onwards. The mother volunteered the statement that the lesions had changed colour from blue to red. No sign of a general reaction. No further treatment needed.

Case 7.—Acute appendicitis, large burst abscess, and peritonitis in a female aged 40. She appeared to be doing well for three days after operation, then became worse—abdomen distended, anxious facial expression, wound drain diminished, temperature and pulse rising. Vaccine given, with prompt improvement within sixteen hours; there was profuse outpouring of pus. The patient looked much better, and distension disappeared. A second dose was given forty-eight hours later. Prompt reaction caused the patient to feel very unwell, and for a few hours the medical attendant had grave doubts as to her recovery. Next day she seemed better than ever, and eventually made a complete recovery.

Case 8.—Phagedænic primary sore of penis; widespread septic infection of groins; advanced pyorrhoea. The patient was in a filthy personal condition. Streptococcus isolated by blood culture. Vaccine given on two occasions without clinical benefit; no reaction at any time. Ulcerative endocarditis found at necropsy.

Case 9.—Acute osteomyelitis in a patient aged 10. Bone opened up and drained, but temperature remained for seven days after operation at about 103° , with no sign of subsiding. Vaccine given, and temperature fell to 98° six hours later, and was 97° in another eight hours.

Case 10.—Pyæmia. Patient was ill three weeks before vaccine was given. No reaction or benefit after both fractions of the first dose, so a week's interval was allowed to elapse before giving a second dose. Again no reaction or improvement, so another week's interval was ordered. The third dose was then given, with the same result. All three doses were the same size.

Case 11.—Primary sore of penis with severe balanitis, becoming gangrenous in one part. Vaccine caused a small slough to separate,

with total disappearance of the surrounding swelling, and left a small clean sore. No sign of a general reaction.

Case 12.—Subacute cellulitis of leg. Chronic indurated area which was subject to periodical acute exacerbations. Vaccine cleared up the condition promptly. No general reaction observed.

Case 13.—Streptococcal septicaemia. Patient had been ill for fourteen days when residual vaccine was first given. A previous dose with the common toxin-containing vaccine produced a reaction which lasted several days and from which no good result was derived. The first dose of the author's vaccine produced no result of any sort, and a second dose four days later also failed to make any impression. The temperature remained at about 105° all the time. The patient died four days after the second dose was administered.

These cases cannot be called promising from the standpoint of collecting a good "record" of results, but the author has acted in accordance with the belief that a remedy can only be tested thoroughly when it is given an opportunity to fail as well as to succeed. In no instance was there the smallest sign of an undesirable result due to the vaccine.

One fact is very clear: vaccine treatment can cure when other remedies have been tried and have failed, but the best results can only be expected when the vaccine is given as early as possible in the disease. It cannot be expected to resuscitate those with the death rattle already in their throats, although on occasion it seems to have come tolerably near to it.

The most important feature of all these cases is that in no case where the fractional dose system was followed did any alarming general reaction occur. The cases demonstrate the author's main contention: it is possible to administer a vaccine to a patient suffering from acute sepsis in such a manner that the vaccine is under control and the individual susceptibility of the patient prevented from causing a disastrous reaction. The system in brief is:

1. The largest dose that most chronic walking cases can tolerate without a general reaction is ascertained.

2. One-third of this dose is called the standard dose, and is administered to an acute case in two halves.

3. If after the first fraction there is neither a general reaction nor improvement, the second fraction should be given within twenty-four hours.

4. If the first dose requires both fractions, the amount of the dose should be considered to be that of the sum of the two fractions.

5. The second dose should not be given for at least four days after the completion of the first dose, and can be of the same size as the first or increased by not more than 20 per cent. If a result has been obtained with the first dose, it is doubtful if an increase of amount in the second is of much advantage.

6. It is as well to prepare two stock vaccines—one containing the streptococcus, pneumococcus, *Staphylococcus aureus*, and Hoffmann's bacillus; the other having the colon bacillus substituted for the pneumococcus, but otherwise the same as the first vaccine.

The author is much indebted to Mr. Roy R. Kerr, F.R.C.S., for providing some of the cases, and most particularly for his close clinical observation, which was largely instrumental in furnishing the facts upon which the system of dosage advocated has been built.

THE PARASITOLOGY AND CLINICAL ASPECTS OF MALARIA IN ANTIGUA.

BY

W. M. McDONALD, M.R.C.S., L.R.C.P.,
DISTRICT M.O. AND M.O.H., ANTIGUA, WEST INDIES.

THERE has not, hitherto, been any investigation into the parasitology of malaria in Antigua. The following observations, based on a record of 300 positive blood films, may be of value in providing an accurate indication of the nature of the infection with which we have to deal. Of the 300 positive films which were analysed 61 per cent. showed *Plasmodium falciparum* (malignant tertian), 28 per cent. *Plasmodium malariae* (quartan), and 11 per cent. *Plasmodium vivax* (benign tertian). The predominant infection, therefore, is *Plasmodium falciparum*.

The form of parasite most frequently found was the young form, but it is worthy of note that the classical signet-ring form was conspicuous by its rarity. The majority of young parasites seen were in the tenue form. In most cases in

which the ring form was retained, the usual form seen was a linear or circular central nucleus, surrounded by a blue body, which, as it grows, shows a thickening at one side. The largest subtertian rings seen were about one-quarter of the size of the red cell; these can be differentiated from a benign ring, first, by the sharply defined outline and decided thickening on one side (the side farthest from the nucleus), a benign ring of the same size being flimsy and ill defined, and showing a uniform thickening in contrast to a one-sided thickening. Secondly, it can be differentiated by its effect on the red cell which contains it; in a subtertian infection the containing cell is normal in size and colour—that is to say, where there is general anisocytosis or anisochromia, the affected cell may vary in size or colour within the same limits as unaffected cells, whereas, when the parasite is benign, the affected cell is always swollen, paler, and distorted, and often shows Schüffner's dots. The subtertian ring, moreover, frequently has a double nucleus, whereas this is very rarely seen in a benign ring. The difference in colour, size, and shape of the affected cells is, however, so marked that there is no difficulty in distinguishing between a benign and a subtertian ring form.

The greatest difficulty occurs in distinguishing between a young quartan and a young subtertian parasite. The young quartan often has a double nucleus, and the affected cell is generally slightly smaller than normal. I have found that the young quartan shows a more intense colouring than the subtertian, and that it does not develop the characteristic one-sided thickening of the protoplasm that the subtertian does. In a quartan infection it is usual to find two or three stages of growth in the same film, whereas in a subtertian rings only are found, or rings and crescents. I have found on only two occasions the sporulating form of the subtertian parasite in the peripheral blood; the spores were smaller and paler than in the similar stage of quartan or benign parasite; one showed thirty small spores and the other seventeen, while the pigment was black, whereas in quartan it shows yellowish-brown in the sporulating stage. Sporulating bodies are commonly found in quartan infection, the number of spores varying from 5 to 15, but averaging 8 to 10, and the spores are large and stained deep red. In the benign parasite the spores vary from 12 to 25 in number, they are much more crowded together, and the pigment is neither central nor clumped. In fact, in my experience the benign tertian parasite never assumes what is known as the "rosette" form; this form is limited to quartan and subtertian infections. Crescents were found in 32 per cent. of the subtertian cases. This is of interest in view of the fact that many authorities believe that crescents are very rarely seen in the peripheral blood.

In a recent article in the *Lancet* by Majors Pratt-Johnson and Gilchrist, it is stated that "crescents are not found in the untreated or in acute cases of the disease." My own experience does not agree with this statement. The cases in which crescents were found were all untreated cases, or at any rate cases which had not been recently treated, and whereas crescents alone are not found in acute cases, yet in many cases of acute illness crescents are found at the same time as the rings. In only three cases did I see Maurer's dots. It would appear that these dots are not dependent on differences of staining, but on some particular factor which has not yet been determined. Pigmented mononuclear cells were found in about 60 per cent. of cases. The chief characteristic of the differential count is a marked increase of large mononuclears and large lymphocytes, a reduction of small lymphocytes to about 3 per cent. of the total count, and a low polymorphonuclear percentage. Eosinophils are normal or slightly increased. The chief characteristic of the red cells is a marked anisocytosis; this is always present and varies only in degree. Abnormal changes, such as polychromasia and basophilic stippling, are sometimes seen.

Clinical Aspects.

The most striking characteristic of malaria in Antigua is the association of the subtertian parasite with clinically mild symptoms. The generally accepted view is that this parasite shows far greater pathogenicity than either of the other two varieties. Here, however, in Antigua, although *Plasmodium falciparum* is the commonest variety found, yet its malignant manifestations are apparently limited almost entirely to young children. I say "apparently" because I believe that further investigation will prove that the malignant nature of this parasite manifests itself locally, not so much in malignant attacks of fever as in the sequelae which, developing at

varying intervals after a malarial paroxysm, have not hitherto been attributed to their true cause, which is malaria. I may mention in this connexion that I have recently seen three cases of nephritis, one of which was fatal, in which *P. falciparum* was found in the blood.

As a general rule, however, this infection is associated in the adult with mild symptoms. The cases under review are patients who have been well enough to walk some miles to the surgery for treatment. The temperature is low, between 99° and 101°. Cases which exhibit a definite forty-eight hours' periodicity are the exception rather than the rule. The usual type is a low fever exhibiting a mild quotidian paroxysm and sometimes a double quotidian. As a rule the spleen is not enlarged, and subjective symptoms are not marked, the patient complaining of slight pains in the limbs and general malaise, which, however, is not sufficiently marked to prevent him from working. Young children, on the contrary, suffer severely and are often acutely ill as a result of malarial infection. I have seen many cases of convulsions and coma in young children, due to infection with *P. falciparum*. The younger the child the more acute are the symptoms. In the series of cases under review there are only five fatal cases, all being children under ten years, all being associated with convulsions or coma, and all being due to *P. falciparum*. This infection is undoubtedly a very grave factor of infantile mortality.

Plasmodium malariae (quartan) exhibits a slightly different symptomatology. Patients seldom seek medical aid for a simple quartan infection. Practically all the cases which have come under my notice exhibit a quartan triplex infection, in which parasites of several different stages of growth are found at the same time, and the resulting pyrexia is quotidian. The patient, as a rule, complains of nothing beyond the fever, which clears up readily under quinine treatment, but relapses more frequently and more persistently than other infections.

Benign tertian infection shows a sharp paroxysm and a more definite periodicity, but yields more readily to treatment than either of the other two varieties. I have seen only three cases of mixed infections, the parasites found in one case being crescents, malignant rings, benign rings, and benign sporulating bodies; the other two were malignant and benign, and malignant and quartan, respectively.

Treatment and Relapses.

Owing to the fact that the cases under review are surgery patients who cannot be kept under observation or followed up it is impossible to give a scientifically complete treatment to any case, but as far as the treatment of the actual malarial attack is concerned I find that the essentials of successful treatment are (1) initial dosage with calomel and salts, and (2) large doses of quinine. The initial dosage with calomel and salts is of the greatest importance, as this makes the action of the quinine much quicker and surer. In the cases of adults I give 30 to 45 grains of quinine daily. Cases clear up very quickly under this treatment. Patients are recommended to continue quinine treatment, but as a matter of fact they never do. It is not possible, therefore, to give any treatment with a view to avoiding relapses. In this connexion it is of interest to note that relapses do not appear to be more frequent than in places where energetic anti-relapse treatment is undertaken. I have records of relapses in about 8 per cent. of the total number of cases. This record is not scientifically accurate, as the cases are not under observation, and many others may have relapsed without coming up for treatment. The greatest number of recorded relapses are quartan cases, with malignant next; and in this particular series none of the benign cases have relapsed. Of the comparatively small proportion of cases which I have been able to keep under observation I can write with certainty of the absence of relapse in the following:

Malignant (acute) treated by one intramuscular injection and 30 grains quinine daily by mouth	2 cases.
Treated 18 months ago; no relapse	1 case.
Treated 13 months ago; no relapse	1 "
Malignant; treated for 3 days only, 14 months ago; no relapse	1 "
Malignant; treated 10 months ago	1 "
Malignant; treated 3 months ago	1 "
Quartan; treated 9 months ago	1 "
Quartan; treated 5 months ago	1 "
Quartan; treated 14 months ago—one intramuscular and oral treatment	1 "

These ten cases did not receive any quinine treatment after the initial attack had subsided. In the small proportion of

cases treated by intramuscular injection four relapses are recorded. In no case did any local trouble occur as a result of intramuscular injection.

General Remarks.

In concluding this report I would draw attention to the connexion between the malaria incidence and the death rate. Our death rate is more than twice as great as that of England and Wales; and while deaths recorded as directly due to malaria provide only 2½ per cent. of the total, yet due weight should be attached to the probability that malaria, on account of its effect in lowering the resisting power of the body to other infections, is a contributing factor to many more deaths than those of which it is the direct or recorded factor, and that, further, there are strong indications that malaria sequelae are responsible for many deaths which are recorded as due to other causes. In view of these facts it cannot be too strongly represented that vigorous anti-malarial measures should be instituted throughout the island with a view to (a) lowering our excessively high death rate and (b) improving the quality and quantity of our labour supply.

THE APPLICATION OF FASCIA LATA IN PLASTIC SURGERY.

BY

CYRIL H. CUFF, M.B., B.S. DUNELM., F.R.C.S. EDIN.,
RESIDENT SURGEON, VICTORIA HOSPITAL, ST. LUCIA, BRITISH
WEST INDIES.

In a previous article¹ I ventured to enumerate some of the uses to which the fascia lata might be applied in plastic surgery. I now wish to illustrate, by a series of cases, some of the suggestions already put forward. The operations subjoined have all been carried out during the last two years, and the results, on the whole, have been encouraging. Technique must be most careful, and every effort in reconstruction should be based on strictly anatomical lines wherever possible. I have also found it advantageous to have no "slack" after suturing the fascia lata *in situ*, but rather to have the graft under slight positive tension.

CASE I.

A man was admitted to hospital suffering from a faecal fistula in the left iliac fossa the result of an old gunshot wound of abdomen. The fistula communicated with the descending colon, and was about the size of a shilling. There was no spur, and the finger could be passed freely in both directions. Operation took place on January 27th, 1920. The opening was surrounded by a free incision one-sixth of an inch from edge, the bowel mobilized, and margin of the opening infolded by a series of Lembert sutures. A portion of fascia lata from the thigh, 2½ inches by 4 inches, was sutured with medium catgut over the deficiency in the oblique muscles from the rectus sheath medially to the remains of the external oblique aponeurosis laterally. Tension in the skin wound was obviated by a pedicle flap turned down from the left hypochondriac region. The wound healed under one dressing, and when last heard of in December, 1920, the patient was doing well.

CASE II.

A man, who had sustained a gunshot wound of left forearm, was admitted with partial wrist-drop and inability to extend fingers or thumb at terminal phalanges. He had a long scar along the upper half of the posterior aspect of forearm, and some obvious loss of muscle substance. An operation was performed on May 18th, 1920, a long exploratory incision being carried over the scar, extending to the wrist. Upper portion of the extensor muscle was found to be healthy; the lower portion and communis tendons in the upper part were an adherent mass of fibrous tissue. Tendons just above posterior annular ligament were healthy. The intervening scar tissue was freely excised, and the communis and extensor longus pollicis tendons identified below. The extensors of the carpus were freed from the overlying scar tissue. A fascial flap 4 inches by 1½ inches was cut from left thigh and split in its lower half. The upper portion was sutured to the healthy belly of the extensor communis digitorum and longus pollicis, the hand and fingers being in a position of dorsiflexion. The lower divided portion was sutured to the communis tendons and longus pollicis. The hand was put up in a long "cock up" splint, sutures removed at the end of three weeks, and gentle passive movement commenced. At the end of two months a fair degree of movement in the fingers was obtained, and even more in the thumb, but on the whole the benefit was less than anticipated.

CASE III.

A man was admitted with an old gunshot wound of the middle third of left shin. There was a painful adherent scar, about 2 inches long, about the middle of the subcutaneous border of tibia. At the operation on April 22nd, 1920, the scar was freely excised, also the underlying scar tissue. The edges of wound were

undercut and two parallel tension incisions made, 1 inch from sides of wound. A flap of fascia lata 2 inches by 1 inch was sutured to the periosteum, with the fatty side subcutaneous, and the wound closed without tension with three sutures of "bipped silk." The result was very satisfactory, the new scar remaining elevated and painless.

CASE IV.

A man was admitted suffering from an old gunshot wound of the mandible with non-union and loss of substance on the left side. There was a gap of 1 inch midway between the symphysis and angle. The fragments were largely subcutaneous, as there was loss of masseter and facial muscles, and the skin was adherent to the bone. Operation was performed on July 5th, 1920, a curved incision being made, commencing at the lobule, to the symphysis. Fragments were freed from skin and scar tissue, sclerosed tips excised and a gutter cut in them in the usual manner with an Albee saw. A graft was cut from the tibia 2½ inches long and secured in the gutter with kangaroo tendon. A strip of fascia lata 4 inches by 1½ inches was then carefully wrapped around the graft and sutured to periosteum of fragments. The skin was closed with horsehair. At first there was some serous discharge, which lasted for nearly a month, but finally the wound closed and the result was very satisfactory. The fascia lata probably saved the graft in this case.

CASE V.

A man was admitted to hospital suffering from hernia of the right lung, through the third interspace, about the nipple line. There was a tympanitic swelling, about the size of a cocoanut, and a marked impulse on coughing, reducible on pressure. The conditions had followed upon an old injury to the chest by the shaft of a cart. Operation was performed on September 8th, 1920, a large curved incision being made from the anterior axillary fold, across the right chest and down towards the ensiform cartilage. The pectoralis major was split about its centre, and the sac (formed by the parietal pleura) dissected off the thin and stretched pectoral muscle and ribs. The third and fourth ribs were found to be widely separated. During this dissection, the sac was accidentally incised and air rushed in. The opening in the pleura was closed with a purse-string suture of catgut, and the sac, after being separated, was invaginated through a defect in the intercostal space. Following upon a suggestion of Professor Rutherford Morison, advantage was taken of a lipoma, situated over the right deltoid. This was dissected out, split into two portions, and inserted into the defect. A sheet of fascia lata 4 inches by 2½ inches was stretched firmly over this and sutured to the divided and separated periosteum of the rib above and below. The pectoral muscle was then adjusted and the wound closed. A report dated January 12th, 1921, states that the chest in the position of rest is quite flat, and with the hardest cough the patient is only able to produce a small impulse without any bulge.

CASE VI.

A pensioner was admitted with "drop-foot," the result of nerve injury consequent upon a gunshot wound of the thigh. Previous operations for nerve suture and for tendon transplantation had failed. At an operation performed in June 29th, 1920, the tendo Achillis was lengthened subcutaneously, the base of the fifth metatarsal and tubercle of scaphoid exposed through small incisions, and the crest of the tibia and fibula exposed through a 3-inch curved incision at the junction of lower and middle thirds. The aponeurosis of the tibialis anticus and peronei was split in the line of incision, and the periosteum of both bones exposed. Fascia lata of right thigh was exposed through a 12-inch incision, and a portion 12 inches by 2 inches removed. This was split from below upwards for 7 inches, thus resembling a pair of trousers. The upper portion was sutured deeply to the periosteum of the tibia and fibula, and to the split aponeurosis. The lower bifurcated portion was pulled down by tunnelling the subcutaneous tissues with forceps, and a limb drawn into each of the two lower wounds. Here with the foot in full dorsiflexion all slack was taken up, and the two strips sutured to the periosteum of the scaphoid and the fifth metatarsal respectively, and to the plantar fascia. The wounds were closed with silkworm gut and a back splint applied. In January, 1921, he could walk well without any special boot.

CASE VII.

A man was admitted with partial rupture of the short head of the biceps of the left arm, the injury having occurred three months before admission. On examination, there was a small hard nodule at the junction of upper and middle thirds of the left arm, on the medial aspect, and above this, a definite hollow. Movements of flexion and supination were weak and very painful.

An operation was carried out on September 20th, 1921, a long incision being made from the coracoid, along the antero-internal aspect of the upper arm to the junction of middle and lower thirds. The scar in the biceps was located just about the junction of the two heads. The site of the scar was firmly adherent to the underlying brachialis anticus. Adhesions were separated, and the tendon of the short head, which was torn and slack, tightened up. A strip of fascia lata, 4 inches by 2½ inches, was then wrapped round the site of rupture, and secured with catgut sutures. A further small portion was sutured over the scarred area on the brachialis. The wound was closed in two layers, and the arm slung in a "dog collar" in semiflexion. At the time of writing the arm is gaining power daily, and movements are free from pain.

I have selected these cases as fair examples of the practical uses to which the fascia lata may be put. It has, of course, been utilized for many other purposes, and doubtless others will yet be devised.