On September 12th vomiting was less, and she was able to recline slightly; the opium mixture was now given every three hours.

On September 13th sulphate of magnesia 30 grs. and tinct. opii π 20 were given every three hours, and the bowels acted the same day. The circumference at the umbilicus was reduced to 33 inches.

On September 14th the local dullness over the cæcum was becoming more defined, and on September 16th the dullness in the flanks was disappearing and the patient able to lie down. Sulphate of magnesia, iron, and strychnine were ordered three times daily. The bowels were moved twice daily and nourishment was taken well. In the dullness over the cæcum, which was an irregular circle 6 inches in diameter, fluctuation was detected about September 25th, and under the influence of poultices gradually pointed.

On September 27th the patient stated that she passed about an eggcupful of matter *per rectum*.

On October 3rd 18 ounces of a bad-smelling purulent fluid—specific gravity 1026—were drawn off from the area of cæcal dullness with an ordinary aspirator under antiseptic precautions. There was a quantity of gas drawn off at the same time, and this was probably already in the fluid, as the needle was not moved about either before or after the pus began to flow.

On October 9th she was aspirated again and 8 ounces of pus drawn off. After this she made an excellent recovery, has enjoyed perfect health ever since, and is able to look after her household duties, including washing.

REMARKS.

Jenner declared perforation to be always fatal. In this case the perforation was due to some half-dozen unripe plums the patient had surreptitiously obtained from a child, and eaten on the afternoon of September 9th. At 10 o'clock that evening she was seized with violent pain and vomiting, which grew worse during the night. Among other points of interest, the following are worthy of notice:

1. The sudden elevation of temperature when the salol was inadvertently stopped, and the sudden fall when recommenced.

2. The harmlessness of giving sulphate of magnesia with opium three days after perforation and the probable good effects it had on the effusion in the flanks.

3. The abrupt termination of the course of the fever after the perforation, the temperature not reaching 99° after the seventeenth day.

4. The comparative good health and appetite existing even with more than a pint of pus in the abdominal cavity.
5. The good results following aspiration without drainage.

A CRITICAL ANALYSIS OF THIRTY CASES OF DISLOCATION OF THE SHOULDER TREATED BY MANIPULATION.

By JOHN LYNN THOMAS, F.R.C.S. Eng.,
Cardiff.

THE majority of the thirty cases were treated by me whilst house-surgeon at the Cardiff Infirmary under favourable conditions to try various adjuncts when simple means failed. The time it took to reduce the dislocation varied from fifteen seconds to six minutes; when the latter time was exceeded other means had been resorted to. Twenty-four cases were reduced without any trouble at the first attempt on the day of the accident; they were dislocations into the axilla, four in females, and the rest in men of various occupations requiring fairly strong arms—coal trimmers, masons, carpenters, common labourers. The method employed was a modification of that described by Professor Kocher, of Bern, at the International Medical Congress of 1881.

Of the six failures at the first attempt, No. 1 was of eight weeks' standing; various methods under chloroform narcosis failed, and the patient was discharged with a useful movable false joint. Nos. 2 and 3 were recent dislocations in spare non-muscular men; manipulation repeatedly failed, and the heel method was at once successful. In No. 4 manipulation, heel method, and circumduction failed; under chloroform narcosis reduction by manipulation was effected in thirty seconds. In No. 5, after the patient was anaesthetised, manipulation again failed, but the heel method was successful. In No. 6 repeated manipulation failed; traction with the patient's arm round the operator's waist was successful.

Of the twenty-four successes two deserve notice. One was that of a circus performer, and the most muscular man I ever treated. Reduction took place in less than two minutes, although there was much muscular irritability; the other case was of three days' standing, in a man aged 60, with much bruising of the axillary folds from repeated attempts at reduction with the heel in the axilla by competent practitioners. It was reduced in 45 seconds in the first stage of manipulation.

The method employed was the following. With the patient lying on his back the operator grasps the patient's hand in

one hand, lays hold of the arm immediately above the elbow with the other, flexes the elbow to a right angle, and then gently and firmly rotates outwards the arm as far as it will go (*first stage*).

Rotation outwards is maintained, and the elbow is carried to the patient's side (*second stage*).

Then, if necessary, the elbow is carried forward close to the curve of the thorax (*third stage*).

One piece of strapping fixes the elbow to the side, and the hand is carried in a sling, which is more comfortable and wholesome to the patients than the ordinary bandaging.

Several of the dislocations were reduced in the first stage of the manipulation, the majority in the second stage, and the rest in the third. The further movement of the elbow to the sternum and rotation inwards of the arm I found unnecessary and of no avail in the unsuccessful cases.

I have seen Kocher's plan tried in a few cases, but the result was not so satisfactory as with the patients lying on their backs. Manipulation succeeds in muscular as well as in non-muscular patients, and the failures are generally attributed to some unusual irregularity in the rent of the capsule of the shoulder-joint. I think they are also partly due to a defect in the method itself, and that is, want of traction to counteract the direct longitudinal pull of the muscles, which sometimes prevents the head of the humerus descending sufficiently towards the rent in the capsule, by which it is to retrace its steps within the joint again.

It is very difficult to decide which of the many methods used for reducing dislocations of the shoulder gives the best result under all conditions, but I think that manipulation is by far the easiest, most scientific, and neatest, as it requires no special preparation of the patient, operator, nor surroundings, whilst the probability of success¹ at first attempt is high.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

THE POSITION OF THE PATIENT DURING EXCISION OF TUMOURS OF THE RECTUM.

A WEEK ago I removed a tumour of the rectum, about $1\frac{1}{2}$ inch by 1 inch, or rather larger, situated so high up the bowel that it was impossible without enlarging the anus to estimate either its exact size or position. Instead of placing the patient in the lithotomy position I turned him on his face close to the side of the table; the right thigh was extended, but the left hip was flexed, the thigh hanging over the side of the table, and the knee resting on a chair.

The ease with which the operation was performed, as compared with the difficulties which are experienced when the patient is in the lithotomy position, prompts me to ask you to insert this note, because, as far as I know, the latter is the most common, if not the invariable one used, at present. The operation consisted in making a mesial incision as far as the level of the third foramen of the sacrum, removing the coccyx, and separating the attachment of the sacrosclatic ligaments as far as the third foramen on the left side. The rectum was then slit up from behind, and the growth pulled down and removed with scissors. An opening of considerable size was made in the peritoneum, but as this could be perfectly well seen it was stitched up with the greatest ease. All bleeding vessels could be at once picked up without difficulty, and all the steps of the operation could be accomplished with much greater precision than I have ever found possible in the lithotomy position.

Those who have not adopted this plan will, I am sure, find it worthy of a trial, and if it has been put in practice by others I hope that the absence of any account of it in the textbooks will be sufficient excuse for thus drawing attention to it.

Wimpole Street, W.

RICKMAN J. GODLEE.

¹ In Braithwaite's *Retrospect of Medicine*, vol. c, Dr. Powers, of New York, gives 77 per cent.

AÆRATED MILK.

I WISH to add my testimony to that of Surgeon-Major Goldsmith regarding the therapeutic value of aërated milk, as described in the *BRITISH MEDICAL JOURNAL* for July 1st. When Dr. Goldsmith informed me some months ago regarding the details of his method, I at once saw how useful it was likely to be in a class of cases with which I have had occasion to deal, namely, the climatic disease, regarding which I have written as psilosis of the tongue and intestine (Sprue). Into the *rationale* of milk diet in the treatment of that disease I need not now enter; suffice it to say that it is the only treatment deserving of serious consideration. Unfortunately in a certain proportion of cases the patient expresses himself as being quite unable to take milk; and there are cases in which, when he overcomes his dislike, the milk seems for a time to disagree. I have known for years that in some of these cases koumiss can be taken and well borne; but the objection I have to koumiss is that it is evidently less nutritious than pure milk. A patient whose hunger pangs are relieved for an hour and a-half by a certain quantity of milk, will often crave for nourishment in a quarter of an hour after an equal quantity of koumiss. I have found, by experience, in several recent cases that aërated milk combines the advantages of koumiss and of pure milk. I shall probably have an opportunity of publishing details of some of these cases in due time, and in the meantime content myself with simply putting the fact on record.

One patient, whose objection to milk seemed to be insuperable, was trained through koumiss and aërated milk into a full diet of pure milk, and is now quite well.

Another patient, a very chronic case, who had no difficulty in taking milk, but in whom diarrhoea continued unchecked, lost the diarrhoea within three days on aërated milk, which she has continued to take with satisfactory results for six weeks.

A nurse who has taken charge of two of my patients tells me that it is very necessary to have two seltzogenes, in order that one may be thoroughly soaked and cleansed while the other is in use.

Surgeon-Major Goldsmith's method will, I believe, be found to be of great practical value in the varied class of cases in which the nearest approach to rest for the intestinal canal is given by keeping the patient alive on an exclusive milk diet.

Queen Anne Street, W.

G. THIN, M.D.

LIGATURE OF THE BROAD LIGAMENT.

PERHAPS the two following cases may be of interest to those of your readers who have followed the correspondence on "Ligature of the Broad Ligament."

CASE I.—Mrs. C., aged 33, consulted me in January, 1891. She was suffering from excessive menorrhagia, the periods occurring every three weeks or oftener, and being attended with severe pain. She had lost weight and was too weak to do her own housework. Her father had died of cancer of the neck, and one of her brothers of cancer of the tibia. By vaginal examination the uterus was found to be somewhat enlarged and immovably fixed. On February 19th, 1891, I opened the abdomen, expecting to find double salpingitis with extensive adhesions, but instead I found a rounded tumour (the uterus) densely adherent at its base and growing to the rectum and Douglas's pouch. Its removal was impossible, and looking upon it as a case of cancer, with a view of arresting the hæmorrhage and retarding growth I tied the right broad ligament; the left I could not find. Up to the present the operation has fully answered its purpose. On looking up my patient this morning, she told me that since the operation she had been stronger and able to do her work. Her periods come on regularly every four weeks and last six days, the first four days being rather free. She had quite lost the severe pains she used to have. The tumour had extended to within an inch of the umbilicus, and filled the left side of the pelvis.

CASE II.—Mrs. H., aged 47, was sent to me by Dr. Dyson, of Honley, with a view to hysterectomy for epithelioma of the cervix. Her history was that since Christmas, up to which time she had been quite well, she had had a continuous blood-stained watery discharge. Examination revealed a