large increase of the recorded insane in the pauper class, the private or well-to-do class had undergone diminution both in the admissions and the numbers under care. In 1879 the ratio of private lunatics per 10,000 of population was 2.97, which has fallen by steady annual decreases to 2.61; and the admissions of this class have diminished (Dr. Cobbold) in the same progressive manner, from 1.07 per 10,000 in 1882 to 0.92 in 1892. These facts would be a strong argument against the enormous increase theory, but the facts are so complicated and uncertain that it would be unfair to urge this as a proof of an actual diminution in the occurrence of mental defect in the richer classes, although it certainly negatives the theory of large increase.

If the general or any special causes of insanity, the result of the intensified conditions of civilisation, were in operation, they should produce a steady and progressive development of the numbers of the insane. This, however, is not the case, but the contrary is the fact. The rate of increase, as Dr. Merson has pointed out, has fallen from 512 per million in 1859-69 to 361 in 1869-79, and 212 in 1879-89.

The increase of certificated insanity, on the contrary, has not been steady and progressive, but in waves; a large wave (as previously described) corresponding to the capitation grant in 1874, subsiding in 1885, eleven years later, to the level of 1872. The wave may be more easily traced in the smaller areas of counties, the manufacturing and urban preceding the agricultural, but is also very distinct in the statistics of Ireland and Scotland, where, as Dr. Clouston has pointed out, its influence was quite as marked as in England, but occurred a year or two later.

From these facts the conclusions may be drawn that there is no definite evidence of the increased incidence of any special causes of mental disorder, or of the development of new or special forms of disease; that the accumulation of the insane in asylums has been mainly coincident with the operation of the parliamentary capitation grant, and that the rate of accumulation is undergoing a progressively rapid decline.

Further, it seems probable that the increased numbers of the insane in asylums and elsewhere may be accounted for by the greater survivorship in asylums (overriding even a diminished production in the earlier periods of life), by the transfer of paupers and criminals to the lunatic class, and by the increased accuracy of the census returns.

The ultimate conclusion is that the evidence at present available admits of the interpretation that there is no real excess in the amount of occurring insanity, but is even compatible with the possibility of an absolute diminution, which has certainly occurred in the earlier decades of life, and is probable in the well-to-do classes.

THE EMPLOYMENT OF ELECTRIC LIGHTING CURRENTS FOR MEDICAL PURPOSES.

By H. LEWIS JONES, M.D.,
Medical Officer in charge of the Electrical Department,
St. Bartholomew's Hospital.

WITH the steady advance of electric lighting both in London and country towns the application of electric lighting currents to medical and surgical purposes is coming more and more into notice. Already many medical men use the current from the mains in the treatment of their patients; and more would do so were it not for a widespread belief that there is some risk of accidents occurring, though with properly chosen apparatus and with ordinary care there is no danger.

The current supplied by the companies is in some cases an alternating current and in others a continuous one.¹ The pressure at which it enters houses is usually 100 volts, and, as this is too high for medical purposes, some device must be employed before the current can be applied to a patient. When the supply is an alternating one, it is very easy by means of a transformer to reduce the voltage to any required degree, and then there is no difficulty. A Woakes's or similar transformer enables the operator to obtain the most delicate graduation of the voltage, and it is perfectly safe

and easy to handle, and when properly wound it will heat cauteries, light small surgical lamps, or yield an "induction" current for the testing or treatment of paralysed muscles.

The customary pressure for cauteries is 4 volts, that proper for the electric bath is 6 to 8, and for applying "induction currents" to patients is 10 to 16. With the alternating supply the use of a properly graduated transformer² to lower the voltage is in every way better than to use the full pressure of the mains and to choke down the current by means of resistances. On the continuous current systems it is a complicated and expensive matter to reduce the voltage by transforming contrivances, and resistances are therefore more often employed for the purpose of reducing the magnitude of the current. These resistances, however, are not very satisfactory.

Five milliamperes from a source of electricity at 100 volts is more unpleasant than the same current from a battery of low voltage, especially at the moment of first contact with the skin; for whatever the magnitude of the interposed resistance may be, the electrodes before contact with the body are at the full potential of 100 volts, and when the contact is made the high pressure expends itself in quickly breaking down the skin resistance, with more pain and shock than when the resistance is gradually overcome by a lower voltage. The drier the skin the greater is the discomfort. When the skin is thoroughly moistened and reddened this effect is much less marked; though practically I have found that infants and young children are far less tolerant of current from a high than from a low voltage source, even if the current be the same in both cases.

Another objection to the use of resistances to reduce the current is that several separate types and sizes are required for the different applications of the current; a rheostat suitable for a cautery will not do for surgical lamps, nor will one suitable for the latter do for galvanic applications to patients. In the use of electric light mains for heating cauteries, regulation by a rheostat is dangerous, for any accidental fusing of the platinum cautery wire is likely to be followed by the formation of an electric arc at the broken point, an accident which might lead to the most serious consequences. Moreover, the contacts in the handles of the cautery instruments as at present made are apt to give trouble by sparking and burning away.

There is a plan of avoiding these risks by distributing the current along two parallel circuits, one of which carries the cautery, the resistance of the two circuits being so adjusted that the current through the cautery cannot exceed a certain value. In this way the danger of the formation of an arc is avoided, but the apparatus wastes a large amount of current.

By far the best way of utilising the continuous current supply is to charge accumulators with it in the first place, and then to take out current from the accumulators. This plan requires a smaller outlay in plant and apparatus, and the accumulators being portable the operator is not tied down to one spot as he is by a fixed installation, but can carry his charged accumulators to his patient when his patient cannot come to him.

Although accumulators are most unsatisfactory when they are not properly attended to, this is the fault, not of the accumulator itself, but of its treatment. Under proper conditions of recharging a secondary cell behaves very well, though it soon spoils if left uncharged for three or six months at a stretch; such treatment as that will ruin any accumulator.

By means of accumulators the continuous 100 volt electric lighting current can be made available for medical purposes, at a smaller outlay, and with fewer complications than by any arrangement of resistances. The Lithanode Company³ make excellent secondary cells which weigh less than 8 ozs. a-piece, and indeed they have cells weighing as little as 2½ ozs. each, which are perfectly suited for galvanisation of patients, for electrolysis, or for the testing of muscle and nerve and so on.

¹ The St. James and Pall Mall, Westminster, Kensington and Knightsbridge, Chelsea, Charing Cross and Strand, and St. Pancras Companies supply a continuous current. The Metropolitan, the London, City of London, and House to House Companies supply an alternating current.

² A very good transformer is made by Messrs. Miller and Woods, of 2, Gray's Inn Road, and is figured in the *Electrician* for September 29th, 1893.

³ The Lithanode and General Electric Company, Limited, 64, Millbank Street, Westminster, S.W.

Everyone who has a continuous current electric lighting supply available can keep these cells in first rate order with no more complicated apparatus than a switch, an incandescent lamp and a pair of connecting cords, and the larger accumulator cells needed for the cautery or for surgical lamps can be kept charged as easily. All that is required is to connect the cells with the electric light supply for an hour now and then with a 16- or 32-candle power lamp in the circuit to regulate the rate of charging. For a complete outfit a general practitioner should have three batteries, one of 15 or 20 small cells arranged in a battery box with current collector and galvanometer; one 2-cell accumulator for his cautery capable of discharging at a rate of 20 ampères, and one of 4 or 5 cells, capable of discharging 1 ampère, for his electric lamps. For this last purpose nothing is better than the small portable sets which can be bought ready-made for use as travelling or bedroom reading lamps; they weigh only 4 lbs. or 5 lbs., and are easily modified to suit the conducting wires belonging to all surgical incandescent lamp instruments.

The dangers to be feared in the use of electric light currents for patients are those arising from carelessness on the part of the operator, such as connecting up to the wrong terminals, forgetting to put in proper resistances, and the like. The fear of sudden and unexpected rise of pressure in the mains need not be considered, as it is almost, if not quite, impossible for such a thing to happen under the existing Board of Trade rules. With an 8-candle lamp permanently fixed in the circuit with the patient, his body cannot become the path of a current of really dangerous magnitude; indeed it is doubtful whether a pressure of 100 volts could do any serious injury to a human being except under special circumstances of application, such as are outside of the ordinary methods of electrical treatment.

A SPORADIC CASE OF CHOLERA.

By L. W. DARRA MAIR, M.D.LOND., D.P.H.

THE following is an account of the cholera case in Mitcham.

H. B., male, aged 53, a resident in Mitcham, and employed at a local corn factor's, was first taken ill at 10.30 P.M. on September 11th, with rather severe diarrhœa. This continued throughout the night, and also during the next day (September 12th), although its severity was not sufficient to prevent him partially attending to his work. On returning home in the evening, he complained of the diarrhœa for the first time to his friends, and "feeling queer" went immediately to bed (about 7 P.M.). Although he had taken "diarrhœa mixture" purchased from a chemist, the diarrhœa now became very much intensified, and was accompanied by severe cramps. A medical man was sent for at 10.30 P.M.

The medical attendant reports that the cramps were very severe, affecting chiefly his arms and legs, and causing him to call out frequently, and to ask to be rubbed. His expression was anxious, but there was no cyanosis. The pulse was 120, feeble. The temperature in the axilla was 97°. The tongue was moist, and coated with whitish fur. He complained much of thirst. About a quart of watery evacuation was seen, but its colour was obscured by the presence of red stained paper. No pain was experienced on pressing the abdomen. There had been some vomiting earlier. The general condition was one of moderate collapse. The voice was not hoarse. The medical attendant was called to him at 5 A.M., and he also saw him at 8.30 A.M. the next day, September 13th, and soon after in consultation with another medical man. The case was thereupon reported to me, and I saw him at 1.45 P.M., when I found him in a state of collapse, very drowsy, and much cyanosed about the legs and feet and hands, the skin of the latter being puckered up in a singular manner, presenting the appearance of prolonged soaking in water, as seen in washerwomen's fingers. The limbs were quite warm. The face was very pinched, the expression indicated great pain, but there was no marked cyanosis, the lips being perhaps a little dusky. The pulse was 120, but stronger than the general appearance of the patient pointed to. The tongue was coated with thick brown fua, but was moist. The temperature was 97°. Very little,

if any, urine had been passed since 10.30 the previous night, but the diarrhœa had abated considerably, and there had been no vomiting since 5 A.M. No motion had been saved; but some liquid was squeezed out of some sheets which had been placed beneath him. This liquid was like dirty milk and water, and was carefully sealed up in a bottle. He had a severe attack of cramp while I was there, which made him call out with a vigorous voice to be rubbed. Instructions were given that nothing was to be taken out of the room except by the inspector, who was told to call periodically and burn any stained articles that had been used.

He was last seen by the medical attendant at 9 P.M., and he reports as follows: "Temperature 102.4°; pulse 120, exceedingly feeble; respirations 28. Tongue moister and cleaner. Motion passed at 8 P.M., 1½ ounce, of creamy consistency, much bile-stained, chocolate colour, and contained many epithelial scales and blood corpuscles. Very little urine voided. Is drowsy, but rousing frequently and then restless, throwing his limbs about, and turning irritably in bed. Hands cold and rather blue; feet and legs warm, and much less blue than in the afternoon. Voice quite hoarse. Cramps less frequent and less severe. No vomiting. Still great thirst. Face pinched, but not blue."

Death took place at 2.30 A.M. on September 14th.

A necropsy was made at 2 P.M. on September 14th, by Dr. Shelswell, the medical attendant, in the presence of Dr. Seaton, the county medical officer, and myself. Rigor mortis was well marked; the abdomen only was examined; the intestines were distended, pressed against the abdominal wall, and showed much injection of the capillaries. There was no sign of peritonitis. The right kidney was examined; the cortical portion was rather wasted; the medullary portion was apparently healthy; the capsule peeled easily except at one spot. The gall bladder contained about half an ounce of bile. The urinary bladder was quite empty. About one foot of the lower ileum was tied *in situ*, removed and sealed up in a bottle, and the stomach, spleen, left kidney, and suprarenal capsule were also removed and sealed up.

The evacuation, which had been obtained by squeezing the sheets, had been previously taken to the Local Government Board, and the specimens above mentioned, together with the motion passed at 8 P.M. on September 13th, were taken there also immediately on the conclusion of the necropsy.

It was reported that the evacuation first mentioned gave the usual indications of the presence of the comma bacillus, but that in the latter, owing to the fact that Dr. Klein did not receive them until the next morning, when the material was decomposed, the culture experiments were negative, although there could be seen in cover glass specimens bacilli similar to the comma.

Dr. Seaton and I made very careful inquiries, and only elicited the following facts. The patient had not been away from Mitcham for many weeks, except to Tooting for a walk on the Sunday before his illness (September 10th), with other members of his family. He and his family dined off roast pork on that day, all of them (five) partaking of it, and this was also eaten cold by all on the next day. None of the other members of the family were taken ill in any way. He admitted having drunk water from a pump in the yard in which he worked, though he had been told it was only for cattle. The following is an analysis by Dr. Stevenson:

The water was free from odour, of a yellow colour, and slightly turbid. It contained, and yielded, in grains per gallon:

Total solid matters	107.52	Ammonia	0.076
Loss on ignition	7.28	Albuminoid ammonia	0.013
Combined chlorine	9.35	Oxygen consumed	0.121
Nitrates	3.63	Hardness	65°
Nitrites traces.			

The turbidity of the water was due to a slight admixture of chalky earth, and a few living and moving organisms. No specific disease-producing organisms were detected. This is a very impure water, evidently a surface water that has percolated through an impure soil. It is also a very hard saline water. I am of opinion that the water is unfit for domestic use, and that its use for drinking purposes is likely to prove injurious to health.

The only other clue that was found was that the patient had been engaged on September 9th with several other men in receiving a consignment of Russian oats that had just arrived from St. Petersburg. On inquiry, however, it was found that the oats had arrived in bulk, and were unloaded