LECTURES ON INJURIES TO THE EPIPHYSES AND THEIR RESULTS.

Delivered at the Royal College of Surgeons.

By JONATHAN HUTCHINSON, JUNR., F.R.C.S.,

Professor at the College,
Assistant Surgeon and Demonstrator of Practical Surgery at the
London Hospital.

LECTURE II.

THE LOWER EPIPHYSES OF THE HUMERUS.

BEFORE considering the injuries to the lower epiphyses of the humerus a few words must be devoted to the complex subject of their anatomy. At birth the epiphysal disc is a fairly straight line, crossing the bone just above the two epicondyles, the olecranon fossa being immediately over it. Ossification in the capitellum is said to begin during the third year; I have found it at eighteen months, and Farabeuf puts it as usually occurring during the second year. The epiphysis during the first few years increases steadily in width,¹ but comparatively little in depth, so that the olecranon fossa is seen to become more distant from the epiphysal disc, which is more and more curved both from side to side and from before backwards. In the fifth year a nucleus appears for the internal epicondyle, and the latter becomes more and more distanced from the bulk of the lower epiphysis. A third nucleus for the trochlea appears in the eleventh or twelfth year, fusing soon with that of the capitellum, and with the shaft in the sixteenth or seventeenth year, the internal epicondyle (we may here disregard a small separate nucleus for the external epicondyle) uniting about a year later. From these dry anatomical details we can deduce several points of interest:—

1. The lower epiphysis is most likely to be detached *en masse*—that is, taking with it both epicondyles—before the age of 13 or 14 years (Farabeuf wrongly says that it cannot occur after 4 or 5, and that to assert the opposite is a surgical enormity).

2. The epiphysis increasing so slowly in depth comes to lie more and more in the elbow-joint, and as a rule its detachment means a wound of, and more or less hæmorrhage into, this joint.

Fig. 1.—Vertical section through elbow-joint at age of 18 months, showing the epiphyses of the humerus and ulna, and their relation to the joint cavity.

3. The marked downward curvature of the epiphysal line, the distancing from it of the olecranon and coronoid fossæ, and increasing comparative width of the epiphysis, make it

¹ During the first two or three years the epiphysis is usually from 7 to 8 mm. in depth, and at 16 it is only about 10 mm.

extremely probable that a detachment of the latter will only partially follow the epiphysal disc, and that it may be rather a fracture than a true separation. We possess a certain amount of pathological evidence on this much disputed point.

In one case recorded by Clarke² (the patient aged 12) death occurred from gangrene due to a bonesetter's tight bandage. The detached epiphysis was found not to include the epicondyles.

In a second case,³ aged 10, where the same untoward result led to a necropsy, the detachment was above the epicondyles, and followed the epiphysal line closely. In a third, aged 2, under my own observation, death occurred from croup at the end of fifteen days after the accident. The specimen is in the Royal College of Surgeons Museum, and shows that the line of separation follows only half-way the epiphysal disc, diverging for the other half into the diaphysis (see Fig. 4).

Fig. 4.—Detachment of lower epiphysis of humerus, specimen obtained a fortnight after the accident. The line of separation has passed on the inner side into the diaphysis. Bony union was occurring with some lateral displacement.

In a fourth case, aged 13, primary amputation was performed. The detachment followed the epiphysal line exactly except at the extreme external part.

My father and one or two other surgeons have recorded cases of compound separation in which it was found on operation that the epiphysal line had been followed fairly closely. Like Schüller, I have performed a great many experiments on the cadaver, and found that before five or six years it is easy to detach the lower epiphysis of the humerus *en bloc* by various forced movements, but that after that age the separation is usually in part a fracture. As regards the question of shortening in actual practice the point is of no great importance, for the lower epiphysal line of the humerus may practically be disregarded after the tenth year, being only responsible for one centimetre of the subsequent increase in length.

It may be stated that probably a clean separation exactly at the epiphysal disc is uncommon after the age of 5 or 6, but that it may occur some years later, and that in the cases, over fifty in number, now summarised,⁴ it was present probably only in the minority. The youngest patient was 18 months old, the oldest 14 years. Suppuration never occurred unless the injury had been compound from the first, or had become so owing to sloughing of the over-stretched skin. In the great majority there is, when the patient comes under care, more or less deformity due to backward tilting or complete displacement of the epiphysis. This we must ascribe to the acting force being usually transmitted through the radius and ulna from in front, as in a fall on the hand, and partly to the action of the triceps muscle. Not infrequently the surgeon has also to correct a lateral displacement of the

² *Dublin Path. Trans.*, 1868, p. 111.

³ Reeve, of Ohio, quoted by Hamilton in his *Fractures and Dislocations*.

⁴ Twenty-four simple cases, 15 of them under my own observation; 23 complicated, mostly compound, cases; 5 cases of old separation of the epiphysis.

epiphysis either outwards or inwards, due to direct violence on one or other side of the joint, and very occasionally the epiphysis is driven forwards instead of backwards.⁵

The deformity resembles closely enough a backward dislocation at the elbow-joint, and hence frequent mistakes in diagnosis. The antero-posterior measurement of the elbow is increased, the olecranon projects, the triceps is abnormally curved. But the end of the diaphysis has not the characteristic outline of the trochlea and capitellum, the elbow can be hyperextended beyond the straight line, and, as a rule, the epicondyle preserves its normal relation to the olecranon. There is slight shortening of the arm; the deformity can be readily reduced with slight crepitus, and as readily recurs. It has been asserted that nearly all the supposed cases of dislocation of the elbow in early life (that is, under 16) are really examples of detachment of the epiphysis. This, however, is a mistake, for the commonest age for the dislocation is from about 5 to 15 years. It will be presently noted that many, although not all, of the cases of backward or outward dislocation of the elbow in young subjects are complicated by detachment of the internal epicondyle.

The forearm is usually in a position midway between full pronation and supination, and comparatively helpless. If there is detachment without any displacement the best sign is an antero-posterior movement or "give" in the bone just above the elbow, obtained by moderate pressure, lateral movement of the epiphysis being more difficult to elicit.

Fig. 5.—From a case of detachment of lower epiphysis of humerus, showing the backward displacement so often met with.

Amongst the complications we have to notice rapid effusion into the joint and swelling around it, the formation of callus in front of the joint which (if, as often happens, the reduction is imperfect) may be extravagant in amount, extending up the bone for some inches, and even sometimes pressing on the brachial artery; and injury to one or other of the three great nerves in the neighbourhood. Hamilton records one case in which the separation occurred at the age of 4; persistent deformity resulted, and numbness in the area supplied by the ulnar nerve lasted into adult life. In another case he operated at the age of 9 on account of forward projection of the diaphysis, complete loss of pronation and supination, persistent flexion of the wrist and fingers, and paralysis of both median and ulnar nerves. The latter had been displaced in front of the elbow, and was replaced at the operation, which does not seem to me to have been a very thorough one; some bone was merely cut away. No improvement followed.

Volkman treated a case of compound separation of the epiphysis, reducing the protruding diaphysis. The result was fair as regards movements of the elbow, but permanent paralysis of the musculo-spiral remained. In one of my own cases pulsation at the wrist was stopped by the forward projecting diaphysis, and there were marked signs of pressure on the median nerve. These symptoms disappeared on reduction, and recovery was perfect but that flexion was limited to a right angle. In a case recorded by Wright pulsation at the wrist ceased on the eleventh day owing to pressure between the exuberant callus and an anterior splint. The case recovered, but with persistent backward displacement of the epiphysis.

To form a correct idea as to the prognosis of simple detachment of the epiphysis diagnosed at the time of the accident, and treated with splints in the flexed position, I venture to quote my own cases—14 in number. In 6 the result as regards absence of deformity and the elbow movements was practically perfect; in the remaining 8 a certain amount of limitation of extension and flexion—usually about 20 degrees in either direction—with thickening of the lower third of the humerus, remained when the patient was last seen. In only

one was the power of rotation in the least affected, and it need hardly be said that a range of flexion amounting to 80 or 90 degrees is, whilst not ideal, of extreme value to the patient. Such results in unselected cases do not incline one to the view so strongly advocated by Bardenheuer, Illingworth, and others that the treatment of such injuries in the flexed position is wholly wrong and unsurgical. Let us candidly examine the arguments on both sides.

First, the advocates of treatment in full extension might say that the favourable results just mentioned are not the rule. We can answer this by adducing several cases in which great stiffness has followed the use of straight splints, a stiffness far more prejudicial than when it occurs with the elbow flexed. With the fear of this before them, some have advocated passive motion from a very early date. Union will be firm, provided the displacement has been fairly reduced, at the end of three weeks, and it is probably unwise to attempt it much before. To do as Warren⁶ recommends—to employ no splints and to perform so-called "passive" motion daily from the time of the accident is simply barbarous, and a plan which we may leave to surgeons on the other side of the Atlantic.

Secondly, it is said the backward displacement of the lower fragment which is so common is best corrected in the extended position, and hence there will be less callus thrown out. Perhaps this is true; my own experiments on the dead subject after detachment of the epiphysis went in the opposite direction, there being a greater tendency to tilting when the elbow was extended than flexed.

Thirdly, it is undoubtedly easier to put up the elbow in the proper relation of the arm and forearm axes when extended. In case after case which has been carefully treated in flexion the "extension angle"—about 170 degrees open outwards—has been converted into an angle open inwards (see Fig. 6). It may at once be freely conceded that it is extremely difficult to prevent some lateral tilting of the epiphysis if an internal angular splint is used. *Per contra*, it may be said that full extension is an irksome position, especially if effusion comes on in the elbow-joint, and that to employ it without much discomfort almost necessitates keeping the patient in bed.

Fig. 6.—The left arm of a case of separation of the lower epiphysis of the humerus, seen from in front, when fully extended. Good union was obtained, but the alteration in the axes of the arm and forearm due to lateral displacement of the epiphysis is made evident by the dotted line (the normal position).

The whole question is still unsolved, and I shall be excused for dwelling on it here by the extreme importance of obtaining the best possible result in these injuries to the elbows of children. Personally, I believe in the virtues of a well-moulded splint applied to the back of the arm and forearm in the flexed position, and whilst extension is made on the lower fragment through the forearm. The back splint with an interruption opposite the elbow is also very useful.

As regards compound cases, where the diaphysis has protruded through the skin, we have records of 15 cases only, too small a number to draw any strong conclusions from. In six of these the diaphysis was reduced more or less perfectly, one recovered with an almost perfect arm, one with thrombosis of the brachial, a third with paralysis of the musculo-spiral. The other three ended in amputation, one primary, two secondary, owing to gangrene from the artery being torn or obliterated by splint pressure. In three cases a more or less complete excision of the joint was performed; one recovered with very good movements, one with a stiff elbow, the third

⁵ In one case under my care this forward displacement was well marked.

⁶ Warren, *Transactions of the American Medical Association*, vol. 1, p. 174

with a fairly movable one. There seems to be no object in removing the epiphysis unless it is broken into fragments. In six cases the projecting diaphysal end was cut off in order to procure reduction; four of these ended in a most satisfactory result, one recovered with an ankylosed elbow, the sixth underwent amputation at the end of three weeks because no union had occurred—surely a premature proceeding. In two of the fifteen cases an action for malpraxis was brought. One feels certain that these results do not represent what may fairly be expected of the resources of surgery at the present time. Finally, we have to note two cases of operation for old deformity due to backward and inward displacement of the epiphysis. Both were under the care of Professor Bardenheuer, who divided the bone and removed part of the diaphysis. In both great improvement was obtained.

Included in the lower epiphysis of the humerus during the first few years are the two projections, the epicondyles, which later on ossify by separate nuclei, and one of which (the internal) deserves special consideration. The external epicondyle is so small and so extremely rarely injured that we may disregard it. The internal epicondyle by the fourth or fifth year has attained sufficient size to make its separate detachment probable, and between this date and the eighth year ossification takes place in it. Fusion with the rest of the bone occurs in the seventeenth to the nineteenth year. The age of the patients whose records I have collected (38 cases of detachment of the epicondyle) ranged between 8 and 18 years. Small as the epiphysis is (never more than 16 or 17 mm. in vertical measurement and half that in width) there are certain points about its separation and the complications attending the latter that make it worthy of more notice than its size would seem to warrant. In the first place, when the epicondyle is separated the elbow-joint is probably always opened; secondly, union takes place by bone only in rare cases; thirdly, the attachment of the strong internal lateral ligament and of several muscles to the epicondyle accounts for the frequency with which it is dragged off when lateral strain is brought to bear upon the elbow, and also for the ease with which true dislocation follows, once this small epiphysis has been detached. Finally, and this is the most important point of all, the epicondyle is in one sense the guardian of the ulnar nerve at the elbow, and when displaced is liable to exert most injurious pressure upon this nerve. In a paper published in the *BRITISH MEDICAL JOURNAL* last year I brought forward several instances of this complication, and expressed the belief that the difficulty in obtaining good movement after this accident was largely due to every attempt at flexion of the joint increasing the pressure upon the nerve.

A boy came under my care at the London Hospital who had simply detached the epicondyle, had been treated most ably at one of the other hospitals (where passive motion had been very frequently used, and he had been three times anaesthetised for this purpose). It was now three months since the accident; he could not bend the elbow 10 degrees from the straight line; had complete anaesthesia in the ring and little fingers, and wasting of the hand muscles supplied by the ulnar nerve. I excised the epicondyle (which was found to be the entire epiphysis), and at the operation found it to be displaced downwards, stretching the ulnar nerve over it, and united only by fibrous tissue to the humerus. After this the power of flexion was almost perfectly regained, and his anaesthesia disappeared; the muscles regained their power more slowly.

A second, practically identical case, I have since operated on with an equally satisfactory result.

We have only one displacement to consider, that in the downward direction, sometimes also a little forwards or backwards, and by any ordinary method of treatment there is practically only one result to expect—fibrous union. To my mind the only chance of obtaining bony union in good position after displacement of this epiphysis would be to fix it with an aseptic steel needle, but I am not aware that this has ever been tried. Whether the elbow is put up flexed or extended seems to make little or no difference as regards the displacement; Pézerat and Bardenheuer recommend the latter position, and all are agreed that gentle passive motion should be begun from one to three weeks after the accident. The amount of trouble to the surgeon and the patient that an apparently simple case of detachment of this insignificant epiphysis may cause is quite surprising. At the same time it may at once be admitted that in many cases a very good functional result is obtained. Thus, in 10 cases under my observation almost perfect range of movement in the elbow-joint was ultimately secured in the 7 which were followed up long enough to decide. Perseverance in the performance of passive motion

was required in nearly all for some weeks. We have already noticed the coincidence of dislocation of the elbow, either backwards or outwards, with detachment of the internal epicondyle. The explanation is of course the same that accounts for Pott's fracture at the elbow being so often a mixed fracture and dislocation. A violent strain upon the internal lateral ligament of the ankle finds the weak spot at the epiphysal disc between the epicondyle and the rest of the bone; once the former is detached, there is practically nothing to hinder the remainder of the force producing a true dislocation. Hence a small practical deduction: when the surgeon has reduced a dislocation of the forearm in a young subject, let him examine the internal epicondyle. The chances are at least even that he will find it has been detached, and this will make it the more important to prevent the dislocation recurring by the proper use of a splint, and to obtain union of the little fragments of bone (fibrous though it will probably be) in such a position as to interfere least with the ulnar nerve and the elbow-joint.

SYMMETRICAL TALIPES DORSALIS IN AN ACEPHALOUS FŒTUS.

By JOSEPH GRIFFITHS, M.A.CANTAB., M.D.EDIN.,
F.R.C.S.ENG.,

Assistant to the Professor of Surgery in the University of Cambridge.

DURING the past summer Dr. Grove, of St. Ives, kindly sent me an acephalous fœtus, in which there was also a condition of spina bifida extending from the skull to the mid-lumbar region of the column. The spine was strongly bent forwards in the dorsal region, as is not infrequently the case in spina bifida. The spinal cord could not be discerned in any part of its length, but the spinal nerves could be displayed by careful dissection. The fœtus was in other respects normal and well formed, but the feet were symmetrically deformed in an unusual manner, as may be gathered from the accompanying sketches taken at the time. (See Fig. 1.)

Fig. 1.—Talipes dorsalis, left foot.

This deformity at first sight resembled that known as talipes calcaneus, but an inspection of the sketches will show that the heel is not drawn down, but slightly drawn up, and the whole bone (os calcis), with that part of the foot, is hori-