Upregulation of Cytokines Is Detected in the Placentas of Cattle Infected with *Neospora caninum* and Is More Marked Early in Gestation When Fetal Death Is Observed[†]

Anne Rosbottom,¹ E. Helen Gibney,^{1,2} Catherine S. Guy,¹ Anja Kipar,² Robert F. Smith,³
Pete Kaiser,⁴ Alexander J. Trees,¹ and Diana J. L. Williams^{1*}

Veterinary Parasitology, Liverpool School of Tropical Medicine/Faculty of Veterinary Science, University of Liverpool, Pembroke Place, Liverpool L3 5QA, United Kingdom¹; Department of Veterinary Pathology, Faculty of Veterinary Science, University of Liverpool, P.O. Box 147, Liverpool L69 7ZJ, United Kingdom²; Department of Veterinary Clinical Science, Veterinary Teaching Hospital, Leahurst, Neston CH64 7TE, United Kingdom³; and Institute for Animal Health, Compton, Berkshire RG20 7NN, United Kingdom⁴

Received 8 November 2006/Returned for modification 30 December 2006/Accepted 12 February 2008

The protozoan parasite *Neospora caninum* causes fetal death after experimental infection of pregnant cattle in early gestation, but the fetus survives a similar infection in late gestation. An increase in Th1-type cytokines in the placenta in response to the presence of the parasite has been implicated as a contributory factor to fetal death due to immune-mediated pathological alterations. We measured, using real-time reverse transcription-PCR and enzyme-linked immunosorbent assay, the levels of cytokines in the placentas of cattle experimentally infected with *N. caninum* in early and late gestation. After infection in early gestation, fetal death occurred, and the levels of mRNA of both Th1 and Th2 cytokines, including interleukin-2 (IL-2), gamma interferon (IFN- γ), IL-12p40, tumor necrosis factor alpha (TNF- α), IL-18, IL-10, and IL-4, were significantly ($P < 0.01$) increased by up to 1,000-fold. There was extensive placental necrosis and a corresponding infiltration of CD4⁺ T cells and macrophages. IFN- γ protein expression was also highly increased, and a modest increase in transforming growth factor β was detected. A much smaller increase in the same cytokines and IFN- γ protein expression, with minimal placental necrosis and inflammatory infiltration, occurred after *N. caninum* infection in late gestation when the fetuses survived. Comparison of cytokine mRNA levels in separated maternal and fetal placental tissue that showed maternal tissue was the major source of all cytokine mRNA except for IL-10 and TNF- α , which were similar in both maternal and fetal tissues. These results suggest that the magnitude of the cytokine response correlates with but is not necessarily the cause of fetal death and demonstrate that a polarized Th1 response was not evident in the placentas of *N. caninum*-infected cattle.

Neospora caninum is a protozoan parasite that causes abortion in cattle worldwide (11). Cattle may abort following transplacental transmission of an infection acquired either exogenously by ingestion of oocysts from the feces of dogs or endogenously by recrudescence of the latent encysted bradyzoite stage of the parasite to the rapidly dividing tachyzoite stage (41). The mechanism by which *N. caninum* causes abortion is unknown, but an immune-mediated pathogenesis has been proposed (34).

The protective immune response to *N. caninum* in mice follows a T-helper 1 (Th1)-type profile, with the expression of interleukin-12 (IL-12) and gamma interferon (IFN- γ) being critical for survival (5, 19, 26). Similarly, the immune response in cattle, as investigated by in vitro studies of peripheral blood cells, is characterized by expression of high levels of IFN- γ (28, 44) by CD4⁺ T cells (1, 28, 44). Furthermore, IFN- γ inhibited intracellular multiplication of *N. caninum* in culture (17), and Staska et al. (39) showed killing of *N. caninum*-infected autol-

ogous targets by CD4⁺ cytotoxic T cells, indicating that these are likely control mechanisms in vivo.

Conversely, it has been suggested that the immune system in pregnancy is modulated so that the expression of Th2-type cytokines predominates (36, 42), and this bias is evident locally at the fetomaternal interface. Lin et al. (25) demonstrated the expression of high levels of IL-4 and IL-10 relative to IFN- γ at the fetomaternal interface of mice, and $\gamma\delta$ T cells isolated from human early pregnancy decidua secreted high levels of IL-10 and transforming growth factor β (TGF- β) relative to stimulated peripheral blood mononuclear cells (PBMC) (31). Furthermore, the expression of Th1-type cytokines and a deficiency of Th2-type cytokines at the fetomaternal interface has been associated with fetal loss in an abortion-susceptible mouse mating combination (9, 10), and defective production of Th2-type cytokines has been reported in decidual T cells in unexplained recurrent abortions (32). These data clearly suggest that the Th1-type immune response to a protozoan parasite during pregnancy might be detrimental to the health of the fetus, a theory that is borne out by studies in both mice and humans. After infection with *Leishmania major*, resistant C57/BL6 mice that express high levels of IFN- γ , IL-2, and tumor necrosis factor alpha (TNF- α) experience high rates of resorption (23), and the expression of Th1-type cytokines in response

* Corresponding author. Mailing address: Veterinary Parasitology, Liverpool School of Tropical Medicine, Pembroke Place, Liverpool L3 5QA, United Kingdom. Phone: 44 151 705 3142. Fax: 44 151 705 3373. E-mail: williadj@liv.ac.uk.

[†] Published ahead of print on 24 March 2008.

to placental malaria infection is associated with intrauterine fetal growth retardation (14, 29).

Thus, it is possible that *N. caninum* infection in pregnant cattle results in a Th1-type immune response at the fetomaternal interface that might precipitate fetal death. Experimental infections have shown that the time of a transplacental infection determines whether or not a cow aborts. An intravenous infection with *N. caninum* tachyzoites in early gestation results in fetal death, but the fetus survives a transplacental infection in late gestation, although it becomes infected (44). We have therefore used this system to compare the expression of Th1 (IL-2, IFN- γ , IL-12, TNF- α and IL-18), Th2 (IL-4), and regulatory (IL-10 and TGF- β 1) cytokines at the fetomaternal interface of cows whose fetuses died after *N. caninum* infection and in cows where transplacental transmission occurred and the fetus survived. We compared the relative contributions of maternal and fetal tissues to cytokine levels and correlated this with pathological alterations and inflammatory infiltration in the placenta. This is the first report describing the levels of both Th1 and Th2 cytokines at the fetomaternal interface of cattle following an experimental *N. caninum* infection.

MATERIALS AND METHODS

Animals. Twenty-four Friesian-Holstein maiden heifers were purchased commercially and confirmed negative for evidence of exposure to *N. caninum* by an antibody enzyme-linked immunosorbent assay (ELISA; Mastazyme; Mast Diagnostics, Liverpool, United Kingdom) (43, 45). Cattle were also confirmed negative for bovine viral diarrhea virus antigen and vaccinated against *Leptospira hardjo* (Leptavoid-H; Schering-Plough Animal Health, Uxbridge, United Kingdom), bovine viral diarrhea virus (Bovidec; Novartis Animal Health, Royston, United Kingdom), and infectious bovine rhinotracheitis virus (Bovilis IBR marker; Intervet, Milton Keynes, United Kingdom). After testing and vaccination, cattle were housed in dog- and fox-proof accommodations. Estrus was synchronized by using progesterone-releasing intravaginal devices (CEVA Animal Health, Chesham, United Kingdom), and cattle were artificially inseminated by using standard techniques, followed by housing with a bull. Pregnancy was confirmed by transrectal ultrasonography at approximately day 35 of gestation.

Parasite infection and postinfection monitoring of fetal viability. Cattle were divided into two groups of 12, and half of each group ($n = 6$) were infected at either day 70 or day 210 of gestation with 10^7 tachyzoites of *N. caninum* Nc Liverpool (3) in phosphate-buffered saline (pH 7.2). Tachyzoites were grown in Vero cells and harvested by passage through a 25-gauge needle to disrupt the Vero cells, followed by filtration through Sephadex PD10 columns (GE Healthcare, Amersham, United Kingdom) to remove Vero cell debris as described previously (45). Control animals ($n = 6$ per group) were infected with Vero cells treated in the same way as those that were infected with tachyzoites. After infection, fetal viability was monitored by ultrasonographic detection of the fetal heartbeat. Cattle infected on day 70 were ultrasound scanned rectally, twice weekly for the first 2 weeks postinfection and daily thereafter until fetal death; cattle infected on day 210 were scanned twice weekly by ultrasound transabdominally. Peripheral immune responses to *N. caninum* were analyzed as described previously (44, 45). Blood was collected into heparin-treated and plain tubes (Becton Dickinson, Oxford, United Kingdom) for analysis preinfection and 1, 2, and 3 weeks postinfection. PBMC were isolated by using Lymphoprep (density 1.077 g/ml; Axis-Shield, Oslo, Norway) and used to assay *N. caninum* antigen-specific proliferation. IFN- γ in the PBMC culture supernatants was measured by ELISA using a Bovigam IFN- γ ELISA kit (CSL, Victoria, Australia). Serum *N. caninum*-specific antibodies were measured by using a Mastazyme serum antibody ELISA (Mast Diagnostics).

Euthanasia and tissue collection. Euthanasia was carried out by using a captive bolt pistol, followed by pithing, in accordance with the United Kingdom Home Office License. Cattle infected on day 70 were euthanized 3 weeks after infection, at day 91 ± 1 of gestation (control animals), or within 24 h of detection of fetal death (infected animals). Cattle infected on day 210 were euthanized 3 weeks after infection, at day 231 ± 1 of gestation. Placental and fetal tissues were collected within 60 min of euthanasia. For analysis of cytokine mRNA expression by real-time reverse transcriptase PCR, full-thickness pieces of manually sepa-

FIG. 1. H&E-stained sections of uninfected bovine placenta. (A) Intact placentome. Black arrowheads indicate the maternal epithelium of a crypt containing fetal cotyledonary tissue. (B) Caruncle where the fetal cotyledon has been manually removed. White arrowheads indicate maternal endometrial epithelium surrounding a crypt with little remaining fetal material. (C) Fetal chorionic epithelium that has been manually removed from a placentome.

rated caruncle (maternal placenta) and cotyledon (fetal placenta) were placed in a 5 $\times$ volume of RNA-Later (Sigma, Poole, United Kingdom). For ELISA, the same tissues were stored at -70°C . To assess the separation, morphological examination of hematoxylin-and-eosin (H&E)-stained sections of caruncle and cotyledon was performed. Cotyledons consisted of 100% fetal material, and caruncles consisted of $>90\%$ maternal tissue with a small amount of fetal villus tissue remaining in the crypts (Fig. 1). For light microscopy, including immunohistology, full-thickness samples of placentomes were collected into 4% buffered paraformaldehyde (pH 7.4) or embedded in cryoembedding gel (OCT Tissue-Tek; Pelco International, Redding, United Kingdom) and snap-frozen in isopentane cooled in liquid nitrogen.

RNA extraction and reverse transcription. A total of 100 μg of tissue (caruncle or cotyledon) was homogenized by using a glass homogenizer into 1 ml of TRIreagent (Sigma), and total RNA was extracted as directed. Then, 1 μg of total RNA was reverse transcribed in a 20- μl volume containing 100 IU of Moloney murine leukemia virus RNase-H $^{-}$ point mutant reverse transcriptase

TABLE 1. Sequences of primers and probes used for real-time reverse transcription-PCR

Target	Primer orientation or probe ^a	Probe or primer sequence (5'-3') ^b	Exon boundary	Accession no. ^c
28S	P F R	AGTAGCTGGTTCCCTCCGAAGTTTCCCT GGCGAAAGACTAATCGAACCA CGAGAGCGCCAGCTATCCT		AF154866†
IL-2	P F R	AGCTCTCCAGGATGCATACATTTGACTTTTACG GGAGAAAGTTAAA[]AATCCTGAGAACCT TTCTAGTAAACACTTAAGATGTTTCAATTCTG	1-2	M12791†
IL-4	P F R	AGATTCCTGTCAAGTCCGCCAGG* GCCACACGTGCTTGAACAAA TCATTCACAGAACAGGT[]CTTGCTT	3-4	U14131
IL-10	P F R	TGCCAAGCCTTGTCGGAATGATCC TGCTGGATGACTTTAAG[]GGTTACC GGCATCACCTCTTCCAGGTA	2-3	U00799†
IFN- γ	P F R	TCATCCAC[]CGGAATTGGAATCAGCTTTTT* GCAGCTCTGAGAACTGGAGGA TGGCTTTGCGCTGGATCT	3-4	Z54144
TNF- α	P F R	CCTCAAGTAACAAGCCGGTAGCCCA TTCAAACACTCA[]GGTCCTCTTC GGAGAGTTGATGTCGG[]CTAC	2-3 3-4	AF011926
IL-12p40	P F R	ACTCTTAGCTTTGGGTT[]CTTTCTGATCCTTT* ATGGAATTTGGTCCACTGATATT GTGAAGTGTCAGAATAATCCTTT	2-3	U11815†
IL-18	P F R	TCCTTTAAG[]GAAATGAATCCTCCTGATAACA GTCTACTCTCTCCTGTGAGAACAA AATATGATGTCACTTTCTTCATTATCA	4-5	NM_174091†

^a P, probe. Primer orientation: F, forward; R, reverse.

^b *, The probe is reverse complement. The point where the primer or probe crosses the intron-exon boundary is indicated by "[]."

^c Accession numbers are for bovine genomic DNA unless indicated by "†," where they are for bovine cDNA since the bovine genomic DNA sequences are not available.

enzyme, a 0.5 mM deoxynucleoside triphosphate mix, 0.5 μ g of random primer, 20 IU of RNasin, and 1 $\times$ reaction buffer, resulting in a final concentration of 50 mM Tris-Cl (pH 8.3), 3 mM MgCl₂, 75 mM KCl, and 10 mM dithiothreitol (all from Promega, Southampton, United Kingdom). RNA was incubated with random primer for 5 min at 70°C and then cooled on ice before the addition of the remaining reagents and incubation for 10 min at 25°C, followed by 60 min at 42°C and a final denaturation for 10 min at 70°C. Reactions were diluted 1:5, and 5 μ l was used in the PCRs to analyze cytokine mRNA expression. Where cytokine expression was not detected, repeat reactions were analyzed at a dilution of 1:2. To analyze expression of the housekeeping gene 28S rRNA, reactions were diluted a further 1:100.

Primer design and real-time PCR. Primers and probes for 28S, IL-2, IL-4, and IFN- γ were designed by using Primer Express Software (PE Applied Biosystems, Warrington, United Kingdom). Those for TNF- α , IL-12p40, and IL-18 were designed by using Primer-3 software (<http://frodo.wi.mit.edu/cgi-bin/primer3/primer3 WWW.cgi>). Details of the primers and probes are given in Table 1. For all cytokines, one primer or probe crossed an intron-exon boundary to prevent detection of genomic DNA. Where information regarding intron-exon boundaries was unavailable for cattle (IL-2, IL-10, IL-12p40, and IL-18), intron-exon boundaries were deduced by comparison with mouse and human cDNA and amino acid sequences. Deduced boundaries were confirmed by PCR amplification of bovine genomic DNA using primers flanking either end of the suspected intron-exon boundary and sequencing of the PCR products. Cytokine and 28S probes were labeled with the fluorescent reporter dye 5-carboxyfluorescein (FAM) at the 5' end and with the quencher *N,N,N,N'*-tetramethyl-6-carboxyrhodamine (TAMRA) at the 3' end. Real-time PCR was performed by using a qPCR probe master mix (Eurogentec, Seraing, Belgium) in a 12.5- μ l volume containing 100 nM probe and primer concentrations individually determined by optimization experiments (Table 2). Amplification and detection of specific products was performed by using the Corbett Rotorgene 3000 (Corbett Research, Sydney,

Australia) with the following cycle profile: 96°C for 5 min, followed by 40 cycles of 94°C for 20 s and 59°C for 1 min. Quantification was based on increased fluorescence detected due to hydrolysis of the target-specific probes by the 5' nuclease activity of the *Taq* polymerase during PCR amplification. For each gene, a standard curve was constructed by using plasmids into which the full-length cDNA was cloned. A six-point duplicate standard curve was included in each run (Table 2), and from this an expression value in arbitrary units was calculated. Expression was then normalized to that of the housekeeping gene 28S rRNA, to give a relative expression value for each cytokine. All runs included duplicate no-template controls. Preliminary studies (data not shown) showed cytokine mRNA expression to be relatively consistent across 10 placentomes from one placenta; therefore, cytokine mRNA expression was measured in two caruncles (maternal) and one cotyledon (fetal) from each animal. For each cytokine in each tissue (cotyledon and caruncle), the median of the six control animals was calculated. The level of each cytokine in the cotyledon and caruncle in the two groups of infected cattle (day 70 and day 210) was then normalized to the median of the relevant control (uninfected) group for each tissue and each time point.

Measurement of TGF- β 1 and IFN- γ by ELISA. A total of 100 mg of frozen, separated cotyledon or caruncle was allowed to thaw in 2 ml of CellLyticMT cell lysis solution (Sigma) containing 5 μ l of protease inhibitor cocktail (Sigma) and then homogenized, and the supernatants were collected by centrifugation at 4°C for 10 min at 15,000 $\times$ g. The total protein in the supernatants was measured by using a Bio-Rad protein assay (Bio-Rad, Hemel Hempstead, United Kingdom) according to the manufacturer's instructions. TGF- β 1 was measured by using a TGF- β 1 E_{max} Immunoassay System (Promega, Southampton, United Kingdom) according to the manufacturer's instructions for detection of active TGF- β 1 (omitting the optional acid activation step). IFN- γ was measured by using a Bovigam IFN- γ ELISA kit as described previously (44).

TABLE 2. Standard curve data from six duplicate 10-fold dilutions of 1-µg/ml plasmid stocks^a

Target	Primer concn (µM)	Log dilutions	Mean C_T	Mean R^2	Mean slope	CV _{max} %	<i>n</i>
28S	0.3	10 ⁻¹ to 10 ⁻⁶	6.65–23.71	0.997	-3.43	3.26	6
IL-2	0.6	10 ⁻³ to 10 ⁻⁸	16.18–34.35	0.998	-3.52	2.57	6
IL-4	0.4	10 ⁻³ to 10 ⁻⁸	15.96–33.15	0.997	-3.61	3.43	8
IL-10	0.2	10 ⁻³ to 10 ⁻⁸	16.55–33.99	0.998	-3.61	4.64	7
IFN-γ	0.3	10 ⁻³ to 10 ⁻⁸	14.05–31.77	0.999	-3.54	2.25	8
TNF-α	0.4	10 ⁻³ to 10 ⁻⁸	13.23–29.24	0.998	-3.20	3.77	5
IL-12p40	0.4	10 ⁻³ to 10 ⁻⁸	15.09–32.47	0.999	-3.45	3.43	5
IL-18	0.6	10 ⁻³ to 10 ⁻⁸	15.86–33.52	0.998	-3.51	2.24	4

^a The mean C_T (i.e., the threshold value at which the change in reporter dye crosses the significance threshold), R^2 (coefficient of regression), and slope and the maximum coefficient of variation of all points (CV_{max}%) for *n* separate runs of each target are shown. The significance threshold for all runs was set at 0.01. In all reactions, the probe concentration was 100 nM; primer concentrations were individually determined for each gene and are as shown.

Light microscopy. Samples of placentomes fixed in paraformaldehyde for 24 to 48 h were routinely embedded into paraffin wax. Sections (3 to 5 µm thick) were prepared and either stained with H&E for evaluation of histological changes or used for immunohistology.

Immunohistology. T-cell (CD3, CD4, and CD8), B-cell (CD79a), and monocyte/macrophage (myeloid/histiocyte) antigens were detected by immunohistology in two placentomes from each animal. IFN-γ was detected by immunohistology in two placentomes from one animal infected on day 70 of gestation.

Staining for CD3, CD79a, myeloid/histiocyte antigen, and IFN-γ was performed on formalin-fixed and paraffin-embedded tissue sections by using anti-human polyclonal and monoclonal antibodies previously tested for their suitability in bovine tissues (16, 18). Endogenous peroxidase was inactivated for 30 min in 0.5% hydrogen peroxide in methanol, and sections were washed in Tris-buffered saline (TBS; 0.1 M Tris-HCl with 0.9% NaCl [pH 7.2]) before antigen retrieval (Table 3) and washing in TBS.

Staining for CD4 and CD8 antigens was performed on frozen sections (10 µM thick), which were air-dried before fixing for 10 min in ice-cold acetone. Endogenous peroxidase was inactivated as described above but with a 15-min incubation only.

Immunostaining was performed as follows. Sections were incubated for 15 to 18 h at 4°C with primary antibody (Table 3), washed for 5 min in TBS, and incubated for 30 min at room temperature with secondary antibody (Table 3). Binding of the secondary antibodies was detected by the Vectorstain ABC kit (Vector Laboratories, Peterborough, United Kingdom) according to the manufacturer's instructions or by using the rabbit or mouse PAP method as described previously (22). Slides were then incubated for 10 min, with stirring, with 3,3'-

diaminobenzidine tetrahydrochloride (DAB; Fluka, Buchs, Switzerland) with 0.01% hydrogen peroxide in 0.1 M imidazole buffer (pH 7.1), counterstained for 30 s in Papanicolaou's hematoxylin, and rinsed in running tap water before the slides were dehydrated, cleared, and mounted. Consecutive sections from each tissue were used as negative controls in which the primary antibodies were replaced by TBS. A bovine lymph node served as the positive control for each marker.

Statistical analysis. Statistical analysis of differences in cytokine expression was carried out with the Mann-Whitney U test using GraphPad Prism version 4.01 for Windows (GraphPad Software, San Diego, CA). The data are expressed as the median (plus range) or the mean ± the standard error of the mean (SEM). For the purpose of statistical analysis, samples in which cytokine expression was below detectable levels were assigned a value of 0.

RESULTS

Fetal death occurs in cattle infected with *N. caninum* at day 70 but not at day 210 of gestation. In animals infected at day 70 of gestation, fetal death occurred 22.7 ± 1.2 days later, and animals were euthanized for tissue collection within 24 h of detection of fetal death. No fetal death occurred prior to euthanasia of the dam in animals infected at day 210 of gestation. Peripheral immune responses to *N. caninum* were as described

TABLE 3. Antibodies and immunological methods^a

Primary antibody	Pretreatment	Blocking agent	Primary antibody dilution	Secondary antibody, dilution	Detection system ^b
Rabbit anti-human CD3	Protease ^d	50% SS	1:5 in 20% SS	Swine anti-rabbit immunoglobulin G, 1:100 in 20% SS	Rabbit PAP, 1:100 in 20% SS
Mouse anti-bovine CD4 (clone ILA-11)	None	Undiluted horse serum	1:10	Horse anti-mouse-biotin, 1:100	ABC
Mouse anti-bovine CD8 (clone IL-105)	None	Undiluted horse serum	1:10	Horse anti-mouse-biotin, 1:100	ABC
Mouse anti-human CD79a (clone HMS4)	Microwave ^e	Undiluted horse serum	1:50	Horse anti-mouse-biotin, 1:100	ABC
Rat anti-human myeloid/histiocyte antigen (clone MAC387)	Protease ^d	10% rat serum	1:100	Rat anti-mouse immunoglobulin G ^g , 1:100	Mouse PAP, 1:500 ^g
Mouse anti-bovine IFN-γ ^c (clone MCA1964)	Citrate buffer ^f	10% rat serum	1:50	Rat anti-mouse immunoglobulin G ^g , 1:100	Mouse PAP, 1:500 ^g

^a Sera and antibodies were diluted in TBS. Unless otherwise indicated, antibodies were from DakoCytomation, Glostrup, Denmark, with the exception of ILA-11 and IL-105, which were supernatants from monoclonal antibodies cultured in-house. SS, swine serum.

^b PAP, peroxidase anti-peroxidase method; ABC, avidin-biotin-peroxidase complex method.

^c Obtained from Serotec, Oxford, United Kingdom.

^d That is, treatment with 0.05% protease (type XXIV, bacterial; Sigma) for 5 min at 37°C.

^e That is, treatment for 10 min at 800 W in 10 mM EDTA (pH 9.0).

^f That is, treatment for 30 min at 97°C in citrate buffer (pH 6.0).

^g Obtained from Jackson ImmunoResearch Laboratories, West Grove, PA.

FIG. 2. Cytokine mRNA expression in placentas approximately 3 weeks after intravenous infection of cattle with 10^7 *N. caninum* tachyzoites at day 70 of gestation. Expression in the maternal caruncles (A) and fetal cotyledons (B) is shown. The data are represented as individual points for each of six control and six infected animals, although it is not possible to distinguish all individual points due to their close proximity. Expression below detectable levels (IL-4, four controls; IL-12p40, four controls; IL-2, three controls and one infected animal) was assigned as "0" for statistical purposes but omitted from the graphs. Statistical analysis data (infected compared to controls) is omitted from graphs A and B for clarity and indicated as follows: A, $P < 0.01$, all cytokines; and B, $P < 0.01$, all cytokines except IL-2 (not statistically significant).

previously (44), confirming infection with viable parasites at both days 70 and 210 of gestation. PBMC from infected animals proliferated and secreted IFN- γ in response to *N. caninum* antigen from 1 week postinfection and seroconverted by 2 weeks postinfection (data not shown). An aliquot of the same parasites used for infection of cattle was used to infect Vero cells. Parasite growth was observed after 5 days, confirming parasite viability.

All cytokine mRNA is greatly increased in the bovine placenta after infection with *N. caninum* at day 70 of gestation. To investigate the possible role of cytokines in fetal death after experimental infection of cattle with *N. caninum* at day 70 of gestation, we measured the levels of IL-2, IFN- γ , IL-4, IL-10, IL-12p40, TNF- α , and IL-18 mRNA in bovine placentas. In maternal tissue (caruncles), all cytokine mRNA was significantly ($P < 0.01$) increased compared to that in uninfected animals (Fig. 2A). IL-4 and IFN- γ were most highly upregulated by 228 (110 to 1,729)- and 785 (287 to 3,641)-fold, respectively. Also highly upregulated were the IL-12p40, IL-2, and TNF- α mRNA, whose levels increased by 33 (10 to 215)-, 50 (15 to 221), and 55 (13 to 391)-fold, respectively. IL-10 mRNA increased by 15 (7 to 97)-fold, and IL-18 mRNA was upregulated 7 (5 to 25)-fold. Cytokine mRNA was also signif-

FIG. 3. Percent contributions of caruncle and cotyledon to the total relative expression (caruncle plus cotyledon) of cytokine mRNA in the placenta after *N. caninum* infection at day 70 of gestation. The mean plus the SEM ($n = 6$) was plotted. **, $P < 0.01$, cotyledon compared to caruncle.

icantly ($P < 0.01$) upregulated in the fetal cotyledon after *N. caninum* infection at day 70 of gestation, with the exception of IL-2 (Fig. 2B). The most highly upregulated were IL-10, IFN- γ , and TNF- α mRNA, which increased by 138 (75 to 479)-, 156 (20 to 10,508)-, and 190 (22 to 3,957)-fold, respectively. IL-18, IL-4, and IL-12p40 mRNA were upregulated by 9 (5 to 12)-, 6 (3 to 264)-, and 3 (2 to 49)-fold, respectively. In general, cytokine levels in cotyledons were lower and more variable than in caruncles; increases in IL-4 and IL-12p40 mRNA were detected in only two of the six controls, and IL-2 mRNA was detected in only three controls and five infected cotyledons.

In cattle infected at day 70 of gestation, all cytokine mRNA levels except for IL-10 and TNF- α were significantly higher in the maternal side than in the fetal side of the placenta. Further comparison of the levels of cytokine mRNA in the caruncles and cotyledons after *N. caninum* infection at day 70 was undertaken to determine the relative contributions of maternal and fetal tissues to the responses seen. The levels of IL-12p40, IL-2, IL-4, IFN- γ , and IL-18 were significantly higher ($P < 0.01$) in caruncles than in cotyledons (Fig. 3). Caruncles had $99\% \pm 0.3\%$, $94\% \pm 3\%$, $93\% \pm 5\%$, $92\% \pm 5\%$, and $86\% \pm 3\%$ of the total amounts of each respective cytokine mRNA. In contrast, caruncles had $56\% \pm 9\%$ and $61\% \pm 8\%$ of the total IL-10 and TNF- α mRNA, respectively, and there was no significant difference between the levels in caruncles and the levels in cotyledons.

All cytokine mRNA is modestly increased in the placenta after infection of cattle at day 210 of gestation. To compare placental cytokine levels when *N. caninum* infection results in fetal death to that when the fetus survives infection, we examined the quantity of cytokine mRNA at the fetomaternal interface 3 weeks after infection of cattle at day 210 of gestation. All cytokine mRNA levels except for IL-2 were significantly ($P < 0.01$ for all except IL-18, for which $P < 0.05$) increased in the maternal caruncle, but to a much lesser extent than that seen after infection at day 70 (Fig. 4A). As seen after infection at day 70, IL-4 and IFN- γ mRNA were the most highly upregulated, but by only 10.1 (1.1 to 17.8)- and 16.3 (6.7 to 65.2)-fold, respectively. IL-12p40 and TNF- α mRNA were upregulated with respective increases of 3.5 (1.8 to 15.4)- and 2.7 (1.6 to 3.4)-fold, and IL-10 and IL-18 mRNA levels increased by 2.0 (1.6 to 3.4)- and 1.4 (1.2 to 2.0)-fold, respectively. Cy-

FIG. 4. Cytokine mRNA expression in placentae approximately 3 weeks after intravenous infection of cattle with 10^7 *N. caninum* tachyzoites at day 210 of gestation. Expression in the maternal caruncles (A) and fetal cotyledons (B) is shown. The data are represented as individual points for each of six control and six infected animals, although it is not possible to distinguish all individual points due to their close proximity. Expression below detectable levels (IL-4, three controls; IL-10, one control and two infected animals; IFN- γ , one control; IL-2, three controls) was assigned as "0" for statistical purposes but omitted from the graphs. Statistical analysis data (infected compared to controls) is omitted from graphs A and B for clarity and indicated as follows: A, $P < 0.01$ except for IL-18, where $P < 0.05$ (IL-2 is not statistically significant); and B, $P < 0.05$, IL-4 and IFN- γ only.

tokine mRNA in the cotyledons was at the limit of detection and, as such, variable (Fig. 4B). Nevertheless, IL-4 and IFN- γ mRNA levels were significantly ($P < 0.05$) increased, and there was a trend toward significance in the increase in TNF- α mRNA in cotyledons. IL-2 and IL-4 mRNA levels were undetectable in three control samples. IL-10 mRNA was undetectable in one control and two infected cotyledon samples, and IFN- γ mRNA was undetectable in one control sample.

A similar Th1/Th2 cytokine ratio is present in the placenta after *N. caninum* infection at days 70 and 210 of gestation. We compared the placental cytokine balance as a ratio of IFN- γ to IL-4 after *N. caninum* infection at day 70 or 210 of gestation. The ratios of IFN- γ to IL-4 in the caruncles were 2.9 ± 0.25 and 2.9 ± 0.77 after infection at days 70 and 210, respectively; there was no significant difference between the two.

Active TGF- β 1 is present in the bovine placenta and protein levels increased after infection at day 70 of gestation. Since TGF- β 1 protein levels correspond poorly to mRNA levels (24), active TGF- β 1 in bovine placentas was measured by ELISA. TGF- β 1 was detectable both at days 70 and 210 of gestation (Fig. 5), and the levels increased after infection at

FIG. 5. Expression of the active form of TGF- β 1 in the placentas of cattle 3 weeks after an intravenous infection with *N. caninum* at day 70 or day 210 of gestation. TGF- β 1 in extracts from separated caruncles and cotyledons was measured by ELISA. The data are shown as means plus the SEM ($n = 6$). Statistically significant differences between groups indicated are shown as follows: *, $P < 0.05$; and **, $P < 0.01$.

day 70; this increase was significant ($P < 0.05$) in caruncles (Fig. 5A). At day 70 of gestation, the expression of TGF- β 1 was higher in cotyledons than caruncles, but this difference was only significant ($P < 0.05$) in control animals (Fig. 5A). The expression of TGF- β 1 at day 210 of gestation was similar in cotyledons and caruncles, and there was no increase after infection (Fig. 5B).

The expression of IFN- γ protein is increased in the placenta after *N. caninum* infection at days 70 and 210 of gestation. To determine whether changes in mRNA expression correlated with levels of protein, the levels of placental IFN- γ were measured by ELISA. Very low levels of IFN- γ protein were detectable in uninfected placentas (0.0025 ± 0.0004 and 0.004 ± 0.001 ng/mg protein in caruncles of day 70 and day 210 control cattle, respectively) (Fig. 6). However, the levels of protein increased significantly ($P < 0.01$) after infection at both days 70 and 210 of gestation, with fold increases in proportion to those seen for the expression of mRNA. IFN- γ levels in caruncles reached 2.1 ± 0.9 and 0.13 ± 0.07 ng/mg after infection at days 70 and 210, respectively.

***N. caninum* infection of cattle at day 70 of gestation results in extensive necrosis, placental infiltration of CD4 $^{+}$ T cells and macrophages, and expression of IFN- γ in necrotic areas.** To determine whether the observed increases in cytokine levels were associated with pathological changes and recruitment of particular cell types into the placenta, histological examination and immunohistological staining for T cells, B cells, and mono-

FIG. 6. Expression of IFN- γ in placentas 3 weeks after an intravenous infection of cattle with *N. caninum* at day 70 or day 210 of gestation. IFN- γ protein in extracts from separated caruncle and cotyledon was measured by ELISA. The data are shown as means plus the SEM ($n = 6$). **, $P < 0.001$, infected compared to controls.

cytes/macrophages were performed. We also used immunohistology to localize the cellular source of IFN- γ . In uninfected control animals, the villus epithelium was intact, and the presence of leukocytes was restricted to occasional T and B cells in placentomes and, in the day 210 controls, scattered macrophages in the maternal interstitium. After infection at day 70 of gestation, extensive, multifocal villus epithelial cell necrosis was present in all placentomes of all animals (Fig. 7A). Macrophages were numerous around blood vessels at the base of caruncles and in association with villus epithelial cell necrosis but were also present in moderate numbers throughout the interstitium (Fig. 7B). There was an intense diffuse lymphocyte infiltration that was more pronounced at the bases of the caruncles (Fig. 7A). It was composed of CD3⁺ T cells (Fig. 7C), the majority of which were CD4⁺ T cells (Fig. 7D). Expression of IFN- γ was observed in maternal epithelial cells, cell-free within necrotic areas, and in mononuclear cells adjacent to areas of necrosis (Fig. 7E and F). After infection at day 210 of gestation, histological changes were restricted to occasional small areas of villus epithelial necrosis. In general, no more than 2 of the 10 placentomes examined were affected, and each placentome had fewer than three or four small areas of necrosis. Again, macrophages were observed in association with the necrotic areas. Scattered macrophages were also present in the maternal interstitium in numbers similar to those observed in day 210 control animals. CD3⁺ T cells, which consisted of both CD4⁺ and CD8⁺ cells, were present in low numbers, sparsely disseminated throughout the placentomes and maternal interstitium, and more numerous around ne-

crotic areas. Sparsely scattered CD79a⁺ B cells were present in placentas collected at both time points, and there were no differences between control and infected animals.

DISCUSSION

Here we report for the first time a dramatic increase in the quantities of a number of cytokines at the fetomaternal interface among cattle in which the fetus died after an experimental infection with *N. caninum* tachyzoites in early gestation. In contrast, cytokine mRNA levels were only modestly increased at the fetomaternal interface of cattle infected in late gestation in which the fetuses survived. The cytokines that were upregulated included those implicated as potential abortifacients, such as IFN- γ (20, 23), TNF- α (13, 23), IL-12p40 (46), and IL-2 (40). However, the levels of Th2 and regulatory cytokines, including IL-4, IL-10, and TGF- β 1, also increased in the placenta after *N. caninum* infection. The cytokine balance, as indicated by the IFN- γ /IL-4 ratio, was similar after infection in early and late gestation, but the magnitude of expression of all cytokines was very different, with strong cytokine expression occurring in the placentomes of cattle whose fetuses were killed as a result of infection.

These results, showing an increased expression of both Th1 and Th2 cytokines after *N. caninum* infection, contrast with previous suggestions that a Th1-biased response to *N. caninum* occurs in infected cattle (27). However, the data are consistent with studies in *N. caninum*-infected mice for which a mixed systemic cytokine response has been described (19, 26, 35) and recent work from our laboratory showing an increase of both IFN- γ and IL-4 mRNA in PBMC in response to *N. caninum* infection of cattle (37). Moreover, in vitro work suggests that an immune response polarized toward Th1 or Th2 is uncommon in cattle (7). Significantly, we have shown that here that the placental cytokine balance in infected cattle, after infection in early and late gestation, does not support the traditional view that the cytokine environment of pregnancy is biased toward the expression of Th2 type cytokines (36, 42).

There was widespread necrosis and inflammation of the placenta in the day 70 group that was absent in the day 210 group. We have shown elsewhere (15) that this necrosis predominantly affects the cotyledons and is associated with high numbers of parasites. Moreover, the day 70 fetal tissues (notably the heart, liver, and neural tissues) showed evidence of extensive parasitosis, whereas it was minimal in the day 210 group. It is likely that the placental necrosis observed in the day 70 animals contributed to fetal death, but why there is such a difference in the extent of parasite dissemination and necrosis in the day 70 group compared to the day 210 group is not clear, particularly since both groups were infected with the same dose of parasites and killed at the same time (3 weeks) after infection. It is possible that the high levels of IFN- γ produced in the placenta in response to the presence of the parasite are pathogenic (36, 38, 40, 47) or, alternatively, that the necrosis is due directly to parasite mediated cell damage, in which case the immunocompetence of the fetus itself may be significant. Fetal immunocompetence develops gradually through gestation, but *N. caninum* antigen specific fetal responses have only been described from midgestation onward (1, 4). Thus, it is likely that after the parasite has crossed the

FIG. 7. Bovine placenta after challenge with *N. caninum* at day 70 of gestation induced fetal death. (A) Multiple small areas of villus epithelial cell necrosis (arrows), associated with mononuclear infiltration and numerous lymphocytes in the maternal interstitium. H&E stain was used. (B) Myeloid/histiocyte antigen-positive macrophages are the dominant cells in association with focal necrosis (arrow). (C) The majority of infiltrating cells in the maternal interstitium are CD3⁺ T cells (arrows). N, focal area of necrosis. (D) The majority of infiltrating lymphocytes are CD4⁺. (E) IFN- γ is expressed weakly by maternal epithelial cells (short arrows) and is seen cell-free (arrows) in areas of necrosis. (F) Individual mononuclear cells (the arrow indicates a lymphocyte) in the maternal interstitium adjacent to areas of necrosis express IFN- γ . For panels B to F, immunohistology was performed using the peroxidase anti-peroxidase method with Papanicolaou's hematoxylin counterstain.

placenta and invaded fetal tissues, its multiplication and dissemination is controlled in the immunologically mature day 210 fetus, whereas the day 70 fetus becomes heavily parasitized. We have suggested elsewhere (15) that this results in large numbers of parasites reinvading the placenta via the fetal circulation in the day 70 group, stimulating a profound inflammatory response in the placenta. This may ultimately cause

fetal death either by the direct cytotoxic effects of proinflammatory cytokines and immune CD4⁺ T cells or by parasite-mediated destruction of fetal trophoblast cells. Both of these mechanisms could potentially lead to placental insufficiency and fetal death in a way similar to that proposed for *Toxoplasma gondii* (8).

By analyzing the levels of cytokines in maternal caruncle

compared to fetal cotyledon tissue we showed that, with the exception of IL-10 and TNF- α , the majority of the cytokines detected were of maternal origin. For IL-12, IL-2, IL-4, and IFN- γ , <8% of the total mRNA detected was in the cotyledons. Nevertheless, for the day 70 group IFN- γ , TNF- α , IL-4, and IL-10 were significantly upregulated in the infected cotyledon compared to the control cotyledon tissue. It is possible that these cytokines were expressed by the fetal trophoblast cells directly or by infiltrating maternal mononuclear cells. We showed by immunohistology that IFN- γ -producing, maternally derived mononuclear cells were present in areas of damaged villous epithelium and so could be the source, particularly of IL-4 and IFN- γ . However, there is evidence that fetal trophoblast cells can express IL-10 and TNF- α (6, 21), and therefore these cytokines could have been either of fetal origin or expressed by infiltrating maternal macrophages, which have also been shown to produce IL-10 and TNF- α (33).

By immunohistology we showed there was an influx of CD4⁺ T cells, probably of maternal origin, into the caruncle; this influx was most marked in the day 70 group. These cells were probably a significant source of cytokines, and we showed by immunohistology that they expressed IFN- γ , corroborating the work of Maley et al. (27). CD4⁺ cytotoxic T cells have been described in *N. caninum*-infected cattle (39). We also show for the first time that IFN- γ was also upregulated in maternal caruncle epithelium. These observations suggest that the increase in cytokine expression and recruitment of macrophages and CD4⁺ T cells to the placenta were to control parasite multiplication. Similarly, in mice infected with *Listeria monocytogenes*, upregulation of IL-12, TNF- α , IL-18, and IFN- γ in the placenta is thought to be essential for effective control of the bacterium and in preventing abortion associated with *L. monocytogenes* infection (2).

Active TGF- β 1 protein was present in control and infected fetal and maternal tissue at both days 70 and 210 of gestation, confirming the results of a previous study (30) and the suggestion that it is involved in placental remodeling (12). In the cattle infected at day 70, higher levels of TGF- β 1 were detected in the cotyledons in both uninfected and infected cattle compared to the day 210 group, but there was a significant increase in expression of TGF- β in the day 70 caruncles after infection. Regulatory T cells have been implicated in controlling the immunological microenvironment at the maternofetal interface, and TGF- β is a signature cytokine for this population (48). It is possible that the expression of TGF- β is upregulated to control the recruitment of cytotoxic T cells that could potentially harm the fetus.

In conclusion, *N. caninum* infection in early gestation, which resulted in fetal death, was associated with a much greater increase in placental cytokine expression, placental necrosis, and inflammatory cell infiltration compared to cattle infected later in gestation when the fetus survived. There was no evidence of a marked polarization of the immune response in the placenta, and the expression of Th1 (including, most markedly, IFN- γ), Th2, and regulatory cytokines increased. Fetal death could have resulted from the fetocytotoxic effects of cytokines such as IFN- γ , TNF- α , or IL-2, or the immune response in the placenta could have been generated to protect the conceptus from direct parasite-mediated cell damage.

ACKNOWLEDGMENTS

We are grateful to the farm staff at the University Animal Husbandry Farm for their dedicated care of the animals used in this study and to the technical staff at the Histology Laboratories, Department of Veterinary Pathology, for performing routine processing of tissues. We also thank D. Bainbridge, Department of Veterinary Anatomy, University of Cambridge, for helpful discussions.

This study was funded by the Wellcome Trust (grant GR066695MA), and E.H.G. was supported by a Postgraduate Research Studentship from the University of Liverpool and by Novartis Animal Health, Ltd.

REFERENCES

- Andrianarivo, A. G., B. C. Barr, M. L. Anderson, J. D. Rowe, A. E. Packham, K. W. Sverlow, and P. A. Conrad. 2001. Immune responses in pregnant cattle and bovine fetuses following experimental infection with *Neospora caninum*. *Parasitol. Res.* 87:817–825.
- Barber, E. M., M. Fazzari, and J. W. Pollard. 2005. Th1 cytokines are essential for placental immunity to *Listeria monocytogenes*. *Infect. Immun.* 73:6322–6331.
- Barber, J. S., O. J. Holmdahl, M. R. Owen, F. Guy, A. Ugglä, and A. J. Trees. 1995. Characterization of the first European isolate of *Neospora caninum* (Dubey, Carpenter, Speer, Topper, and Ugglä). *Parasitology* 111(Pt. 5):563–568.
- Bartley, P. M., E. Kirvar, S. Wright, C. Swales, I. Esteban-Redondo, D. Buxton, S. W. Maley, A. Schock, A. G. Rae, C. Hamilton, and E. A. Innes. 2004. Maternal and fetal immune responses of cattle inoculated with *Neospora caninum* at mid-gestation. *J. Comp. Pathol.* 130:81–91.
- Baszler, T. V., M. T. Long, T. F. McElwain, and B. A. Mathison. 1999. Interferon-gamma and interleukin-12 mediate protection to acute *Neospora caninum* infection in BALB/c mice. *Int. J. Parasitol.* 29:1635–1646.
- Bennett, W. A., S. Lagoo-Deenadayan, N. S. Whitworth, M. N. Brackin, E. Hale, and B. D. Cowan. 1997. Expression and production of interleukin-10 by human trophoblast: relationship to pregnancy immunotolerance. *Early Pregnancy* 3:190–198.
- Brown, W. C., A. C. Rice-Ficht, and D. M. Estes. 1998. Bovine type 1 and type 2 responses. *Vet. Immunol. Immunopathol.* 63:45–55.
- Buxton, D., J. S. Gilmour, K. W. Angus, D. A. Blewett, and J. K. Miller. 1982. Perinatal changes in lambs infected with *Toxoplasma gondii*. *Res. Vet. Sci.* 32:170–176.
- Chaouat, G., M. A. Assal, J. Martal, R. Raghupathy, J. F. Elliott, T. Mosmann, and T. G. Wegmann. 1995. IL-10 prevents naturally occurring fetal loss in the CBA $\times$ DBA/2 mating combination, and local defect in IL-10 production in this abortion-prone combination is corrected by in vivo injection of IFN- γ . *J. Immunol.* 154:4261–4268.
- Chaouat, G., E. Menu, D. A. Clark, M. Dy, M. Minkowski, and T. G. Wegmann. 1990. Control of fetal survival in CBA $\times$ DBA/2 mice by lymphokine therapy. *J. Reprod. Fertil.* 89:447–458.
- Dubey, J. P., and D. S. Lindsay. 1996. A review of *Neospora caninum* and neosporosis. *Vet. Parasitol.* 67:1–59.
- Dungy, L. J., T. A. Siddiqi, and S. Khan. 1991. Transforming growth factor- β 1 expression during placental development. *Am. J. Obstet. Gynecol.* 165:853–857.
- Fest, S., A. C. Zenclussen, R. Joachim, E. Hagen, H. U. Demuth, and T. Hoffmann. 2006. Stress and substance P but not the substance P-metabolite SP5-11 trigger murine abortion by augmenting TNF- α levels at the fetomaternal interface. *Scand. J. Immunol.* 63:42–49.
- Fried, M., R. O. Muga, A. O. Misore, and P. E. Duffy. 1998. Malaria elicits type 1 cytokines in the human placenta: IFN- γ and TNF- α associated with pregnancy outcomes. *J. Immunol.* 160:2523–2530.
- Gibney, E. H., A. Kipar, A. Rosbottom, C. S. Guy, R. F. Smith, U. Hetzel, A. J. Trees, and D. J. L. Williams. 2008. The extent of parasite-associated necrosis in the placenta and foetal tissues of cattle following *N. caninum* infection in early and late gestation correlates with foetal death. *Int. J. Parasitol.* 38:579–588.
- Gutierrez, M., F. I. Forster, S. A. McConnell, J. P. Cassidy, J. M. Pollock, and D. G. Bryson. 1999. The detection of CD2⁺, CD4⁺, CD8⁺, and WC1⁺ T lymphocytes, B cells and macrophages in fixed and paraffin embedded bovine tissue using a range of antigen recovery and signal amplification techniques. *Vet. Immunol. Immunopathol.* 71:321–334.
- Innes, E. A., W. R. Pantton, J. Marks, A. J. Trees, J. Holmdahl, and D. Buxton. 1995. Interferon gamma inhibits the intracellular multiplication of *Neospora caninum*, as shown by incorporation of 3H uracil. *J. Comp. Pathol.* 113:95–100.
- Johnson, L., J. Gough, Y. Spencer, G. Hewinson, M. Vordermeier, and A. Wangoo. 2006. Immunohistochemical markers augment evaluation of vaccine efficacy and disease severity in bacillus Calmette-Guerin (BCG) vaccinated cattle challenged with *Mycobacterium bovis*. *Vet. Immunol. Immunopathol.* 111:219–229.
- Khan, I. A., J. D. Schwartzman, S. Fonseka, and L. H. Kasper. 1997. *Neo-*

- spora caninum*: role for immune cytokines in host immunity. Exp. Parasitol. 85:24–34.
20. Kim, S., D. S. Lee, K. Watanabe, H. Furuoka, H. Suzuki, and M. Watarai. 2005. Interferon-gamma promotes abortion due to *Brucella* infection in pregnant mice. BMC. Microbiol. 5:22.
 21. King, A., P. P. Jokhi, S. K. Smith, A. M. Sharkey, and Y. W. Loke. 1995. Screening for cytokine mRNA in human villous and extravillous trophoblasts using the reverse-transcriptase polymerase chain reaction (RT-PCR). Cytokine 7:364–371.
 22. Kipar, A., S. Bellmann, J. Kremendahl, K. Kohler, and M. Reinacher. 1998. Cellular composition, coronavirus antigen expression and production of specific antibodies in lesions in feline infectious peritonitis. Vet. Immunol. Immunopathol. 65:243–257.
 23. Krishnan, L., L. J. Guilbert, T. G. Wegmann, M. Belosevic, and T. R. Mosmann. 1996. T helper 1 response against *Leishmania major* in pregnant C57BL/6 mice increases implantation failure and fetal resorptions. Correlation with increased IFN- γ and TNF and reduced IL-10 production by placental cells. J. Immunol. 156:653–662.
 24. Li, M. O., Y. Y. Wan, S. Sanjabi, A. K. Robertson, and R. A. Flavell. 2006. Transforming growth factor-beta regulation of immune responses. Annu. Rev. Immunol. 24:99–146.
 25. Lin, H., T. R. Mosmann, L. Guilbert, S. Tuntipopipat, and T. G. Wegmann. 1993. Synthesis of T helper 2-type cytokines at the maternal-fetal interface. J. Immunol. 151:4562–4573.
 26. Long, M. T., T. V. Baszler, and B. A. Mathison. 1998. Comparison of intracerebral parasite load, lesion development, and systemic cytokines in mouse strains infected with *Neospora caninum*. J. Parasitol. 84:316–320.
 27. Maley, S. W., D. Buxton, C. N. Macaldowie, I. E. Anderson, S. E. Wright, P. M. Bartley, I. Esteban-Redondo, C. M. Hamilton, A. K. Storset, and E. A. Innes. 2006. Characterization of the immune response in the placenta of cattle experimentally infected with *Neospora caninum* in early gestation. J. Comp. Pathol. 135:130–141.
 28. Marks, J., A. Lunden, D. Harkins, and E. Innes. 1998. Identification of *Neospora* antigens recognized by CD4⁺ T cells and immune sera from experimentally infected cattle. Parasite Immunol. 20:303–309.
 29. Moormann, A. M., A. D. Sullivan, R. A. Rochford, S. W. Chensue, P. J. Bock, T. Nyirenda, and S. R. Meshnick. 1999. Malaria and pregnancy: placental cytokine expression and its relationship to intrauterine growth retardation. J. Infect. Dis. 180:1987–1993.
 30. Munson, L., A. Wilhite, V. F. Boltz, and J. E. Wilkinson. 1996. Transforming growth factor beta in bovine placentas. Biol. Reprod. 55:748–755.
 31. Nagaeva, O., L. Jonsson, and L. Mincheva-Nilsson. 2002. Dominant IL-10 and TGF- β mRNA expression in γ 8T cells of human early pregnancy decidua suggests immunoregulatory potential. Am. J. Reprod. Immunol. 48:9–17.
 32. Piccinni, M. P., L. Beloni, C. Livi, E. Maggi, G. Scarselli, and S. Romagnani. 1998. Defective production of both leukemia inhibitory factor and type 2 T-helper cytokines by decidual T cells in unexplained recurrent abortions. Nat. Med. 4:1020–1024.
 33. Pyeon, D., K. L. O'Reilly, and G. A. Splitter. 1996. Increased interleukin-10 mRNA expression in tumor-bearing or persistently lymphocytotic animals infected with bovine leukemia virus. J. Virol. 70:5706–5710.
 34. Quinn, H. E., J. T. Ellis, and N. C. Smith. 2002. *Neospora caninum*: a cause of immune-mediated failure of pregnancy? Trends Parasitol. 18:391–394.
 35. Quinn, H. E., C. M. Miller, and J. T. Ellis. 2004. The cell-mediated immune response to *Neospora caninum* during pregnancy in the mouse is associated with a bias toward production of interleukin-4. Int. J. Parasitol. 34:723–732.
 36. Raghupathy, R. 1997. Th1-type immunity is incompatible with successful pregnancy. Immunol. Today 18:478–482.
 37. Rosbottom, A., C. S. Guy, E. H. Gibney, R. F. Smith, J. F. Valarcher, G. Taylor, and D. J. Williams. 2007. Peripheral immune responses in pregnant cattle following *Neospora caninum* infection. Parasite Immunol. 29:219–228.
 38. Sollwedel, A., A. Z. Bertoja, M. L. Zenclussen, K. Gerlof, U. Lisewski, P. Wafula, B. Sawitzki, C. Woiciechowsky, H. D. Volk, and A. C. Zenclussen. 2005. Protection from abortion by heme oxygenase-1 up-regulation is associated with increased levels of Bag-1 and neuropilin-1 at the fetal-maternal interface. J. Immunol. 175:4875–4885.
 39. Staska, L. M., T. C. McGuire, C. J. Davies, H. A. Lewin, and T. V. Baszler. 2003. *Neospora caninum*-infected cattle develop parasite-specific CD4⁺ cytotoxic T lymphocytes. Infect. Immun. 71:3272–3279.
 40. Tezabwala, B. U., P. M. Johnson, and R. C. Rees. 1989. Inhibition of pregnancy viability in mice following IL-2 administration. Immunology 67:115–119.
 41. Trees, A. J., and D. J. Williams. 2005. Endogenous and exogenous transplacental infection in *Neospora caninum* and *Toxoplasma gondii*. Trends Parasitol. 21:558–561.
 42. Wegmann, T. G., H. Lin, L. Guilbert, and T. R. Mosmann. 1993. Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon? Immunol. Today 14:353–356.
 43. Williams, D. J., H. C. Davison, B. Helmick, J. McGarry, F. Guy, A. Otter, and A. J. Trees. 1999. Evaluation of a commercial ELISA for detecting serum antibody to *Neospora caninum* in cattle. Vet. Rec. 145:571–575.
 44. Williams, D. J., C. S. Guy, J. W. McGarry, F. Guy, L. Tasker, R. F. Smith, K. MacEachern, P. J. Cripps, D. F. Kelly, and A. J. Trees. 2000. *Neospora caninum*-associated abortion in cattle: the time of experimentally-induced parasitaemia during gestation determines fetal survival. Parasitology 121(Pt. 4):347–358.
 45. Williams, D. J., J. McGarry, F. Guy, J. Barber, and A. J. Trees. 1997. Novel ELISA for detection of *Neospora*-specific antibodies in cattle. Vet. Rec. 140:328–331.
 46. Zenclussen, A. C., R. Joachim, E. Hagen, C. Peiser, B. F. Klapp, and P. C. Arck. 2002. Heme oxygenase is downregulated in stress-triggered and interleukin-12-mediated murine abortion. Scand. J. Immunol. 55:560–569.
 47. Zenclussen, A. C., A. Sollwedel, A. Z. Bertoja, K. Gerlof, M. L. Zenclussen, C. Woiciechowsky, and H. D. Volk. 2005. Heme oxygenase as a therapeutic target in immunological pregnancy complications. Int. Immunopharmacol. 5:41–51.
 48. Zenclussen, A. C., K. Gerlof, M. L. Zenclussen, S. Ritschel, A. Z. Bertoja, S. Fest, S. Hontsu, S. Ueha, K. Matshushima, J. Leber, and H. D. Volk. 2006. Regulatory T cells induce a privileged tolerant microenvironment at the fetal-maternal interface. Eur. J. Immunol. 36:82–94.