Published in final edited form as:

J Acquir Immune Defic Syndr. 2006 April 15; 41(5): 642–650.

Sexual Risk Reduction for Persons Living with HIV: Research Synthesis of Randomized Controlled Trials, 1993–2004

Blair T. Johnson, PhD^a, Michael P. Carey, PhD^b, Stephenie R. Chaudoir, BA^a, and Allecia E. Reid, BA^c

^a Center for Health/HIV Intervention and Prevention, University of Connecticut, Storrs, CT

^b Center for Health and Behavior, Syracuse University, Syracuse, NY

^c Department of Psychology, Arizona State University, Tempe, AZ

Abstract

Objective—To conduct a meta-analytic review of interventions to reduce HIV+ individuals' sexual risk.

Design—Studies were included if they examined a deliberate sexual-risk-reduction strategy in a sample that included HIV+ participants; used a randomized controlled trial design; measured condom use or number of sexual partners following the intervention; and provided sufficient information to calculate effect size (ES) estimates.

Method—Reports were gathered from computerized databases, by contacting individual researchers, by searching relevant journals and conference proceedings, and by reviewing reference sections of obtained papers. Data from 15 studies (N=3,234 participants) available as of November 30, 2004 were included. Effect sizes (ES) were standardized mean differences.

Results—Across the studies, intervention participants exhibited lowered sexual risk relative to control participants on condom use (Mean ES, 0.16; 95% CI, 0.08 to 0.25) but not for number of sexual partners (Mean ES, −0.01; 95% CI, −0.16 to 0.14). Interventions were more successful at increasing condom use if the sample included (1) fewer men who have sex with men (MSM) or (2) younger samples, and (3) when interventions included motivational and skills components.

Conclusions—Behavioral interventions reduced sexual risk especially if they included motivational and skills components. Such interventions have been less effective for older samples, suggesting the need for further refinement to enhance their efficacy. Motivation- and skill-based interventions have not yet been tested with HIV+ MSM who, in general, appear to have benefited less from extant risk-reduction interventions.

Persons infected with HIV can now routinely expect to live longer and healthier lives. With extended longevity, however, comes the challenge of adopting safer sexual practices for many years. Evidence suggests that the majority of HIV+ persons do reduce their risk behavior once they learn that they are infected with HIV. For example, Weinhardt *et al.*'s [1] meta-analysis of HIV counseling and testing (HIV-CT) programs found that, subsequent to HIV-CT, HIV+ participants increased condom use and reduced unprotected intercourse more so than did HIV − and untested participants. A more recent survey of 3,723 HIV+ people yielded similarly encouraging results. In this study, nearly 85% of HIV+ persons reported that they did not engage in risk behavior with uninfected partners [2].

Although many HIV+ persons reduce risk behaviors subsequent to learning that they are infected, a minority of HIV+ persons find this challenge difficult. For example, in Weinhardt *et al.*'s [2] sample, 13–19% reported unprotected vaginal or anal intercourse with partners whose serostatus was negative or unknown. Moreover, 18% of injection drug users reported that they had shared injection needles with other partners. Other studies have reported that a minority of HIV+ persons continue risky sexual or drug-use practices [3]; other individuals who are not aware of their serostatus may also transmit the disease. The route of sexual transmission is the leading cause of the approximately 40,000 new infections in the US annually [4–5]; sexual intercourse between men and women results in most HIV-1 infections acquired by adults in sub-Saharan Africa [6]. Transmission through blood transfusions, injections with infected needles, and scarification are thought to represent only a few infections [6].

Risk reduction programs for HIV+ persons have been implemented in many settings but an evaluation of these programs provides mixed evidence of their efficacy. For example, Cleary *et al.* [7] tested an informational and supportive intervention promoting behavior change with 271 HIV+ blood donors, and reported no advantage of the risk-reduction intervention relative to a control condition. In contrast, Rotheram-Borus *et al.* [8] evaluated a multi-session group-based intervention for HIV+ adolescents, and reported that adolescents reduced the number of HIV– sexual partners by 50% and decreased the number of unprotected sexual acts by 82%.

The current meta-analysis integrates the available evidence to determine the degree to which prevention programs for HIV+ persons are efficacious. We located controlled intervention studies that addressed risk reduction in samples that included HIV+ persons, and obtained effect size estimates of intervention efficacy. Our primary goal was to determine whether sexual risk reduction programs for HIV+ persons help participants to reduce the number of sexual partners and/or increase condom use. Our secondary goals were to identify moderators of intervention efficacy including intervention components that warrant inclusion and sub-groups for whom the interventions need further tailoring. Such information can provide needed guidance for intervention development and prevention research.

Method

Sample of Studies

We searched for randomized controlled trials (RCTs) using three strategies. First, we searched electronic reference databases (MEDLINE, PsycINFO, AIDSLINE, CINAHL, Dissertation Abstracts Online, and ERIC) using search terms related to HIV interventions (e.g., risk reduction, prevention, seropositive) and sexual risk behavior (e.g., condom use). Second, we used the same search terms in Internet search engines such as Yahoo! and Google through May 26, 2004. Third, we checked HIV-related listservs and the NIH database of grant awardees (CRISP), and we sent requests for papers to individual researchers conducting interventions with seropositive individuals. The goal of these supplemental strategies was to ensure the comprehensiveness of the reference database searches. Studies that fulfilled the search criteria and that were available as of November 30, 2004 were included. In some cases, information about study interventions was taken from several publications or unpublished reports. All studies included in our final sample were published reports, although we considered both published and unpublished studies in our search.

Selection Criteria

Studies or portions of studies were included if they (1) examined a deliberate HIV-risk-reduction strategy in a sample that included HIV+ participants; (2) used a RCT design; (3) measured condom use or number of sexual partners; and (4) provided sufficient information to calculate effect size (ES) estimates. Studies were excluded if the intervention(s) focused on

perinatal transmission contexts [e.g., 9], or if they utilized time-series-only designs [e.g., 10]. Consistent with these criteria, studies that did not clearly focus on sexual risk reduction were excluded. Several RCTs with HIV+ participants were excluded because (1) they did not provide critical statistics necessary for the meta-analysis [e.g., 11], (2) did not have a sexual risk-reduction component [e.g., 12], or (3) did not report either condom use or number of partners [e.g., 13]. For example, although Coates *et al.*'s [12] study targeted HIV+ individuals, its intervention focused on coping and made no mention of sexual risk reduction. Studies with samples of both HIV- and HIV+ participants were included if the number of HIV+ participants was ≥ 30 , and if separate effect sizes could be calculated for the HIV+ participants. If the latter were not available in the original reports, we contacted the authors of such studies and requested separate analyses for the HIV+ individuals in their sample; these requests resulted in 4 reports that are not typically known to the literature on prevention with seropositives [14,15,16,17]. Inclusion of these results allowed comparison of whether efforts focused on HIV+ individuals differ from those focused on HIV- or individuals whose HIV serostatus was unknown.

These criteria yielded 15 studies (Figure 1). At the first follow-up, 3,234 HIV+ individuals participated, reflecting a retention rate of 79%. Consistent with meta-analytic convention [1, 18], each intervention was treated as an individual study during analysis.

Study Information

Two raters independently coded the qualitative content of each study to describe the studies and to determine whether variation in ESs could be attributed to features of the studies. Studies were coded for the following dimensions: (1) sample characteristics (e.g., ethnicity, gender, sexual orientation); (2) risk characteristics (e.g., sex trade, drug use, proportion of entire sample HIV+); (3) design and measurement specifics (e.g., number of follow-ups); and (4) content of control and intervention condition(s) (e.g., number of sessions, training of session leaders). Because leading HIV prevention experts recognize the importance of developing interventions based upon a conceptual model of risk behavior [19], we also coded studies according to whether they provide information, motivational, and/or behavioral skills components, consistent with a leading meta-theoretical model of HIV risk behavior [20].

Across all study- and intervention-level categorical dimensions, coders agreed on 60% to 100% of judgments. Categorical variables that were used only for descriptive purposes and were not included as explanatory variables in the current analyses were coded with 90% agreement and average values of .84. Variables that served as explanatory variables were coded with 93% agreement and average values of .86. Disagreements were resolved through discussion.

Effect Size Derivation

We calculated individual ESs for relevant measures reported in each of the separate interventions. Specifically, we analyzed two self-reported sexual outcomes: (1) condom use (unspecified, vaginal, anal, oral) and (2) number of partners. Because results from the four condom use varieties converged closely, effect sizes were averaged for the primary analyses of condom use. For the purpose of this review, condom use was inferred from any measure that implied it (e.g., unprotected acts). Studies nearly always defined outcomes in continuous rather than dichotomous terms; thus, the ES calculated was the standardized mean difference (d). The pooled standard deviation served as the denominator in the ES calculation; in a small minority of cases, the denominator was instead another form of standard deviation (e.g., the standard deviation of the paired comparisons) because the pooled standard deviation was unavailable and could not be calculated from the report. The sign of each effect size was set so that it was positive when the outcome favored risk reduction, and effect sizes were corrected for sample size bias [21]. One report offered intervention statistics separately by gender [7] and another by ethnicity [14]; these reports were treated as individual studies. Effect sizes were

calculated on the measures provided at the first follow-up after the intervention to reduce method variance across studies. We averaged the ESs from multiple measures of the same outcome. When two or more intervals were assessed (e.g., condom use in previous week vs. previous 30 days), we used the interval that most closely matched the time since the intervention ended.

When a study offered a statistic controlling for baseline differences as well as statistics that were not adjusted, we used the former. If a statistic did not control for baseline, but baseline results were reported, Becker's [22] strategy was employed wherein the post-intervention effect size is corrected for any differences between the groups at baseline. When studies reported odds ratios, we transformed them to d using the Cox transformation according to published guidelines [23]. Analyses were performed with both fixed- and random-effects assumptions; analyses to examine whether features of the studies explained variability in the effect sizes followed fixed-effects assumptions [24,25]. Because at least one of the fixed-effects models of study features fully explained the variability in condom use effects, there was no need to incorporate random-effects assumptions in analyses.

RESULTS

Description of Studies

Table 1 lists the studies and their main descriptive features. The studies in the sample appeared between 1993 and 2004. Eleven studies included only HIV+ individuals, whereas the remaining four studies focused on HIV- individuals but also included some participants who were HIV+ ($M = 14\%$ HIV+ participants in these 4 samples). The samples in the studies were primarily male (64%), of African-American background (49%), and averaged 35 years of age. Fourteen (93%) studies were conducted in the U.S., with the majority (73%) of these conducted in medium to large cities. Only two studies sampled those who were known to engage in sex trading or commercial sex work (13%), and only one sampled HIV+ persons who were in drug treatment (7%); no study included incarcerated samples. More frequently, studies sampled populations of men who have sex with men (MSM; 7 [47%]) and those who use recreational drugs (12 [80%]).

All studies used random assignment, with the majority of studies (14 [93%]) assigning individuals (rather than intact groups) to conditions. All studies utilized a pre- and post-test design and included an average of 2.3 ($SD=1.0$) follow-ups after baseline data collection. The initial follow-up, the focus of this synthesis, occurred at a mean of 19.00 weeks post-intervention (range, 0–47 weeks). Only 3 studies took measures immediately and their interventions averaged 22.00 weeks in length; the other studies interventions lasted an average of 11.71 weeks. All interventions provided participants with HIV/AIDS information. Thirteen (65%) provided motivational components (e.g., social support) and 12 (60%) provided behavioral skills training (e.g., for condom use). Interventions included an average of five participants who met for six sessions that averaged 98 minutes each. Controls were typically either a wait-list (8 [40%]) or an HIV/AIDS education comparison (7 [35%]). The latter were typically matched for time/contact, or provided an abbreviated form of the intervention condition; they averaged five participants who met for three sessions of 77 minutes each.

Efficacy of the Interventions

Fourteen interventions were evaluated using only condom use, two interventions were evaluated using only number of partners, and five used both measures. ESs for studies that reported both measures were highly correlated, $r = .73$. Across the 19 interventions that assessed condom use, interventions increased condom use relative to controls, $d = 0.16$, 95% CI 0.08 to 0.25. (Table 2) Yet, the effect sizes varied widely around this mean value, $Q(18) =$

55.15, $P < .001$. (Figure 2) An examination of intervention efficacy within the four types of condom measures (unspecified, vaginal, anal, oral) confirmed this overall result. These results were parallel using either fixed- or random-effects assumptions. Seven of the studies assessed the number of sexual partners; relative to controls, intervention participants did not decrease the numbers of partners they reported, $d. = -0.01$, 95% CI -0.16 to 0.14 . These effects were homogeneous, $Q(6) = 4.62$, $P = .593$. (Table 2)

Intervention Features Associated with Increased Condom Use

Analyses revealed three features of the studies that were associated with smaller or larger efficacy as gauged by condom use. (Table 3) First, interventions were more successful to the extent that they sampled younger rather than older participants. Second, interventions were more successful to the extent that MSM were not included in the sample; interventions with 100% MSM exhibited no significant change. Third, interventions that included both motivational and behavioral-skills components increased condom use compared to interventions that had only one of these components, or neither; unless informational, motivational and behavioral-skills components were included, interventions had no effect on condom use. Model fit was excellent in the case of the MSM model, $Q_{Residual}(10) = 12.16$, $P = .27$, but not as good for each of the age and intervention content models. It is worth noting that no study focused on MSM included all three of the content components.

We also evaluated whether studies that focused exclusively on HIV+ individuals achieved greater efficacy than those that included individuals with unknown or negative serostatus; a comparison revealed no significant difference. Similarly, none of the patterns reported in the preceding paragraph changed when analyses were restricted to samples of 100% HIV+ individuals. The following features of the studies also did not relate to the magnitude of condom use effect sizes: Gender, racial composition, knowledge of serostatus, year of study, whether or not the intervention was designed to focus on HIV+ individuals, whether or not participants were injection drug users, attrition, and time since intervention ended.

Given the relatively brief follow-up intervals, and because decay of intervention effects is an important consideration, an exploratory analysis included the additional condom use follow-up measurements that five studies provided. This analysis, also, showed no change in intervention efficacy over time, $\beta = 0.01$, $P = .93$. Finally, there was no tendency for the effect sizes to depend on the year in which the interventions were conducted, $\beta = -0.20$, $P = 0.14$.

COMMENT

This quantitative synthesis is the first to evaluate the extent to which interventions evaluated with people who live with HIV work to reduce sexual risk behavior. This review focused on condom use and number of partners because these measures are the most commonly used markers of risk behavior. We synthesized RCTs because such studies provide the strongest evidence regarding the efficacy of HIV risk reduction programs. In total, 15 RCTs and 21 interventions qualified for the review (Table 1). Results showed that, with only some exceptions noted below, interventions led to reduced sexual risk behavior in people living with HIV as gauged by condom use (Table 2), an effect that showed no tendency to decay with time. The magnitude of this risk-reduction effect is equivalent to, or larger than, the effects reported in earlier meta-analyses of HIV prevention trials conducted in HIV- samples [e.g., 1,18,26–29], whose mean ESs range from 0.06 [26] to 0.25 [27].

The tendency for interventions to increase condom use relative to controls depended to some extent on the features of the studies. Compared to controls, intervention group members' condom use improved to the extent that samples had fewer men who have sex with men (MSM), in younger samples, and, importantly, when interventions included motivational and behavioral

skills enhancements. (Table 3) Although the overall magnitude of the effect size on condom use was small, it was larger for interventions that included both motivational and skills enhancements, for those that had younger participants, and for those that did not sample MSM. Indeed, interventions had no significant impact when the sample was 100% MSM or when motivational and behavioral skills enhancements were omitted.

Only seven studies examined number of sexual partners as an outcome; these studies revealed no change for intervention members compared to control-group members. This non-significant effect may reflect a restricted range (i.e., floor effect) on this variable, reliance on condom use as a risk reduction strategy, lack of intervention efficacy on this variable, or other factors not addressed in our analyses. Another possibility is that those who test positive for HIV avoid HIV-negative partners and prefer HIV-positive partners, a strategy known as serosorting [2, 30,31,32]. Unfortunately, too few studies did reported partner data in a manner that would permit such an analysis. One of the studies in our sample, which addressed MSM [33] examined just this possibility and found a small but non-significant trend for intervention participants to be less likely to have unprotected intercourse with HIV-negative or HIV-unknown partners. Future research should continue address this important issue.

Previous meta-analyses suggested that individuals who test positive for HIV increase risk-reduction behaviors such as condom use [1], especially with serodiscordant partners. As mentioned, interventions for MSM did not reduce sexual risk behavior compared to control interventions, a pattern that differs from that reached in an earlier meta-analysis of studies with (presumably) HIV- MSM [28]. Yet, because no study in the current meta-analysis provided motivational and behavioral skills to a sample of MSM, it is not presently possible to know whether such enhancements would prove successful with HIV+ MSM. However, a recent review of the literature examining risk reduction for all MSM [45] strongly suggested that notable risk reduction can occur, especially when interventions have a strong skills training component, a conclusion that converges with the current review's findings. Yet their review did not examine the extent to which serostatus related to success. Future research should test whether more comprehensive risk reduction programs (i.e., those with information, skills, and motivational components) results in improved results with HIV+ MSM.

The age-related results in the current analyses were unexpected. Contrary to earlier reviews, we found that interventions were more efficacious with younger rather than older samples [e.g., 18,27-29]. We hypothesize that older samples may have had longer duration partnerships than for those in the younger samples, a factor that is known to increase resistance to change [42]. Future studies should investigate this hypothesis.

Perhaps the most surprising finding of this work is that more than two decades into the epidemic, there have been so few intervention RCTs that focus on people living with HIV. We located only 15 RCTs that addressed the needs of this important population. Of this number, only 11 were explicitly designed to address risk behaviors in HIV+ persons. Although there have been literally hundreds of studies conducted with uninfected populations, the relative paucity of controlled studies with infected persons indicates the urgent need for research in this area, as recognized by the Centers for Disease Control and Prevention when they launched "Advancing HIV Prevention: New Strategies for a Changing Epidemic." This initiative encourages early diagnosis of HIV, increased access for HIV care, and most important to the current context, strengthened HIV prevention services for HIV+ persons.

Acknowledgements

Funding: This research was supported by National Institutes of Health grants R01-MH58563 to Blair T. Johnson and K02-MH01582 to Michael P. Carey.

We thank the study authors who made additional data and analyses available for this investigation. For comments on an earlier draft of this paper, we thank Peter A. Vanable. *Key words:* HIV prevention; Public Health; Evaluation; Meta-Analysis; Research Synthesis; Sexual Transmission of HIV; Behavioral Intervention; Secondary Prevention

References

1. Weinhardt LS, Carey MP, Johnson BT, et al. Effects of HIV counseling and testing on sexual risk behavior: a meta-analytic review of published research, 1985–1997. *Am J Public Health* 1999;89:1397–1405. [PubMed: 10474559]
2. Weinhardt LS, Kelly JA, Brondino MJ, et al. HIV transmission risk behavior among men and women living with HIV in 4 cities in the United States. *J Acquir Immune Defic Syndr* 2004;36:1057–1066. [PubMed: 15247559]
3. Centers for Disease Control and Prevention. Advancing HIV prevention: new strategies for a changing epidemic—United States, 2003. *MMWR* 2003;52:329–332. [PubMed: 12733863]
4. Centers for Disease Control and Prevention. HIV and AIDS – United States, 1981–2001. *MMWR* 2001;50:430–434. [PubMed: 11475378]
5. Centers for Disease Control and Prevention. Cases of HIV infection and AIDS in the United States, 2002. *HIV/AIDS Surveillance Report* 2002;14:1–48.
6. Buve A, Bishikwabo-Nsarhaza K, Mutangadura G. The spread and effect of HIV-1 infection in sub-Saharan Africa. *Lancet* 2002;359:2011–2017. [PubMed: 12076570]
7. Cleary PD, Van Devanter N, Steilen M, et al. A randomized trial of an education and support program for HIV-infected individuals. *AIDS* 1995;9:1271–1278. [PubMed: 8561981]
8. Rotheram-Borus MJ, Lee MB, Murphy DA, et al. Efficacy of a preventive intervention for youths living with HIV. *Am J Public Health* 2001;91:400–405. [PubMed: 11236404]
9. Heyward WL, Batter VL, Malulu M, et al. Impact of HIV counseling and testing among child-bearing women in Kinshasa, Zaire. *AIDS* 1993;7:1633–1637. [PubMed: 8286073]
10. Dancy BL, Marcantonio R, Norr K. The long-term effectiveness of an HIV prevention intervention for low-income African American women. *AIDS Educ Prev* 2000;12:113–125. [PubMed: 10833037]
11. Patterson TL, Shaw WS, Semple SJ. Reducing the sexual risk behaviors of HIV+ individuals: Outcome of a randomized controlled trial. *Ann Behav Med* 2003;25:137–145. [PubMed: 12704016]
12. Coates TJ, McKusick L, Kuno R, et al. Stress reduction training changed number of sexual partners but not immune function in men with HIV. *Am J Public Health* 1989;79:885–887. [PubMed: 2735479]
13. Wyatt GE, Longshore D, Chin D, et al. The efficacy of an integrated risk reduction intervention for HIV-positive women with child sexual abuse histories. *AIDS Behav* 2004;8:453–462. [PubMed: 15690118]
14. Dushay RA, Singer M, Weeks MR, et al. Lowering HIV risk among ethnic minority drug users: comparing culturally targeted intervention to a standard intervention. *Am J Drug Alcohol Abuse* 2001;27:501–524. [PubMed: 11506265]
15. El-Bassel N, Witte SS, Gilbert L, et al. The efficacy of a relationship-based HIV/STD prevention program for heterosexual couples. *Am J Pub Health* 2003;93:963–969. [PubMed: 12773363]
16. The National Institute of Mental Health Multisite HIV Prevention Trial Group. The NIMH multisite HIV prevention trial: reducing HIV sexual risk behavior. *Science* 1998;280:1889–1894. [PubMed: 9632382]
17. Roffman RA, Picciano JF, Ryan R, et al. HIV-prevention group counseling delivered by telephone: an efficacy trial with gay and bisexual men. *AIDS Behav* 1997;1:137–154.
18. Johnson BT, Carey MP, Marsh KL, et al. Interventions to reduce sexual risk for the human immunodeficiency virus in adolescents, 1985–2000. *Arch Pediatr Adolesc Med* 2003;157:381–388. [PubMed: 12695235]
19. Fishbein M. The role of theory in HIV prevention. *AIDS Care* 2000;12:273–278. [PubMed: 10928203]
20. Fisher, JD.; Fisher, WA. Theoretical approaches to individual-level change in HIV risk behavior. In: DiClemente, R.J.; Peterson, J.L., editors. *Handbook of HIV Prevention*. Dordrecht, Netherlands: Kluwer Academic Publishers; 2000. p. 3-55.

21. Hedges, LV.; Olkin, I. Statistical methods for meta-analysis. Orlando: Academic Press; 1995.
22. Becker BJ. Synthesizing standardized mean-change measures. *Brit J Math Stat Psychol* 1998;41:257–278.
23. Sánchez-Meca J, Marín-Martínez F, Chacón-Moscoso S. Effect-size indices for dichotomized outcomes in meta-analysis. *Psychol Methods* 2003;8:448–467. [PubMed: 14664682]
24. Johnson, BT.; Eagly, AH. Quantitative synthesis of social psychological research. In: Reis, HT.; Judd, CM., editors. *Handbook of Research Methods in Social and Personality Psychology*. New York: Cambridge University Press; 2000. p. 496-528.
25. Lipsey, MW.; Wilson, DB. *Practical Meta-Analysis*. Thousand Oaks: Sage Publications; 2001.
26. Albarracín D, McNatt PS, Findley-Klein C, et al. Persuasive communications to change actions: an analysis of behavioral and cognitive impact in HIV prevention. *Health Psychol* 2003;22:166–177. [PubMed: 12683737]
27. Kalichman SC, Carey MP, Johnson BT. Prevention of sexually transmitted HIV infection: a meta-analytic review of the behavioral outcome literature. *Ann Behav Med* 1996;21:84–92.
28. Johnson WD, Hedges LV, Ramirez G, et al. HIV prevention research for men who have sex with men: a systematic review and meta-analysis. *J Acquir Immune Defic Syndr* 2002;30:S118–130. [PubMed: 12107365]
29. Mullen PD, Ramirez G, Strouse D, et al. Meta-analysis of the effects of behavioral HIV prevention interventions on the sexual risk behavior of sexually experienced adolescents in controlled studies in the United States. *J Acquir Immune Defic Syndr* 2002;30:S94–S105. [PubMed: 12107363]
30. Wolitski RJ, Parsons JT, Gómez CA. Prevention with HIV-seropositive men who have sex with men: Lessons from the Seropositive Urban Men's Study (SUMS) and the Seropositive Urban Men's Intervention Trial (SUMIT). *J Acquir Immune Defic Syndr* 2004;37:S101–S109. [PubMed: 15385906]
31. Parsons, JT. HIV-positive gay and bisexual men. In: Kalichman, SC., editor. *Positive prevention: Reducing HIV transmission among people living with HIV/AIDS*. New York: Kluwer Academic/Plenum; 2005. p. 99-133.
32. Weinhardt, LS. HIV diagnosis and risk behavior. In: Kalichman, SC., editor. *Positive prevention: Reducing HIV transmission among people living with HIV/AIDS*. New York: Kluwer Academic/Plenum; 2005. p. 29-63.
33. Herbst JH, Sherba RH, Crepaz N, et al. A meta-analytic review of HIV behavioral interventions for reducing sexual risk behavior of men who have sex with men. *J Acquir Immune Defic Syndr* 2005;39:228–241. [PubMed: 15905741]
34. Lightfoot M, Song J, Rotheram-Borus MJ, et al. The influence of partner type and risk status on the sexual behavior of young men who have sex with men living with HIV/AIDS. *J Acquir Immune Defic Syndr* 2005;38:61–68. [PubMed: 15608527]
35. Witte, SS. HIV prevention with urban, minority couples: promoting the female condom. Unpublished dissertation
36. Fogarty LA, Heilig CM, Armstrong K, et al. Long-term effectiveness of a peer-based intervention to promote condom and contraceptive use among HIV-positive and at-risk women. *Public Health Rep* 2001;116:103–119. [PubMed: 11889279]
37. Gielen AC, Fogarty LA, Armstrong K, et al. Promoting condom use with main partners: a behavioral intervention trial for women. *AIDS Behav* 2001;5:193–204.
38. Kalichman SC, Rompa D, Cage M, et al. Effectiveness of an intervention to reduce HIV transmission risks in HIV-positive people. *Am J Prev Med* 2001;21:84–92. [PubMed: 11457627]
39. Kelly JA, Murphy DA, Bahr GR, et al. Outcome of cognitive-behavioral and support group brief therapies for depressed, HIV-infected persons. *Am J Psychiatry* 1993;150:1679–1686. [PubMed: 8214177]
40. Macneil JM, Mberesero F, Kilonzo G. Is care and support associated with preventive behavior among people with HIV? *AIDS Care* 1999;11:537–546. [PubMed: 10755029]
41. Margolin A, Avants SK, Warburton LA, et al. A randomized clinical of a manual-guided risk reduction intervention for HIV-positive drug users. *Health Psychol* 2003;22:223–228.

42. Richardson JL, Milam J, McCutchan A, et al. Effect of brief safer-sex counseling by medical providers to HIV-1 seropositive patients: a multi-clinic assessment. *AIDS* 2004;18:1179–1186. [PubMed: 15166533]
43. Rotheram-Borus MJ, Swendeman D, Comulada WS, et al. Prevention for substance-using HIV-positive young people: telephone and in-person delivery. *J Acquir Immune Defic Syndr* 2004;37:S68–S77. [PubMed: 15385902]
44. Wingood GM, DiClemente RJ, Mikhail I, et al. A randomized controlled trial to reduce HIV transmission risk behaviors and STDs among Women living with HIV: the WiLLOW Program. *J Acquir Immune Defic Syndr* 2004;37:S58–S67. [PubMed: 15385901]
45. Wolitski RJ, Gómez CA, Parsons JT. Effects of a peer-led behavioral intervention to reduce HIV transmission and promote serostatus disclosure among HIV-seropositive gay and bisexual men. *AIDS* 2005;19:S99–S109. [PubMed: 15838199]

Figure 1.
Selection process of study inclusion in the meta-analysis

Figure 2.

Forest plots of effect sizes for interventions that assessed condom use (top, A) or number of partners (bottom, B). The delta symbol for each effect size is sized proportionally to its weight in the analyses. The confidence interval for each effect size is indicated by its line and for the mean by the width of its diamond. Zero values indicate exactly no difference between the two groups. Effect sizes that significantly favor the intervention appear in green (right side); those that favor the control group appear in red (left side).

Table 1

Descriptive Features of Studies in Sample.

Study	Sample	Setting & Location	Intervention(s), Focus, Duration	Time to Follow-up	Outcomes Measured
Cleary <i>et al.</i> [7]	271 Blood donors 100% HIV+	Blood center, New York, NY	Structured Intervention Program [†] (6 weeks) HIV/AIDS information Focus on sexual risk reduction	46 weeks	Proportion sexually active Condom use for vaginal sex, vaginal sex during menses, anal receptive and active sex, and oral sex Oral-anal sex
Dushay <i>et al.</i> [14]	1300 Drug users* 17% HIV+ 73% Men 50% Black 50% Hispanic	Community outreach centers; Hartford, CT	African-American Culturally Competent Enhanced Intervention (3 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction Puerto Rican Culturally Competent Enhanced Intervention (3 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction	20 weeks	Number of sexual partners and IDU sex partners
El-bassel <i>et al.</i> [15] [‡] ; White [35] [‡]	217 Heterosexual couples* 21% HIV+ 55% Black 39% Hispanic	Primary health care setting; New York, NY	Couples Relationship-Based Intervention (6 weeks) HIV/AIDS information Motivation Behavioral Skills Focus on sexual risk reduction Woman-Alone Relationship-Based Intervention (6 weeks) HIV/AIDS information Motivation Behavioral skills Focus on sexual risk reduction	13 weeks	Frequency unprotected sex acts Proportion protected sex acts Number of sexual partners
Fogarty <i>et al.</i> [36]; Gielen <i>et al.</i> [37] [‡]	322 Women 100% HIV+ 43% Involved in sex trading 6% White 91% Black 1% Hispanic	Outpatient and pediatric HIV clinics & a primary HIV care facility; Baltimore, MD	Enhanced Intervention (26 weeks) HIV/AIDS information Behavioral skills Focus on sexual risk reduction	Immediately	Condom use with main male sex partner

Study	Sample	Setting & Location	Intervention(s), Focus, Duration	Time to Follow-up	Outcomes Measured
Kalichman <i>et al.</i> [38]	328 patients seeking HIV/AIDS care 100% HIV+ 70% Men 22% White 74% Black	Study clinic: Atlanta, GA	Transmission Risk-Reduction Intervention (2.5 weeks) HIV/AIDS information Motivation Behavioral skills Focus on sexual risk reduction	13 weeks	Number of sexual partners Frequency of intercourse and unprotected sexual acts Condom use
Kelly <i>et al.</i> [39]	115 Depressed men 100% HIV+ 94% MSM 62% White 29% Black	Mental health study clinic: Milwaukee, WI	Cognitive-Behavioral Group Intervention (8 weeks) HIV/AIDS information Focus on sexual risk reduction Social Support Group Intervention (8 weeks) HIV/AIDS information Focus on sexual risk reduction	13 weeks	Instances and number of participants reporting unprotected insertive & receptive anal intercourse
MacNeil <i>et al.</i> [40]	154 newly diagnosed HIV+ individuals 100% HIV+ 66% Women 100% Black	Counseling center or homes of participants: Semi-urban Tanzania, Africa	Enhanced Care and Support Intervention (13 weeks) HIV/AIDS information Focus on sexual risk reduction	13 weeks	Disclosed serostatus to sexual partner Condom use for last sexual intercourse Sex with person other than spouse/main partner
Margolin <i>et al.</i> [41]	90 IDU patients in drug treatment 100% HIV+ 70% Men 36% White 49% Black 16% Hispanic	Inner-city methadone maintenance program: New Haven, CT	HIV + Harm Reduction Program (26 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction	Immediately	Unprotected penetrative sex
NIMH [16]	3706 STD clinic patients* 2.5% HIV+ 58% Women 74% Black 25% Hispanic	Inner city, community based clinics: New York, NY; Northern NJ; Baltimore, MD; Atlanta, GA; Milwaukee, WI; Los Angeles, Orange, San Bernardino counties, CA	Small-Group HIV Risk Reduction Program (3 weeks) HIV/AIDS information Motivation Behavioral skills Focus on sexual risk reduction	9 weeks	Frequency of unprotected vaginal or anal intercourse Proportion of condom use Consistent condom use or abstinence
Richardson <i>et al.</i> [42]	886 sexually active patients seeking HIV treatment 100% HIV+ 74% MSM	HIV clinics: CA	Gain-Framed Safer-Sex Intervention (45 weeks) HIV/AIDS information Focus on sexual risk reduction	Less than 30 weeks	Unprotected insertive or receptive anal or vaginal intercourse

Study	Sample	Setting & Location	Intervention(s), Focus, Duration	Time to Follow-up	Outcomes Measured
	86% Men 41% White 16% Black 37% Hispanic 82% HAART		Loss-Framed Safer-Sex Intervention (45 weeks) HIV/AIDS information Focus on sexual risk reduction		
Roffman <i>et al.</i> [17]	548 Gay & Bisexual Men*	Telephone intervention: U.S., Puerto Rico, and Canada	Cognitive-Behavioral Group Counseling Intervention (14 weeks) HIV/AIDS information Motivation Focus on sexual risk reduction	Immediately	Unprotected anal and oral intercourse Proportion of all anal and oral events unprotected
Rotheram-Borus <i>et al.</i> [8]	310 HIV+ youths, ages 12–24 100% HIV+ 72% Men 19% White 27% Black 37% Hispanic	Adolescent clinic care sites: Los Angeles CA; San Francisco, CA; New York, NY; Miami, FL	“Stay healthy” and “Act Safe” Intervention (26 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction	13 weeks	Number of sexual partners Unprotected sex acts No sexual risk pattern
Rotheram-Borus <i>et al.</i> [43]	151 HIV+ youths, ages 16–29 100% HIV+ 69% MSM 78% Men 23% White 26% Black 42% Hispanic 50% HAART	Community agencies: Los Angeles, CA; San Francisco, CA; New York, NY	Telephone Intervention (18 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction In-Person Intervention (18 weeks) HIV/AIDS information Motivation Behavioral skills Equal focus on sexual and drug risk reduction	47 weeks	Number of HIV+ partners Protected sex acts with all partners, HIV+ partners 100% condom use or abstinence
Wingood <i>et al.</i> [44]	366 Women 100% HIV+ 84% Black	HIV/AIDS clinics and health departments: Anniston, AL; Birmingham, AL; Montgomery, AL; Atlanta, GA	“Women Involved in Life Learning from Other Women” (WILLOW) (4 weeks) HIV/AIDS information Motivation Behavioral skills Focus on sexual risk reduction	22 weeks	Frequency of unprotected vaginal sex Proportion who have never used condoms

Study	Sample	Setting & Location	Intervention(s), Focus, Duration	Time to Follow-up	Outcomes Measured
Wolitski <i>et al.</i> [30];	730 HIV+ MSM	Community settings: New York, NY; San Francisco, CA	Seropositive Urban Men's Intervention Trial (SUMIT) (6 weeks)	13 weeks	Unprotected anal and oral intercourse with HIV- or unknown-status partners
Wolitski <i>et al.</i> [45]	100% HIV+ 51% White 23% Black 17% Latino 1% Asian		HIV/AIDS information Motivation Focus on sexual risk reduction		

Abbreviations: HAART, highly active antiretroviral therapy; IDU, injection drug user; MSM, Men who have sex with men; STD, sexually transmitted disease.

* Sample characteristics reported describe a mixed sample of both HIV-seropositive and -seronegative or -unknown participants.

† Intervention was administered to both men and women, but results were stratified into samples of men only and women only; this single intervention was treated as two separate interventions for our current analyses.

‡ Two reports of one study were available and were used to code the study features or to calculate effect sizes.

Table 2
Efficacy of interventions to promote risk reduction at studies' first follow-up assessments.

Outcome	<i>k</i> of interventions	Weighted mean <i>d</i> (and 95% confidence interval)		Homogeneity of effect sizes	
		Fixed effects	Random effects	<i>Q</i>	<i>P</i>
Condom use	15	0.16 (0.08, 0.25)	0.16 (−0.05, 0.37)	80.76	<.0001
Unspecified context	3	0.37 (0.20, 0.54)	0.45 (0.14, 0.77)	5.02	.081
Vaginal	5	0.21 (0.09, 0.33)	0.24 (0.04, 0.45)	7.97	.158
Anal	5	0.15 (0.03, 0.26)	0.16 (0.01, 0.30)	5.31	.257
Oral	19	0.16 (0.09, 0.23)	0.19 (0.05, 0.33)	55.15	<.0001
Averaged	7	−0.01 (−0.16, 0.14)	−0.01 (−0.16, 0.14)	4.62	.593
Number of sexual partners					

Table 3
Features of studies that were linked to the efficacy of interventions to increase condom use.

Study dimension	Patterns		Test statistics	
	Level	Mean <i>d</i> (95% <i>CI</i>)	<i>k</i>	<i>P</i>
Age of sample	20 years 40 years	0.53 (0.22, 0.84) 0.11 (0.02, 0.20)	14	$\beta = -0.35$.018
Proportion of sample MSM	All MSM No MSM	0.04 (-0.07, 0.15) 0.42 (0.24, 0.60)	12	$\beta = -0.66$.002
Inclusion of intervention elements	Only information Information plus either motivational or behavioral skills All three dimensions	0.05 (-0.07, 0.16) 0.12 (-0.02, 0.26) 0.32 (0.20, 0.43)	7 3 9	$Q_B(2) = 10.98$.004

Note. Analyses are based on fixed-effects assumptions; each effect size (*d*) was weighted by the inverse of its variance. Analyses were not undertaken on the sexual frequency outcomes. MSM = Men who have sex with men. *CI* = confidence interval. *k* = number of studies. *P* = probability. Q_B = Homogeneity between categories; when statistically significant, implies differences among the means for the observed levels.