Published in final edited form as:

J Org Chem. 2004 April 2; 69(7): 2569–2572.

Unified Synthesis of C₁₉–C₂₆ Subunits of Amphidinolides B₁, B₂, and B₃ by Exploiting Unexpected Stereochemical Differences in Crimmins' and Evans' Aldol Reactions

Wei Zhang, Rich G. Carter^{*}, and Alexandre F. T. Yokochi[†]

Department of Chemistry, 153 Gilbert Hall, Oregon State University, Corvallis, Oregon 97331

Abstract

The efficient synthesis of the C₁₉–C₂₆ subunit of amphidinolide B₁ and B₂ has been completed using a boron-mediated aldol reaction. The synthesis of the C₁₉–C₂₆ subunit of amphidinolide B₃ has also been accomplished through an unexpected anti aldol reaction using a titanium-mediated process. In addition, the first reported examples of a stereochemical discrepancy between the Evans' boron-mediated oxazolidinone and the Crimmins' titanium-mediated oxazolidinethione aldol reactions are disclosed. A working hypothesis is put forth to explain the results.

The synthetic utility of chiral oxazolidinones has been well-documented in the organic community.¹ This powerful removable auxiliary has been shown to be effective in the construction of a wide variety of carbon–carbon and carbon–heteroatom bonds in a highly stereoselective fashion. Numerous working models have been put forth to explain and predict the resultant stereochemical outcome from these reactions. These models have proven to be highly reliable and general, with only isolated examples of anomalies having been reported.²

In particular, the so-called “Evans-syn” aldol reactions with chiral oxazolidinones using dibutylboron triflate and the appropriate amine base have become the standard by which new asymmetric and diastereoselective reactions are judged against (Scheme 1).³ The recently developed titanium-mediated oxazolidinethione aldol reaction, from the Crimmins laboratory, has been shown to provide comparable levels of selectivity on a wide range of systems.⁴ There are several advantages to the Crimmins' aldol methodology (e.g., relative ease of auxiliary cleavage and use of the logistically easier titanium enolates). One particularly attractive attribute is the flexibility imparted by Crimmins' chelated and nonchelated models which is dependent on the amount of TiCl₄ and amine base (normally (–)-sparteine) that is added. This modification provides access to both the Evans-syn product via the nonchelated model and the non-Evans-syn adduct via the chelated model (Scheme 1). The level of syn/anti selectivity is high (normally > 20:1) and comparable to traditional boron-mediated aldol reactions for both the chelated and nonchelated titanium-mediated conditions. It should be pointed out that Crimmins has shown that the chirality of (–)-sparteine does *not* influence the stereochemical outcome of the transformation.⁴

We were attracted to the application of the Crimmins and/or Evans methodologies for the construction the eastern subunit **15** of the cytotoxic macrolide amphidinolide B₁ (**11**)^{5,6}

rich.carter@oregonstate.edu.

[†]Director of X-ray Crystallographic Facility, Department of Chemistry, Oregon State University, Corvallis, OR 97331. E-mail: alexandre.yokochi@oregonstate.edu.

Supporting Information Available: Crystallographic data for aldol adduct **28** and experimental procedures, including copies of spectral data (¹H and ¹³C NMR), for compounds **15**, **16**, **21**, **23–25**, **28**, **29**, **31–34**, **36–38**, **44–46**, and **49**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(Scheme 2). Amphidinolide B₁ has attracted considerable synthetic interest,⁷ yet the total synthesis of **11** remains an elusive target.⁸ Two additional members of the amphidinolide B family, B₂ (**12**) and B₃ (**13**), have also been reported. These structures only differ from the parent B₁ structure by alternate stereochemistries at C₁₈ and/or C₂₂. Compounds **11** and **12** should be accessible from a common subunit **15** while C₂₂ epimer **13** should be accessible from the *anti,anti* adduct **16**.

While the application of the titanium-mediated aldol methodology has begun to appear,⁹ several important combinations have yet to be fully explored. One such example is the coupling of an *O*-benzyl-protected glycolate such as **2** or **8** with an α -chiral aldehyde **3** to provide a *syn,syn* coupled adduct such as **7** or **10** (Scheme 1). The stereochemical “Felkin” relationship of the C₂₂ and C₂₃ positions (amphidinolide B₁ numbering) would appear to be ideally suited for this transformation as both the auxiliary and the aldehyde appear to be directing the outcome in a complementary fashion. Despite this combination, the examples of a *syn,syn* adduct from an *O*-benzyl-protected glycolate such as **1**, **2**, or **8** are surprisingly rare.¹⁰ In fact, no examples of the aldol reaction depicted with oxazolidinethione **2** or **8** have been reported with α -chiral aldehydes. In this paper, we disclose the first reported examples of these combinations and the resulting synthesis of the C₁₉–C₂₆ subunits **15** and **16** for all amphidinolides B₁–B₃.

Construction of the necessary aldehyde precursor **21** was accomplished in four steps from commercially available Myers auxiliary **18**. The known alkylation¹¹ with the commercially available (*R*)-propylene oxide provided the C_{23,24}-coupled material **19** in 94:6 dr.¹² Subsequent TES protection followed by reduction with BH₃·NH₃/LDA and Ley oxidation¹³ yielded the desired aldehyde **21** (Scheme 3).

Exploration into the aldol reaction commenced with the known oxazolidinethione auxiliary **22**¹⁴ (Scheme 4). Treatment of the prescribed conditions for obtaining nonchelation or “Evans-*syn*” aldol adducts [TiCl₄ (1.0 equiv), (–)-sparteine (2.5 equiv)] provided two diastereomeric aldol adducts **23** and **24** in a 1.5:1 ratio. Unlike as predicted in the Crimmins’ models for this transformation, *none* of the expected Evans-*syn* adduct was observed. Instead, the *anti* adducts **23** (H₂₁–H₂₂ *J* = 9.3 Hz) and **24** (H₂₁–H₂₂ *J* = 8.4 Hz) were isolated in nearly equal amounts.¹⁵ The addition of additives such as NMP or alternate bases (e.g., TMEDA) did not affect the observed stereochemical outcome.⁴ Thwarted by this unexpected result, we turned to an achiral aldehyde (2-butenal) to ensure the protocol was performing as expected. Aldol reaction under the identical conditions [TiCl₄ (1.0 equiv), (–)-sparteine (2.5 equiv)] provided solely the expected Evans-*syn* adduct **25** (H₂₁–H₂₂ *J* = 3.3 Hz) in a 17:1 ratio. One possible explanation would be a mismatched relationship between the directing effect of the auxiliary and the inherent stereochemical preference of the aldehyde. To this end, the experiment was conducted with the achiral auxiliary **26**; however, equal amounts of the two previously observed *anti* stereochemistries (21*R*,22*R* and 21*S*,22*S*) were again the only observable products.

Given that the nonchelation approach provided none of the desired *syn* adduct (e.g., compound **7**), the complementary chelation aldol would appear to be the next logical step (Scheme 5). Using the enantiomeric oxazolidinethione auxiliary **27**, treatment under the chelation conditions [TiCl₄ (2 equiv), (–)-sparteine (1 equiv)] proceeded poorly and in low yield. Crimmins has also reported that the use of less TiCl₄ [(1 equiv), (–)-sparteine (1 equiv)] proceeds via the chelated model.⁴ Treatment using these conditions provided some improvement in the selectivity of the transformation, yielding a more respectable 4:1 ratio of the two *anti* products (**28/29**) (**28**: H₂₁–H₂₂ *J* = 8.8 Hz; **29**: H₂₁–H₂₂ *J* = 9.2 Hz) with *none* of the predicted *syn* adduct. It is important to note that despite the expected directing effect of the auxiliary (e.g., **22** should give the same absolute configuration at C₂₁, C₂₂ under nonchelation conditions as enantiomeric auxiliary **27** gives under chelation conditions), the major product **28** from the chelation-controlled conditions using **27** contained the 21*R*,22*R* stereochemistry

of the **minor** product from the nonchelation conditions with **22**. Alcohol **28** is synthetically useful as it possesses the correct *anti,anti* C_{21–23} stereochemistry for amphidinolide B₃. Interestingly, use of the enantiomeric auxiliary **22** under chelation conditions led to an equal mixture of the adducts **23** and **24**. A similar result was observed with the achiral auxiliary **26**. It became apparent at this juncture that the titanium-based aldol were unable to provide the necessary stereochemical relationship (e.g., **7** or **10**) for amphidinolides B₁ and B₂. We were gratified to observe that use of the auxiliary **30**¹⁶ under boron-mediated conditions did yield the desired Evans-*syn* adduct **31** (H₂₁–H₂₂ *J* = 2.1 Hz) in a 95:5 *syn,syn* and *syn,anti* ratio (72% isolated yield of **31**). To the best of our knowledge, these results represent the first reported examples of the stereochemical divergence between the titanium-mediated oxazolidinethiones and boron-mediated oxazolidinones.¹⁵

Stereochemical assignment of the aldol products **23–25**, **28–29**, and **31** was accomplished via a series of degradation experiments and X-ray crystallographic analysis of **28** (Scheme 6). Reductive removal of the auxiliaries from adducts **24** and **28** yielded an identical diol **32**, thereby confirming **24** vis-à-vis X-ray structure **28**. An analogous path was followed for the adducts **23** and **29** to yield the diol **34**. The stereochemistry of **31** was confirmed via conversion to the TBDPS ether **36** and correlation with the TBDPS ether **33** through TPAP oxidation to the ketone **37**. This degradation also indirectly established one of the two unknown stereocenters of **23** and **29** as 2*S* by assignment of both aldol adducts **28** and **31** as the 2*R* configuration. The 2*S* configuration was confirmed by Mosher ester analysis of **35**.¹⁷ Finally, the stereochemistry of **25** was confirmed by reduction to a known compound **38**.¹⁸

A working hypothesis for the observed stereochemical results invokes the use of the open transition state **40** and boat transition states¹⁹ **41** and **43** to explain the observed stereochemistry (Scheme 7). One possible rationale for the inability of these transformations to proceed through the chair transition states **39** and **42** could be an unfavorable interaction between the benzyloxy substituent and the α -position of the aldehyde.²⁰ As steric bulk at these positions increase, this unfavorable interaction should become more significant. We also hypothesize that the bulk of the benzyloxy substituent may be increased by an aggregation effect. While additional studies are necessary to verify this aggregation effect, we do observe a modest correlation of concentration with diastereoselectivity [0.15 M, 4:1 dr; 0.05 M, 2:1 dr (**28:29**)]. A similar correlation is observed for this transformation by even a slight variation in the ratio of auxiliary to aldehyde. The use of 1.5 equiv of auxiliary **27** with 1 equiv of the aldehyde **21** yielded a 4:1 ratio (**28:29**), while 2.0 equiv of the same auxiliary **27** (1 equiv of **21**) yielded a 2:1 ratio (**28:29**) under the described chelation conditions. Phillips and co-workers have also commented on the sensitivity of titanium-mediated aldol reactions to slight modifications.^{19a} In contrast to the titanium enolates, the boron enolates are unable to aggregate in the Zimmerman–Traxler transition state due to the full valence shell on boron. This important difference does appear to agree with the observed stereochemical results. Finally, an open transition state **40** is put forth to justify the *anti* adduct **23**.¹⁵ This proposed explanation allows for an approach of the aldehyde consistent with the Felkin model.

Completion of the C₁₉–C₂₆ subunits of amphidinolide B₁–B₃ was accomplished in three steps (Scheme 8). Silylation using TESOTf provided the bissilylated compound **44**. Conversion to the thioester using catalytic amounts of KSEt followed by cuprate coupling gave the methyl ketone **15**.²¹ An analogous path was pursued with *anti,anti* adduct **28** to provide the methyl ketone **16**. Interestingly, attempted introduction of TBS protecting group on the adduct **31** led to competitive silyl migration. This migration was not observed with the adduct **28**.

A unified strategy for the synthesis of the C₁₉–C₂₆ subunits of amphidinolide B₁–B₃ **13–15** has been accomplished. The first reported examples of the divergence of the titanium-mediated

oxazolidinethione aldol reaction to provide the anti adducts **23–24** and **28–29** as the sole products have been reported. A working model is put forth to explain the stereochemical results.

Supplemental Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Financial support was provided by the National Institutes of Health (NIH) (GM63723) and Oregon State University. This publication was also made possible in part by a grant from the NIH–National Institute of Environmental Health Sciences (P30 ES00210). We thank Professor Max Dienzer (Mass Spectrometry Facility, Environmental Health Sciences Center, Oregon State University) and Dr. Jeff Morré (Mass Spectrometry Facility, Environmental Health Sciences Center, Oregon State University) for mass spectra data, Roger Kohnert (Oregon State University) for NMR assistance, and Dr. Roger Hanselmann (Rib-X Pharmaceuticals) for his helpful discussions.

References

1. Evans, DA.; Kim, AS. Handbook of Reagents for Organic Synthesis: Reagents, Auxiliaries and Catalysts for C–C Bonds. Coates, RM.; Denmark, SE., editors. John Wiley & Sons; New York: 1999. p. 91–101.
2. (a) Clive DLJ, Yu M. Chem Commun 2002:2380–81. (b) Evans DA, Kaldor SW, Jones TK, Clardy J, Stout TJ. J Am Chem Soc 1990;112:7001–31. (c) Iseki K, Oishi S, Kobayashi Y. Tetrahedron 1996;52:71–84.
3. (a) Evans DA, Bartroli J, Shih TL. J Am Chem Soc 1981;103:2127–29. (b) Evans DA, Nelson JV, Taber TR. Top Stereochem 1982;13:1–115.
4. Crimmins MT, King BW, Tabet EA, Chaudhary K. J Org Chem 2001;66:894–902. [PubMed: 11430110]
5. Ishibashi M, Ishiyama H, Kobayashi J. Tetrahedron Lett 1994;35:8241–42.
6. (a) Kobayashi J, Ishibashi M, Nakamura H, Ohizumi Y, Hirata Y, Sasaki T, Ohta T, Nozoe S. J Nat Prod 1989;52:1036–41. [PubMed: 2607346] (b) Ishibashi M, Ohizumi Y, Hamashima M, Nakamura H, Hirata Y, Sasaki T, Kobayashi J. J Chem Soc Chem Commun 1987:1127–29.
7. (a) Cid B, Pattenden G. Tetrahedron Lett 2000;41:2573–76. (b) Ohi K, Nishiyama S. Synlett 1999:571–72. (c) Ohi K, Nishiyama S. Synlett 1999:573–75. (d) Eng HM, Myles DC. Tetrahedron Lett 1999;40:2275–78. (e) Eng HM, Myles DC. Tetrahedron Lett 1999;40:2279–82. (f) Chakraborty TK, Thippewamy D. Synlett 1999:150–52. (g) Ishiyama H, Takemura T, Tsuda M, Kobayashi J. J Chem Soc Perkin Trans 1999;1:1163–66. (h) Chakraborty TK, Thippewamy D, Suresh VR, Jayaprakash S. Chem Lett 1997:563–64. (i) Chakraborty TK, Suresh VR. Chem Lett 1997:565–66. (j) Lee DH, Lee S-W. Tetrahedron Lett 1997;38:7909–10. (k) Ohi K, Shima K, Hamada K, Saito Y, Yamada N, Ohba S, Nishiyama S. Bull Chem Soc Jpn 1998;71:2433–40.
8. For total syntheses of other members of the amphidinolides: (a) Williams DR, Kissel WS. J Am Chem Soc 1998;120:11198–99. (b) Williams DR, Myers BJ, Mi L. Org Lett 2000;2:945–48. [PubMed: 10768193] (c) Williams DR, Meyer KG. J Am Chem Soc 2001;123:765–66. [PubMed: 11456603] (d) Lam HW, Pattenden G. Angew Chem Int Ed 2002;41:508–511. (e) Maleczka RE, Terrell LR, Geng F, Ward JS III. Org Lett 2002;4:2841–44. [PubMed: 12182569] (f) Trost BM, Chrisholm JD, Wroblewski ST, Jung M. J Am Chem Soc 2002;124:12420–21. [PubMed: 12381177] (g) Fürstner A, Aïssa C, Riveiros R, Ragot J. Angew Chem Int Ed 2002;41:4763–66. (h) Ghosh AK, Liu C. J Am Chem Soc 2003;125:2374–75. [PubMed: 12603108]
9. (a) Crimmins MT, King BW, Zuercher WJ, Choy AL. J Org Chem 2000;65:8499–09. [PubMed: 11112569] (b) Crimmins MT, Choy AL. J Am Chem Soc 1999;121:5653–60. (c) Crimmins MT, Katz JD, McAtee LC, Tabet EA, Kirincich SJ. Org Lett 2001;3:949–52. [PubMed: 11263923]
10. Only four reported examples of this combination have appeared in the literature: (a) Crimmins MT, Katz JD, McAtee LC, Tabet EA, Kirincich SJ. Org Lett 2001;3:949–52. [PubMed: 11263923] (b) Chakraborty TK, Suresh VR. Tetrahedron Lett 1998;39:7775–78. (c) Piscopio AD, Minowa N, Chakraborty TK, Koide K, Bertinato P, Nicolaou KC. J Chem Soc Chem Commun 1993:617–18. (d) Jones TK, Reamer RA, Desmond R, Mills SG. J Am Chem Soc 1990;112:2998–3017.

11. Myers AG, McKinstry L. *J Org Chem* 1996;61:2428–40.
12. It should be noted that this stereochemical combination provides the epimeric stereochemistry at C₂₅ versus the target **11**; however, the alkylation of the enantiomeric (*S*)-propylene oxide proceeds in poor selectivity due to its mismatched relationship to the approaching enolate. This stereocenter will be inverted later in the synthetic sequence.
13. Ley SV, Norman J, Griffith WP, Marsden SP. *Synthesis* 1994:639–66.
14. Crimmins MT, McDougall PJ. *Org Lett* 2003;5:591–94. [PubMed: 12583777]
15. It should be noted that Crimmins has recently reported the development of a titanium-mediated oxazolidinethione method for the synthesis of *anti* aldol adducts through an open transition state; however, this reaction protocol requires the addition of an additional 2.5 equiv of TiCl₄ immediately prior to addition of the aldehyde. See ref ¹⁴.
16. Evans DA, Gage JR, Leighton JL, Kim AS. *J Org Chem* 1992;57:1961–3.
17. (a) Ohtani I, Kusumi T, Kashman Y, Kakisawa H. *J Am Chem Soc* 1991;113:4092–96. (b) Dale JA, Mosher HS. *J Am Chem Soc* 1973;95:512–19. (c) Sullivan GR, Dale JA, Mosher HS. *J Org Chem* 1973;38:2143–47.
18. Reduction using lithium borohydride revealed the known diol which match the reported ¹H NMR and ¹³C NMR spectrum. $[\alpha]_{\text{D}}^{20} = +17.5$ ($c = 0.48$, CHCl₃) vs lit. $+16.8$ ($c = 2.0$ in CHCl₃). Fuhry MAM, Holmes AB, Marshall DR. *J Chem Soc Perkin Trans* 1993;1:2743–46.
19. The preference for a boat transition state in oxazolidinethione aldol reactions has been proposed previously. (a) Guz NR, Phillips AJ. *Org Lett* 2002;4:2253–56. [PubMed: 12074680] (b) Evans DA, Downey CW, Shaw JT, Tedrow JS. *Org Lett* 2002;4:1127–30. [PubMed: 11922799]
20. For recent and somewhat related examples, see: (a) Evans DA, Siska SJ, Cee VJ. *Angew Chem Int Ed* 2003;42:1761–65. (b) Marco JA, Carda M, Díaz-Oltra S, Murga J, Falomir E, Roeper H. *J Org Chem* 2003;68:8577–82. [PubMed: 14575488]
21. White JD, Carter RG, Sundermann KF, Wartmann M. *J Am Chem Soc* 2001;123:5407–13. [PubMed: 11389619]

Scheme 1.

Generalized Examples of Crimmins' and Evans' Aldol Adducts

Scheme 2.
Retrosynthetic Analysis of Amphidinolide B₁

Scheme 3.

Key: (i) ref 11, (*R*)-propylene oxide, 92%, 94:6 dr; (ii) TESCl, Et₃N, DMAP, CH₂Cl₂; (iii) LDA, BH₃·NH₃, THF, 0 °C, 78% over two steps; (iv) TPAP, NMO, CH₂Cl₂, 85%.

Scheme 4.

Key: (i) $TiCl_4$ (1 equiv), $(-)$ -sparteine (2.5 equiv), **21**, CH_2Cl_2 , 0.15 M, $-78^\circ C$, 40 min, 1.25:1 dr (**23**:**24**), 44% **23**, 30% **24**; (ii) $TiCl_4$ (1 equiv), $(-)$ -sparteine (2.5 equiv), 2-butenal, CH_2Cl_2 , 0.15 M, $-78^\circ C$, 40 min, 80%.

Scheme 5.

Key: (i) TiCl_4 (1 equiv), (-)-sparteine (1 equiv), **21**, CH_2Cl_2 , 0.15 M, -78°C , 40 min, 4:1 dr (**28:29**), 53% **28**; (ii) Bu_2BOTf (1 equiv), Et_3N (1.1 equiv), PhMe, 0.15 M, -50 to -30°C , 2 h, 72%.

Key: (i) LiBH₄, MeOH, THF, 0 °C to rt; (ii) TBDPSCl, imid, DMAP, CH₂Cl₂, 0 °C to rt; (iii) (*R*)/(*S*) Mosher acid chloride, DMAP, CH₂Cl₂; (iv) difference in ppm [(*S*)-Mosher ester–(*R*)-Mosher ester, CDCl₃, 400 MHz NMR] shown on structure **35**; (v) TPAP, CH₂Cl₂, molecular sieves.

Scheme 7.
Possible Explanation for Observed Stereochemical Outcome

Scheme 8.

Key: (i) TESOTf, 2,6-lutidine, CH₂Cl₂, 0 °C; (ii) EtSH, KH (cat.), THF; (iii) Me₂CuLi, Et₂O, -50 °C; (iv) TESOTf, 2,6-lutidine, CH₂Cl₂, 0 °C.