Technical Section

TECHNICAL NOTES & TIPS

TECHNICAL NOTES

Retraction of oedematous eyelids with paper-clips

LEO H-H CHENG, PRAGIA KUMAR

Department of Oral and Maxillofacial Surgery, Barts and the London NHS Trust, and Homerton University Hospital NHS Trust, London, UK

CORRESPONDENCE TO

Leo H-H Cheng, Department of Oral and Maxillofacial Surgery, Barts and the London NHS Trust, 3rd Floor, Alexandra House, Royal London Hospital, Whitechapel, London E1 1BB, UK
E: leo.cheng@bartsandthelondon.nhs.uk

BACKGROUND

Examination of ocular content and visual acuity, as part of full facial examination, is often hindered by post-traumatic or post-operative peri-orbital and eyelid oedema or haematoma. In order to exclude rare complications of occult ocular injury or retrobulbar haemorrhage, visualisation of the cornea and eyeball is crucial for making the correct clinical diagnosis. Patients frequently refuse ocular examination by clinicians' gloved fingers due to blood clots sealing the eyelashes together or painful swollen eyelids.

TECHNIQUE

A pair of ordinary paper clips (length 3.2 cm or 7.4 cm) are bent to approximately 90° near the midpoint of the paper clip with gentle finger pressure to create a hand-held retractor (Fig. 1). The paper-clip retractor should be held at the proximal end of the clip in order to allow the exposed wire end to lie vertically away from

Figure 1 An eyelid retractor made from bending a 3.2-cm paper-clip.

Figure 2 Retraction of post-traumatic swollen eyelids with a pair of paper-clips.

patient's eyelid. The paper-clip retractors are then cleaned with antiseptic solution before gently retracting the swollen eyelids. The eyelid will slowly open with a stepwise technique to advance the paper-clip retractors only on the skin surface (Fig. 2).

DISCUSSION

Paper-clip eyelid retractors are simple, cheap, readily available self-made eyelid retractors, and are helpful in opening closed oedematous eyelids in both trauma and postoperative situations. No patients suffered from any undue pressure or complications from the paper-clip eyelid retractors.

A novel use for synthetic ligaments in hip surgery

B HOLROYD, DE FERN

Trauma and Orthopaedic Department, Royal Cornwall Hospital, Truro, Cornwall, UK

CORRESPONDENCE TO

B Holroyd, Trauma and Orthopaedic Department, Royal Cornwall Hospital, Truro, Cornwall, UK
E: benholroyd@hotmail.com

BACKGROUND

Synthetic ligament replacement or augmentation is an accepted part of orthopaedic practice. Their use as a dynamic augmenta-

Figure 1 Bare trochanter.

Figure 2 Ligament sutured onto abductors.

Figure 3 Ligament sutured onto short external rotators.

Figure 4 Securing ligament through intra-osseous tunnels.

tion device around the hip has not been described previously. A bare trochanter is a lesion (Fig. 1) similar in pathology to a rotator-cuff tear in the shoulder¹ and is occasionally seen during an anterolateral approach to the hip. Small defects can be repaired by suturing the abductors onto the bare area, but larger defects require a different strategy. Similar defects can be found after posterior approaches to the hip particularly at the time of revision surgery. We describe the novel use of a synthetic ligament to reconstruct these defects.

TECHNIQUE

The broad-shaped end of a 30-fibre LARS (Corin Group; Cirencester GL7 1YJ, UK) rotator-cuff ligament (1500 N) is sutured onto the remnants of the gluteus medius tendon in the case of a bare trochanter (Fig. 2) or onto the short external rotators in the case of revision posterior approaches (Fig. 3). The two limbs of the Y-shaped ligament are attached onto the proximal femur through intra-osseous tunnels secured with interference screws. The ends of the ligament are tied onto each other to reinforce the reconstruction (Fig. 4).

DISCUSSION

Dynamically reconstructing damaged muscles following hip surgery can be achieved successfully with synthetic ligaments which have inherent strength whilst acting as a scaffold into which native tissue can grow. Previously described repairs have been static tethers² with an inherent hard end-point to movement making the dynamic technique described an attractive proposition.

References

1. Bunker TD, Esler C, Leach W. Rotator-cuff tear of the hip. *J Bone Joint Surg Br* 1997; **70**: 618–20.
2. Barbosa JK, Khan AM, Andrew JG. Treatment of recurrent dislocation of total hip arthroplasty using a ligament prosthesis. *J Arthroplasty* 2004; **19**: 318–21.