and dragged into a criminal court of law, and arraigned before a jury for an act of supposed crime, over which she had not the least command or power of controlling.

Although this is a subject that belongs to the obstetric department only, it may sometimes become a medico-legal question, and is I think, well worth the notice of professors of medical jurisprudence, for in some cases of infanticide and concealment of the birth of a child it may be very properly adduced as mitigating evidence.

Should you concur with me that this, as well as the size of the hand and arm, in the practice of midwifery, are subjects that have been neglected or overlooked by teachers and obstetric practitioners, you will perhaps allow their attention to be directed to it by the insertion of my remarks in an early number of the *Journal*.

Bath, September 24, 1850.

CASES OF FEVER, WITH REMARKS.

By MICHAEL FOSTER, Esq., HUNTINGDON.

It may not be uninteresting to give a brief report of the outbreak of a fever of a fatal character at the Huntingdon Union Workhouse, in January, 1847; more particularly as a fever of the same type, with various modifications, has just now prevailed at Herringford Abbots, a village four miles north-east of Huntingdon. I will give the first case in detail, then a copy of the minutes, as reported to the Board, then a summary, with remarks.

Mary Archer was admitted into the lying-in ward about a fortnight ago, in the family-way by a man named Joseph Martin, of Great Catworth. During this period she was observed to suffer from a depression of spirits, but never complained of being in any way ill. Had been very sick on the Sunday week previous to her confinement, and afterwards occasionally, and on each occasion had brought up a darkish green-coloured fluid.

On Tuesday, the 5th of January, 1847, at one o'clock in the afternoon, the attention of the medical attendant was first drawn to her. He found her very feverish; tongue brown in the centre, with reddened tip and edges; pulse small and weak, not unusually quick; the bowels had been relieved the evening before; had been sick frequently during the day, vomiting dark coloured fluid. At six P.M. was sent for to attend her in labour. Found the head pressing on the perineum, which was very turgid, especially just anterior to the anus, where its colour was almost black; the bag of the membranes protruded, which were also of very dark colour; the head was lying a little across, the left ear being easily felt at the back of the pubes. Learnt that she had been just in this way two hours. At nine P.M. she was delivered of a dead male child, about the full period, (according to her reckoning wanting a week.) The placenta came away immediately. Uterus contracted;

no hæmorrhage; had had no strong pains at any time during the labour; she was sick every now and then afterwards; seemed restless; had constant alternating cold and hot perspirations; and complained of fainting sensations; drew her breath short, and moaned slightly occasionally. Took a cup of tea and toast about ten o'clock; in half an hour afterwards was free from sickness, and appeared more comfortable, and remained so for three quarters of an hour. From this time (eleven o'clock) she was tolerably comfortable, taking a little brandy and water occasionally to allay the sickness, which continued at intervals. Later at night she became restless, breathing short and quick; the neck, face, and lips became very blue; and having rested quietly three quarters of an hour, she raised herself and suddenly expired, at two o'clock in the morning.

Post-mortem, twelve hours after death.—Abdomen contained a large quantity of bloody serum; uterus natural; liver healthy, larger than is usual; stomach much disturbed with flatus; contained a dark coloured fluid, showing patches of inflammation here and there. There was serous fluid effused into the pericardium. Heart healthy and of natural appearance; the right ventricle and auricle were full of very dark coagulated blood; the left were also full of blood, but fluid. Lungs were healthy. The œsophagus and the cardiac orifice of the stomach were dark coloured. There was much venous congestion of the body generally, but especially in the neck. The contents of the stomach were analyzed. Rheinsch's process for arsenic, protochloride of tin, for corrosive sublimate, and other tests, were tried, but no signs of any poison were discovered.

Minutes reported to the Board of Guardians.

"Monday, January 11, 1847.

"I beg to call the attention of the Board to the sudden appearance of fever in this house of a very severe and dangerous character, partaking of the type of bilious remittent at the onset, rapidly leading on to typhoid disorganization of the tissues. The first case was that of Mary Archer, aged 25, who was taken ill, with severe vomiting of dark brownish-green coloured fluid, extreme prostration, premature delivery, and died in the course of twenty-four hours. The *post-mortem* examination showed extensive effusion of turbid fluid into the peritoneal cavity, great enlargement of the liver, and much distension of the stomach (flatus;) its internal surface very red in some parts, and of a dark slate colour in others. The right side of the heart was loaded with blood, quite black, as also were the vessels of the neck and throat. The next case, that of George Bell, aged 12, exhibited the same severe vomiting violent pains of the head and belly, and who died in three days. Since then, and up to this day, nine more cases of greater or less degree of severity have taken place. I have recommended these to be separated as much as possible from the other inmates; chloride of lime to be put into most of the rooms, and large quantities of unslacked lime in all parts of the drains; also the pregnant women to be removed from the house.

"15th.—Six more cases have appeared since Monday.

Three have died, viz., Savage, Howe, and Baram; the last was a weak epileptic patient, quite unable to withstand further illness. Three or four of the other cases are of mild character, and going on favourably; Bond, Rowell, and Selby are still very ill, and upon the whole the fever spreads, but the more recent cases are less severe in their character. No men have been attacked. I reported to Sir John Walsham, according to his request, 'that although there was some room to suspect the fever might have been brought to the house by a Dutch sailor boy (a tramp), yet the evidence was not satisfactory; but when the drainage throughout the whole of the workhouse, and the open and overflowing state of the cess-pools are examined, I have no

hesitation in saying there is quite sufficient to generate fever of the worst kind.'

"January 22nd.—Since last week there have been only two new cases of fever. Elizabeth Wright and Martha Bell, both nurses. They are going on well. There have been three deaths—viz., Ann Saunders, an old debilitated female, Ann Whitehead, a blind person, very weakly, who had suffered long from scrofula, and Lydia Jackson. All the other cases are doing well and promise recovery.—"29th. No new case of fever.

"February.—The state of the workhouse as regards fever is much better. No new cases.

"March 5th.—I think pregnant women may now be admitted."

SCHEDULE A.

Name and age of Patient.	Nature of Disease.	Duration of Disease.	Treatment and Diet.
Mary Archer, aged 26 ...	Childbirth. Bilious fever, severe vomiting, extreme prostration, short rapid respiration, venous congestion, abdomen distended with flatus, effusion into peritoneum, gastritis	Died in 18 hours after seizure	Salines
George Bell, aged 12 ...	Symptoms the same as last, gastritis two days, meningitis one day	Died in 4 days	Ant. Tart. et Mag. Sulph., &c. Hyd. c. Cretâ, Pulv. Doveri; cold lotions
Rebecca Baram, aged 50	All her life subject to epilepsy. Symptoms as the last	Died in 4 days	
Sarah Savage, aged 23	Remittent. Premature delivery after 6 days, peritonitis 3 days, delirious 2 days before death	Died in 9 days	Leeches, blisters, calomel, and wine
Eliza Howe, aged 17 ...	Confinement, followed two days after by fever. Peritonitis	Died in 3 days	Treatment as the last
Ann Whitehead, aged 23	Blind. Scrofulous ulcers, fever, and erysipelas. Delirious	Died in 8 days	Wine during two days
Ann Saunders	Debility and old age. Fever	Died in 4 days	Wine, &c.
Lydia Jackson, aged 35	Remittent. Gastritis, meningitis two days, severe	Died in 7 days	Leeches, blisters, wine, calomel, ointment, lotions
Mary Bond, aged 23 ...	Remittent fever followed by miscarriage after 1 month. Recurrence of fever, mammary abscess, extreme debility, vomiting, &c.	Ill altogether 6 wks. Died three weeks after abortion	
George Gillson ... } Frederick Eates ... }	Attacked mildly	Recovered in 3 or 4 days	Milk, wine, and medicine
Sarah Leach, middle-aged	Bad constitution, fever and severe diarrhœa	Cured in 14 days	Hyd. c. Cretâ, Pulv. Doveri, &c., as the others
Hannah Selby, aged 20	Had gonorrhœa. Attacked severely, peritonitis, delirious	Recovered in 20 days	Leeches, blisters, wine, ammonia, Hydrarg. cum Cretâ, and Dover's powder, &c.
Sarah Hall	Fever ran its course, not severe	Recovered in 14 days	
Ann Johnson	As the last.	Recovered in 14 days	
Emma Heddens		Recovered in about 14 days	
Elizabeth Wright, (Nurse)		Recovered in about 9 days	
Sarah Rowel, aged 14	Attacked severely. Peritonitis	Convalescent in 14 days	Leeches, blisters, &c.
Martha Bell (Nurse) ...	Attacked severely. Erysipelas	Convalescent in 10 days	Wine, &c.
Maria Hall (Hired nurse)	Attacked very severely. Peritonitis, erysipelas, inflammation of the glands of the neck.	Ill from 18 to 20 days	Leeches, blisters, calomel, sinapisms, Hydrarg. c. Cretâ and Dover's powder, ammonia and opium.

SCHEDULE B.—*Summary of Deaths and Recoveries.*

Date.	Name and Duration of Disease.	Nature of Disease.	Name.	The remaining nine females had the fever more or less severely.	Duration of Disease.
Jan. 6th.	M. Archer died in 18 hrs.	Confinement	Elizabeth Wright	Recovered in 9 days	
" 9th.	Geo. Bell .. 4 days	Confinement	Martha Bell	" in 10 days	
" 14th.	Eliza Howe .. 3 days	Old epileptic	Sarah Rowell ...	" in 14 days	
" 15th.	R. Baram .. 4 days	Premature delivery	Sarah Leach ...	" in 14 days	
" 15th.	S. Savage .. 9 days	Blind. Very scrofulous	Sarah Hall	" in 14 days	
" 17th.	A. Whitehead .. 8 days	Weak and old	Ann Johnson ...	" in 14 days	
" 19th.	A. Saunders .. 4 days	...	Emma Eddens ...	" in 14 days	
" 20th.	L. Jackson .. 7 days	...	Hannah Selby ...	" in 20 days	
Feb. 9th.	Mary Bond .. 1 mth.	Abortion fifth month	Maria Hall	" 18 to 20 days	

Of the eleven who recovered, two were boys who were only mildly attacked and soon recovered.

REMARKS.—*The type* may be considered bilious-remittent, great derangement of the biliary system, rapid and destructive action of the serous membranes of the brain and abdomen; the latter, particularly in pregnant women, who seemed especially susceptible, and in some cases erysipelas appeared.

Cause.—No doubt gaseous exhalations from the overflowing drains and open cess-pools in several parts of the premises. So closely did the first case resemble poisoning, that I analyzed the contents of the stomach with the view of ascertaining whether any poison had been administered, and it was whilst musing upon what could be the cause of death that a second case (G. Bell) appeared with analogous symptoms. The same causes in operation at Huntingdon. When the drains were cleansed the fever subsided DIRECTLY. Whether infectious or not, three of the nurses, previously in good health, were attacked, and the fever was confined entirely to the *female* side of the house.

Four pregnant women were seized; all died.* Of

the twenty cases (out of seventy-three females in the house) nine died.

Such was the fatality that much cannot be said for treatment. In the earliest stage the tartrate of antimony, in full doses, was the most beneficial; if not much allayed by this nothing very useful. The most satisfactory steps were the removal of all the inmates that could be moved, and the *non admission of any others.*

Huntingdon, Sept. 19, 1850

Hospital Reports.

QUEEN'S HOSPITAL, BIRMINGHAM.

REPORTS OF CASES ADMITTED UNDER DR. DAVID NELSON.

Extracted from the Clinical Records,

BY W. J. MOORE, ESQ. M.R.C.S., AND L.A.C., RESIDENT MEDICAL OFFICER.

CASE XV.

Case of Thomas Sheridan, aged 20. Temp. nervous. Affected with Phthisis, terminating in death. Admitted March 4.

HISTORY.—This patient stated that he was only conscious of being ill three months before. At Christmas he caught cold, and had since that time been troubled with cough, which had at first been moist, but had become drier under medical treatment. He had of late become pale and lost flesh.

Symptoms.—On examination, the tongue was found red at the point and edges; bowels were costive, and the discharge dark, and the urine was scanty and high coloured; the expectoration was sometimes thick and yellow, and at other times thin and tenacious; he had not sweated since he began to wear flannels; the chief dulness was observable over the right side, anteriorly; pulse 140, respiration, 36.

Diagnosis.—That there was tubercular deposit, chiefly in the right lung, and afterwards excavation and cerebral effusion.

Treatment.—He was a quiet but spirited young man, and was ready to submit to any treatment. The plan decided on was to employ derivatives, expectorants, tonics, and alteratives; and the orders were:—Hab. Hydrarg. Chlorid., gr. v., statim et Ferri. Carb., gr. iij.; Pulv. Aloes, Pulv. Ipecac. utr., gr. j., omni nocte. Ol. Aselli, dr. j., ter die; Tinct. Opii Camph., dr. j., ter die; Acid. Sulph. Aromat. gtt. xxx.; Sulph. Quinæ, gr. j., omni nocte. Ap. ves. lat. dext.

7th.—The cough was much easier, and the expectoration profuse; he had had no sweating, but his tongue was coated, and he felt his throat so enlarged as to impede his breathing and excite cough from time to time; he was also hoarse. On examination of the chest, again the dulness was observed to be most apparent under the centre of the right clavicle, where there

* One pregnant woman was removed and she escaped; the two others were attacked before measures could be taken to remove them, and it was not known that the third was pregnant.