CASES OF
HEPATITIS, DYSENTERY, AND DIARRHŒA,
WITH REMARKS.

By W. T. BLACK,

ASSISTANT-SURGEON TO THE FORCES, SOUTH AFRICA.

CASE I.

THE case of hepatitis occurred in a female, Mrs. M., of a plethoric habit of body, who had been subject to similar attacks while resident in India. The first symptoms inducing a call for medical aid, appeared on June the 11th, 1852, and were of a decided character. They reached their intensity on the 14th, notwithstanding the remedies used; the urine was scanty, thick, bilious coloured; fever high; breathing oppressed; and considerable pain in right hypochondrium. A smart dose of jalep and calomel was then given, which gave much relief, and sweating now took place, which considerably lessened the fever, and may have been caused by the depressing effect of the purgative allowing the diaphoretics previously administered to act easier upon the skin. The purge was repeated the next day, and gave further relief to the local symptoms, and the sweating still continued to keep down the fever. The mouth now showed the effects of mercury, an ointment containing which had been used to dress a blister applied to the side. The intense symptoms now abated, and gradually declined under the use of saline drinks and purgative pills, but the stomach could not bear antimonial medicines since the aggravation of the disease on the 14th, and the patient continued to feel much weakness. From some cause a relapse of the complaint began on the 25th, and the next day had reached as great intensity as ever, with the addition of dyspnoea, and sense of oppression and weight around the the region of the liver; and the bowels became again costive and the perspiration ceased. A large dose of compound jalap powder gave relief to the local symptoms, its action being kept up by a purgative and saline mixture, and perspiration again appeared to subdue the fever. No attempt was made to affect the system with mercury this time, as its action did not even prevent the relapse. The bowels were now kept freely open by colocynth and calomel pills at night, and salines during the day, and as the morbid action seemed to subside, so these means were gradually reduced in strength, and though much constitutional anxiety continued, yet by the middle of July she was quite convalescent. No tonics were given and no addition of stimulating regimen, merely attention being given to keeping the bowels regularly open, so as to allow of nature effecting a healthy reduction of the vascularity of the liver, and a restoration of its normal secretion.

CASE II.

The case of bilious diarrrhœa occurred in a bastard man, (Privulf,) of strong and healthy appearance, and the first symptoms of engorgement of the liver and

portal system showed itself by complaint of great pain in the loins and side, which were relieved by a Dover's powder at night, and a dose of salts next morning, which caused the evacuation of a large quantity of bile by stool. On the 7th of July they appeared again under the form of nausea, pain and soreness of the belly, loss of appetite and a very white coated tongue. After the exhibition of an emetic, the evacuation of bile manifested itself, and bilious diarrrhœa supervened, and continued more or less for several days, sometimes with increase of pain, generally spasmodic, but always attended with soreness of belly. Compound chalk powder and opium pills at night gave most relief. The diarrrhœa gradually declined, and the patient by the 10th, was convalescent.

CASE III.

Heathfield's case showed a combination of dysentery and biliousness, the engorged portal circulation seeking relief by either one mode or the other, according to circumstances. He had been five or six days ill before applying for medical aid. The symptoms were dysenteric, with a very white coated tongue, thirst, nausea, &c. June 27th and 28th he had given him two doses of rhubarb, calomel, and nitrate of potass, which relieved the abdominal symptoms very much, and operated gently, no more blood being passed. An attack of abdominal pain occurred next night, which was relieved by opium. On the 30th the bowels were natural, but nausea was complained of, with loss of appetite continuing, and white tongue, the first and last symptom obtaining alleviation from sedleitz powders. On July 2nd, as debility and anorexia seemed now the leading symptoms, decoction of cinchona was ordered, under which he improved, but it caused costiveness and flatulence, which rhubarb and soda relieved. He became convalescent by the 5th of July.

CASE IV.

Almost similar symptoms attended Mrs. W's. case, but rather less intense, and more chronic, which first came under notice June 19th. Biliousness had been present some days, with abdominal pain and white tongue. Colocynth and calomel pills were first given, and then salts, but on the 22nd dysenteric purging appeared with increase of abdominal pain. Pills of calomel and opium were given, without any effect in checking the sanguineous purging, but the tongue was now cleaner, and compound chalk powder was tried instead, and with success. On the 24th the stomach became disordered and bilious, and the tongue white, which a dose of rhubarb and soda mitigated, but did not open the bowels, which was required to be effected with the administration of rhubarb and calomel. As there still seemed a tendency to dysenteric purging when the bowels were moved but likening mucus instead of blood, compound chalk mixture was again given. Powders of small doses of calomel, opium, ipecacuanha, and gum, were next given, to act as an alterative and restore a proper state of the intestinal secretions and

their evacuations, which succeeded. On July 11th a relapse of dysenteric symptoms appeared with abdominal pain, which was relieved by opium pills, followed next morning by a dose of rhubarb and calomel; and on the 13th, as biliousness of the stomach again appeared, a dose of rhubarb and soda was given to relieve it, and convalescence seemed to be established. Subsequently the mouth became sore, and on the 17th of July a severe attack of pain in the belly, with shivering and fever supervened, with costiveness, which was subdued by pediluvium fomentation, followed next day by doses of sedlitz powders, which operated freely, and relief to all the symptoms took place. Soreness of the mouth, and a continued tendency to salivation, however, manifested itself, otherwise the patient is convalescent.

CASE V.

Pure dysentery marked the next case, in Mr. S., who reported himself sick June 23rd, though he had been suffering for a week previous. The symptoms were sanguineous purging, with its accompaniments, and attended by few constitutional symptoms, except loss of appetite. A powder of rhubarb and calomel was given, and relieved for a time all the symptoms, and as he considered himself able to undertake duty he was not put under any medical restraint. Up to the 6th of July he had four of these doses, but the bloody motions generally returned with straining, a day or two after the taking a powder, but the accessory symptoms had now subsided, the appetite had returned, and the tongue became clean. He was now kept to his room, and pills of calomel and opium regularly administered according to the recurrence of bloody stools, which were finally checked and then allowed to become natural, and by the 15th he was convalescent. No affection of the mouth took place. The compound calomel powder so useful in the former case, was tried in this before the pills, but found ineffectual, and the latter were beneficial here, but inoperative in Mr. W's. case.

CASE VI.

The last case was a decided one of dysentery, and occurred in an European named Jennings, a groom, who had been ill with it occasionally for a month before, and had usually found relief from doses of rhubarb and calomel, but did not attend regularly. On the 7th of July the symptoms were so intense, that he was taken into hospital. Pills of calomel and opium, beneficial in the last case, were here of no avail, and seemed to increase the symptoms. Opium in pill was tried, to relieve the abdominal pain, but without success; also compound chalk powder, but with no better effect. Two grains of calomel and one grain of opium, in powder, were now given every four hours, and relief at the end of twenty-four hours was experienced. There was no fever and the tongue was not very much coated, and half the above dose was tried, with some cinchona added, but the symptoms again returned as bad as ever. The full dose was again put in requisition every six

hours, with good effect, and after twenty-four hours lessened one quarter, without allowing any return of the severe dysenteric symptoms. On the 12th a blister to the abdomen removed all pain, and an injection of warm water was administered that night, with further relief. There was now mercurial fetor of the breath, approaching to salivation, but no affection of the gums. On the 13th, as I considered the amendment produced was not commensurate with the large and repeated doses of calomel, an alteration in the treatment was ventured upon. A large dose of Dover's powder, a pediluvium, and other accessory means were enforced, to produce profuse perspiration that night, and followed next morning by a purgative dose of rhubarb and soda. On the 14th, great relief was experienced and the balance of the circulation and distribution of nervous fluid seemed to have become more naturally established, the stools were now changed and took on the form of diarrhœa. Small doses of Dover's powder and hydrarg. were given night and morning, with the best effect, and compound chalk powder, should diarrhœa be troublesome, in the intervals. The stools became gradually more natural, more fœcal and bilious, and the appetite much improved, and on the 20th of July he was convalescent.

Remarks.—All these six cases will be seen to have occurred in the course of one month, leading to the supposition that similar exciting causes for the diseases were present in each case, or that a single endemic cause prevailed, as they all happened in the same locality at Shilon. The diseases all implicated the same system of organs—two being developed at one extremity of the intestinal tube, another at the other, three being intermediate. In the case of hepatitis, the hyperæmia of the liver was immediately relieved by strong purgatives, diminishing thereby the fluid contents of the portal system. The hyperæmia of the general circulation or fever, was relieved by the production of profuse sweating. The calibres of the vessels, both local and general, were thus allowed an opportunity to exercise their natural resiliency, and return to a natural size. The pressure on the secreting cells of the liver by the enlarged blood-vessels, is coincident with arrest and nonelimination of secretion, which takes place easier when that was removed. The same would occur in the general circulation. As heat is commensurate with an increase of fibrin in the blood, so is there an increase of fibrin in inflammation and fever. Fibrin is also considered as an excrementitious form of albumen, so that the object of medicine would be to endeavour to get this superabundance eliminated from the system. Its retention in the circulation leads to its deposit in organs predisposed, and in some cases with an impediment to their functions. This elimination of the fibrin can be accomplished by increasing the secretory powers of the different organs. As during inflammation little or no secretion is formed, so the biliary matter, not being liberated from the blood, would accumulate in the circulation, its more essential organic constituents would remain unformed, or in the state of fibrin, and the

colouring matter would appear to tinge the countenance, and flow off by the urine.

That the inflammation of the serous surface ever takes place, independent of any diseased action in the substance of the liver in ordinary cases of so-called hepatitis, would, I think, admit of a doubt. Pneumonia may exist independent of pleurisy, though hepatitis and congestion of the liver could not exist long without involving the serous covering, because morbid effusions have easier egress into the vacant air-cells of the former. Mercury seemed to have no ultimately beneficial effect, as a relapse of the complaint occurred after the affection of the mouth by the mineral, and during its continuance. The same effect towards the relief of the disease was induced in the relapse by medicines containing none of that ingredient, and the cure was permanent, without the mouth being again affected. On theoretical grounds, even mercury would be inconsistent with subduing increased action of the liver, unless on the principle of *similia similibus curantur*. Antimony during the height of the disease proved hostile to the stomach, causing vomiting when given either in the fluid or solid form to induce diaphoresis.

In the bilious diarrhoea most probably there exists hyperæmia of the portal circulation, the vessels of the liver not being in a diseased state at all, and therefore do not give way to dilatation; and this is rendered probable, from the paucity of bile existing in the latent state. The treatment of bilious diarrhoea under the above supposition, must evidently be palliative; the hyperæmia has found a vent, and ought not to be unduly checked. The increased quantity of bile flowing off ought to be shielded from injuring the coats of the intestines in its passage through them. Symptoms will announce if it produces morbid effects on the mucous membranes, by the occurrence of soreness if the mucous membrane has been bared of epithelium, and of pain when the exposed subjacent tissue of muscular fibre is thrown into unnatural spasm. Remedies appropriate to these conditions would require to be exhibited to afford relief, as anodynes to diminish sensibility, and to allay the susceptibility of the nervous fibrillæ. Any matter, as peculiar aliments, tending to increase the contents of the portal blood, should, of course, be avoided, such as admit of absorption by the lacteals being preferable to such as enter the veins. The allaying of the force of the disease would be safer than checking quickly its symptoms. An overloaded portal circulation may also cause its hyperæmia to relieve itself indirectly by the effusion of its overloaded contents from the thinner mucous membranes, as from the colon, producing a form of dysentery. The more depending and enlarged portion of its circulation would suffer first, as in the rectum and sigmoid flexure. The circulation in the colon may be presumed at all times less active than that of the small intestines, as it takes less share in digestion and absorption of aliment; it could not, therefore, relieve itself so easily as the upper parts, from having less activity—local activity of circulation having little to do with the force of the heart.

Want of this property, it is well known, also in a morbid sense, leads to congestion. The epithelial lining of the colon is less liable naturally to change than that of the stomach and small intestines. Should a course of diet, as an animal one, be in use, still less would the shedding and reproduction of epithelium exist, than in cases where a vegetable diet was usual. The chief current, and the strongest of the portal circulation, must be in the afferent and efferent vessels of the stomach and small intestines, that of the colon being tributary to, and more sluggish than, this great stream. Should the portal circulation become hyperæmiated, then would this congestion of the colic vessels be increased, and give a tendency to effusion. This morbid event might be either sudden, when a strong external cause was applied, or gradual, and only show itself when the pressure of the blood had reached a maximum, which would be indicated by such a symptom as a sense of expression in the abdomen. The gradual loss of blood would thus tend to prevent the accession of fever, which would have greater cause to arise when the disease was brought on suddenly before the engorged portal blood had found any relief by effusion. Inflammation would, in a sudden case, be more likely to arise also from the same cause, and be followed sooner and easier by destruction of parts, and arrest of the natural functions, which would be less likely to take place so rapidly, or to that extent and intensity, in a protracted case. Effusion of blood in these cases acts as a naturally local abstraction of blood. When the symptoms have once commenced, the nervous action of the intestinal canal is thrown out of its balance, and becomes concentrated on the diseased part, hence irregularity of peristaltic action, and want of action in the small intestines evidenced in the absence of fecal evacuation, and these of a vitiated character. As the portal circulation has now got a vent for its contents, less bile is formed, which would also arise from the loss of harmony in the action of the abdominal sympathetic nervous power. Hence medicines having the power to increase the bile, would divert the morbid effusion of portal blood, and such as tend to allay spasm would allow of the restoration of the nervous balance. Medicines also which increased the secretions and peristaltic motion of the small intestines, would be beneficial on both suppositions; but such as would increase the bile, and act like the latter, would be supposed to afford greater relief to the overburdened portal circulation, and transfer the normal current of blood, as it exists naturally in a greater amount to the small intestines.

As amylaceous articles of diet seem to be more fitted for nutriment in these cases than animal or albuminous vegetable substances, one may conjecture their peculiar fitness arises from their assimilation requiring a greater activity in the secretory powers of all the salivary system of glands than in those of the liver; and besides, from their absorption in the shape of sugar, not adding to the increase of the fibrinous elements of the portal blood.

Mercury in some form seems to exert a curative effect in dysentery, some constitutions bearing calomel, some

agreeing with hydrargyrum cum creta, &c., better. This effect does not appear to be in every case coincident with the establishment of an affection of the entire system, as evidenced in the gums, though witnessed apparently in one of the above cases. In Mr. S. no constitutional affection from the mineral supervened. The influence must then have been entirely confined to the abdominal organs concerned, and could not have passed through the liver; or, if so, must have returned with the bile, upon which secretion perhaps, after all, the necessary effect has to be produced. In other cases I have witnessed the same, and besides, have seen the mouth affected, and still the progress and violence of the disease not checked materially. The case of Mr. S. got well by checking the complaint, that of another by the supervention of a diarrhœa, either from the transference of increased action to the small intestines, or the transmutation of the dysenteric morbid processes in the colon into those of the saline affection. The occurrence of perspiration in Jennings's case was the turn of the disease, the balance of the circulation and nervous fluid seemed to be restored by it at a critical period, so that the organs essential to a restoration of health had become supplied again with a due share of each, the complaint was thus robbed of its means for carrying on the war, and health had an opportunity of resuming its empire. Whether had the affection of the system with the mercury conduced also to this end? That the mineral in the form of calomel had entered the blood, seemed to be established by increase of the saliva, and mercurial fœtor of the breath, yet the dysenteric symptoms did not very materially change for the better. I am inclined, on the whole, to consider from these cases, and from others previously under treatment, that its curative effect is local, and confined to the organs concerned in the disease in the abdomen. It may be sedative, as when calomel is combined with opium, as in Mr. S.'s case, it is slightly stimulating to the biliary and intestinal secretions, so as to restore the balance of the portal circulation. In the case of Mrs. W. and Heathfield, checking the dysenteric symptoms by simple opium, seemed to occasion biliousness of stomach, inferring that the portal blood desired to relieve itself by that channel instead, and when medicines were given to correct the latter symptoms, they did not appear to bring a return of the dysenteric symptoms. Calomel and opium in Mr. S.'s case evidently acted accumulatively, and together with rest, allowed a natural redistribution of the portal circulation and nervous fluid. The difference of the effect produced by different medicines, in these cases, seemed to have no clearer cause than constitutional peculiarity, and the cause of the difference in treatment arose from the difference in circumstances at the time, and the necessity of counteracting symptoms as they arose. The benefit subsequently shown, must have been derived in these cases of dysentery, from the previous exhibition of purgative doses of rhubarb and calomel, which took off the oppression of the portal circulation by increasing the flow of bile, thus relieving it of its load, and in an early case no doubt would be sufficient to afford nature

an opportunity to restore a healthy state of the economy of the large intestines. The diarrhœa produced in Jennings's case, was apparently of yellow bile, such being the colour of the motions, and this seems generally the colour which the subsequent diarrhœa, under the influence of mercurials, takes on. What difference, then, chemically speaking, exists between yellow and dark-coloured bile? Secretion of dark bile, I fancy, is produced in cases of hepatitis, by purgative doses of mercurials. About the same time as these cases above narrated took place, shewing hyperæmia of the portal circulation, other slight ones happened in other people, as biliousness of stomach, where colocynth and blue pill or calomel, followed by sedlitz or salts, effected cures. The causes of this general indisposition it would be almost futile to endeavour to enter upon; it might be put down to increased diet of meat from scarcity of vegetables at this time of the year; it might be from cold, but that would be just as liable to affect any other system of organs, without a predisposition. Heat is out of the question: so is moisture, as the season is otherwise distinguished, being very dry.

The diet is more vegetable in this country in summer and autumn than in winter and spring, though abdominal complaints prevail more in the latter period than in the former, while in England the prevalence occurs in the autumn, when there is more vegetable food. In this country exclusive animal diet is supposed to be the internal cause of these complaints; in England, a superabundance of vegetable food.

An increased vegetable diet would tend to increase the secretory powers of the intestinal canal, and this carried to excess would lead to such an amount as easily turn into inflammation under proper exciting causes, and the disease would be active. The warmth and dryness of the air here would keep up perspiration in ordinary times, while the coolness and moisture of that of Britain would tend to check and cause revulsion of the general fluids to the mucous membranes generally of the body. A diminished vegetable diet or a correspondingly increased animal one, would demand less than ordinary powers from the mucous membranes, as animal and albuminous vegetable food likewise, becomes easily absorbed direct from the stomach, except fat. The intestines, especially the large ones, would have less call upon them to furnish secretion, and the previous afflux of blood to which they had been accustomed in the summer and autumn, would have no vent on the mucous membrane, the volume of the portal circulation being still maintained. The same under the use of an albuminous diet. The portal blood would thus go to the liver, not much changed, as organic changes take place in it for the production of secretion, and would therefore be of a different constitution than otherwise, and there would also be less call for biliary secretions; so from want of local organic activity the portal circulation would become surcharged with unchanged venous blood, less adapted, at the same time, for the purposes of the liver. Its effusion through the thin membranes of the colon, nature may take as one way of relieving itself, and hence the dysentery be induced. In the

former state of disease the hyperæmia would be active, and probably more arterial, here congestive and venous, as the nature of the evacuated blood demonstrates.

As in the pure cases of dysentery, the overcharged portal circulation finds relief by the direct effusion of its contents, and that taking place at a limited and specified extent of the intestinal canal, so engorgement of the liver and vessels of the upper primæ viæ would occur, less than where this direct uncomplicated effusion did not take place, as in the intermediate cases of Privulf, Heathfield, and Mrs. W. In these the congestion of the portal vessels seemed more general, as there was evidently a tendency to disorder of the upper primæ as well as the lower bowels. The implication of the former being further shown by the earlier and deeper white coating imparted to the tongue, irrespective of the symptoms. The tongue, in this pure dysentery, does not seem to become affected till later in the disease, continuing tolerably clean at the commencement, and then results from the effects of local inflammation taking place in the colon, and the subsequent disorder of the whole primæ viæ. This condition of the tongue in the former cases may be conjectured to result from an arrest of the shedding and reproduction of the epithelial lining of its mucous membrane, an event common to it, and that of the stomach, &c. The old epithelium remains attached, none being fresh formed underneath, which would also occur amongst the numerous mucous glands of similar localities, hence checked secretion and congestive states of the capillaries of the mucous membranes are coincident. The removal of such a state of matters would be the restoration of secretion and of the current of blood to a healthy activity.

ON SAVINE.

By EDWARD COPEMAN, M.D.

PHYSICIAN TO THE NORWICH HOSPITAL.

Read at the August meeting of the Norwich Pathological Society.

THE late trial of Mr. Pascoe, a member of the medical profession, for the crime of procuring abortion, excited, as you all know, considerable interest, and, for a time, no little sympathy, on the part of his professional brethren. It has been since but too surely proved that, whatever might at first have been the opinion as to the justice of the sentence, his crime was of a deep dye, and deserved the punishment awarded. But the medical evidence on which he seemed to have been condemned is still open to criticism, and it is with the belief that practical good may result from a consideration of the principal point involved in it,—namely, the propriety, or otherwise, of employing savine as a legitimate medicinal agent, that I now venture to occupy a few minutes of your time with the subject. I do not remember the exact statements made at the trial by the medical witness; but the purport of his evidence was to impress upon the Jury that savine was now never given by medical practitioners for truly

medical purposes; that it was a dangerous medicine; and that, when used, it was in all probability employed with some sinister intent. It, therefore, becomes a proper medico-legal question whether or not savine can and ought to be employed as a medical agent; for the evidence I have referred to was listened to by the Judge, and probably believed by most of those who heard it; and it is important that no wrong impression on the matter should rest on the minds of Judges and Juries in future investigations of the kind.

Savine is an evergreen shrub, belonging to the natural family *conifera*; and its active qualities appear to reside chiefly in an essential oil of much the same constitution as oil of turpentine. All authorities represent it as a drug possessing considerable power upon the uterine organs; but some describe it as an exceedingly useful and efficient emmenagogue, whilst others speak of it as a dangerous and poisonous medicine. "Savine and its essential oil are emmenagogues; the fresh plant is too frequently employed by the peasantry as an abortifacient, but cannot be employed for this purpose without seriously endangering the life of the mother." (Phillip's Translation, Pharm. Lond., 1851.)—"It is, perhaps, the most powerful uterine stimulant of the materia medica, and is occasionally administered in amenorrhœa, though always requiring the utmost caution, lest it induce inflammatory action." (Brande's Manual of Pharmacy.)—"Savine is a powerful stimulant to the uterine organs, and is employed as an emmenagogue with much benefit in amenorrhœa and chlorosis depending on torpor or deficient action of the uterine system. In consequence, however, of its poisonous properties it should be used with caution; its employment is contraindicated where there is the least tendency to irritation or inflammation of the uterus or any of the pelvic viscera." (Neligan on Medicines.)—Nevin and Pereira recommend it as the most effective of all emmenagogues. I will not, however, occupy your time by further quotations, but proceed to the relation of my own recent experience of the drug.

CASE I.

Amelia Woods, about 18 years of age, was admitted as an out-patient at the hospital on the 4th of October, 1851, for chlorosis and amenorrhœa; of a pallid unhealthy complexion, weak circulation, and inclined to be fat. I ordered her a course of steel; but as her attendance was irregular, and she seemed to derive no benefit from the treatment, I admitted her into the house on the 13th of December, and gave her for a time steel and aloetic medicines. These failing, on the 20th of January, 1852, I ordered a mixture composed of Ol. Sabinæ, gttss. iv.; Mucil. Acac., oz. j.; Mist. Camph., oz. iij.; a fourth part twice a day. On the 24th six drops of the oil were put into the mixture; on the 25th, eight; and on the 30th, ten; which dose—namely, five drops daily—she continued to take until the 4th of February, on which day menstruation commenced and lasted four days, in natural quantity. No uncomfortable effect from the medicine was ever com-