ON
A VARIETY OF LATENT PNEUMONIA.

By C. M. DURRANT, M.D.,

PHYSICIAN TO THE EAST SUFFOLK AND IPSWICH HOSPITAL.

Read before the Meeting of the Suffolk Branch, held June 25th, 1882.

MANY cases of a form of pneumonia having presented themselves to my notice, both in hospital and private practice, during the past two years, in which the general symptoms in reference, equally to the patient as to the observer, have been so slight, (the disease at the same time being extensive in its pathological results,) that I have deemed it not an unfit subject to bring before the present meeting.

In the last volume of the "Transactions" of our Association, there is a paper by Mr. Caleb Williams upon "The Present Type and Character of Disease." This paper is one of great value, as setting before the profession in the clearest manner the altered type of disease generally in the present day, as combined with the character which obtained to the majority of disorders some twenty-five years ago.

I have ventured upon this digression—first, because it gives me an opportunity of strongly recommending the perusal of Mr. Williams's paper; and, secondly, because in the very latent form of pneumonia to which I am about to direct your attention, the existence of the present constitutional character of disease exerts, if not a primary, a very strongly modifying influence.

The symptoms as occurring in private practice may perhaps be best detailed by the brief recital of a case which presented itself to my notice a few weeks ago:—

The patient, a middle-aged man, and a stranger to me, commenced the description of his sensations by saying,—“Doctor, I am not well, and I cannot tell what ails me. I am not aware that I have taken cold; I have no cough, and no pain, but still I feel a sensation of weight in my side that reminds me continually that I have a side. These feelings of *malaise* (he continued) have been upon me for nearly three weeks.” Upon inquiring if his breathing were affected, he replied:—“Now you direct my attention to the circumstance, it certainly has been somewhat short at times, but I have not thought much of it.” This gentleman's countenance was dull, and his cheeks reddish; the eyes heavy, and the conjunctivæ slightly yellow; tongue not much coated, and moist; pulse 80, of moderate strength; breathing at the time of examination not accelerated; bowels sluggish; urine occasionally depositing a sediment. His appetite, he said, was “too good,” and he slept well, and the temperature of the skin was but slightly above the natural standard. Upon baring the chest for the purpose of making a sthethoscopic examination, I found that the ribs upon the right side acted less freely than on the left, and over a very large portion, nearly two-thirds of that lung, the respiratory murmur (unaffected by position,) was almost inaudible, while the sound elicited by percussion was characteristically dull to a corresponding extent.

This case required a steady perseverance in the treatment presently to be advised, before a return of the feelings of health could be obtained, and which occurred simultaneously with the restoration of the natural breathing.

I have endeavoured thus to give, in the somewhat graphic language of the patient himself, a description of the very few symptoms, and these the most prominent, which presented themselves in this obscure form of pneumonia, and which, whether occurring in what was formerly more particularly designated the plethoric constitution, or its reverse; or whether, attacking the well-fed among the higher classes, or the labourer and artisan, still the symptoms will be found to bear no proportion to the extent of lesion to which the disease gives rise. The affection under consideration is essentially characterized by negative results,—viz., an absence of those symptoms to which we generally refer as evidencing a serious interference with the healthy function of respiration. For example:—We have seldom rigor complained of at the onset; the countenance, although dull and heavy, is not anxious; debility is almost universally complained of as a leading symptom; there is in general no severe pain; the cough, if there be any, is not distressing, and the amount of expectoration either inappreciable, or if more abundant, it is untinged mucus; the breathing may be short, but not sufficiently hurried to attract much attention; the *alæ nasi* are quiescent; the skin is in general moist, and free from that pungent heat so characteristic of the more acute form of attack; the pulse is either natural in frequency, or but slightly accelerated, rarely above 90; the urine is for the most part free from deposit, and in sufficient quantity. The physical signs, however, are well marked, and will, if carefully sought for, unfailingly indicate the true nature of what, without this aid, must often be doubtful and embarrassing.

The tendency of this form of pneumonia, so far as my own observations have led me to observe, has been towards recovery. Tedious,—very tedious, both to patient and practitioner, it will sometimes prove; but I have not found, provided that the subject of the attack be not already labouring under phthisis, that it either predisposes to, or merges into that malady.

The cause of this disease is not, I believe, so much attributable to cold as to a disordered condition of the primary and secondary assimilative processes, inducing as a consequence a morbid state of the blood itself; this, by depressing nervous power, subsequently causes in its transit through the lung, irritation and congestion, followed by adynamic inflammation, with exudation. The disease may pass through its different stages with so little disturbance to the pulmonary organs, that, as in the case above referred to, the patient will attribute his disordered feelings to any but the right cause.

Since writing these notes I have seen another instance in which the patient, unconscious of the existence of any serious lung affection whatever, (but in whom, nevertheless, more than one half of the left lung was in a consolidated state,) attributed all his discomfort to an affection of the stomach. The treatment that I have

hitherto found to be most beneficial has been chiefly confined to counter-irritation, the guarded introduction of mercury, (with a view to the correction of morbid secretions, and carried to the extreme of very slightly affecting the gums,) the iodide of potassium, and cod-liver oil.

Venesection is in no case called for, as a dangerous degree of depression may be expected to follow the use of depletory measures. Cupping, or even leeches, will seldom be needed, unless an unusual degree of pain be complained of, which is quite contrary to what ordinarily obtains. Blisters will be found to be of signal service, and these must be repeated again and again, not kept open, but permitted to heal; and their renewal, as well as any change of locality, must be influenced by the result of carefully-repeated stethoscopic investigation. Should any indication of an indurated portion of lung remain after the application of from three to five blisters, at weekly intervals, a strong solution of iodine (one drachm of iodine, and half a drachm of iodide of potassium, to one ounce of rectified spirit,) painted, if it can be borne, once daily over the affected spot, will be found at this stage very useful. If the secretions be much deranged, small doses of the hydragrym cum cretâ should be given, slightly, and only slightly, to affect the gums; or what will then be found to be a still better form is the bichloride, in doses of one-sixteenth of a grain, night and morning. Iodide of potassium, with alkaline salines, will be called for, to be followed by cod-liver oil, in doses from a teaspoonful to a tablespoonful thrice a day. This latter remedy has given me much satisfaction in the treatment of pulmonary indurations of an adynamic character; and it has exerted a very marked influence in promoting the absorption of morbid exudation. This, I presume, it can alone effect by improving the condition of the general health.

ON THE

TREATMENT OF ERYSIPELAS BY THE MURIATED TINCTURE OF IRON.

By W. H. RANKING, M.D., CANTAB.,

PHYSICIAN TO THE NORFOLK AND NORWICH HOSPITAL.

Read before the Meeting of the Suffolk Branch, June 25th, 1852.

It has always been my opinion that the communications most suitable to the brief reunions which take place at these anniversaries of the several Branches of the Provincial Medical and Surgical Association, are those which are able to convey practical information in few words; for this reason I have selected for your attention a case which has recently come under my notice, and in which a method of treatment not commonly followed has proved eminently successful. I allude to the treatment of erysipelas by the muriated tincture of iron. The case is as follows:—

Charlotte Andrews, aged 23, a fat strumous girl, was

admitted into the Norfolk and Norwich Hospital under my care, in April, for engorged and suppurating cervical glands, which were treated by generous diet, cod-liver oil, free lancing, and subsequent dressing with the iodide of lead ointment, an application which I may state in passing, appears to me to be more suitable to scrofulous ulcers than any other with which I am acquainted. Under this treatment her progress was all that could be desired, until the 26th of May, when my attention was called to an erysipelatous redness over the right breast. This quickly extended, and in spite of the assiduous endeavour on the part of our intelligent house-surgeon, to limit its boundaries by the nitrate of silver, it soon invaded the head and face, inducing those serious symptoms which are observed in these cases. As soon as the disease fully declared itself, no time was lost, as is my invariable custom, in sustaining the powers of the patient by wine, beef-tea, and ammonia, the local application being flour.

On the 28th, the disease had continued to extend, and as the patient was losing strength the ammonia was replaced by quinine.

On the 29th, the report is that the erysipelas had occupied the entire head and face, and was creeping down the back. The patient was delirious, with dry tongue and feeble pulse of 130.

On the 30th she was still worse, the tongue was more dry and dark, and diarrhoea was added to the other unfavourable symptoms. Wine was given in increasing quantities, so that she took more than a bottle in the twenty-four hours.

June 1st.—She was still more depressed; pulse 140, fluttering; the face livid, and she appeared to be fast sinking. At this time I determined to administer the muriated tincture of iron, as recommended by Mr. Bell, of Edinburgh, and did so, as will be seen, with the best results. The dose was forty minims in water, every three hours. In the evening, after three doses, the pulse was still 140, but had more resistance to the finger, and she was left for the night, with orders to continue the medicine and wine.

On the 2nd there was a most marvellous change for the better. The tongue, which the day before was dry and dark, was now moist and cleaner, the pulse had sunk to 120, and the patient was able to raise herself in bed. The face was desquamating and her only complaint was urgent thirst, which was gratified with water *ad libitum*.

3rd.—I continued the iron in doses of thirty-six drops. The improvement was still more manifest; the face had become more natural in appearance, and she was pronounced out of danger. From this time her recovery was uninterrupted.

Knowing how difficult it is to establish a medical *fact*, I am prepared to find that some of my hearers may dispute the agency of the iron in this girl's recovery. On this point I would only say, that prior to her taking that medicine she lost ground hourly, in spite of the freest exhibition of wine, ammonia, and quinine; and that after three or four doses a perceptible advantage had been gained, which advantage was rendered indisputable on the