may contain some later in the day. This form of albuminuria has no tendency to develop into chronic inflammation of the kidneys, and may last for years without getting worse. Such patients should avoid everything that could irritate the kidneys.

4. *Functional Albuminuria*.—This name I should like to apply to that form which appears in diabetes without any other symptom of chronic disease of the kidney, and where the only cause discoverable is the increased flow of sugar from the kidney. This albuminuria is always slight, very rarely are there more than traces of albumen present. There are no casts or other morphotic elements. This form of albuminuria is chiefly to be found in diabetics who have been passing a large quantity of sugar for a considerable time. If the amount of sugar excreted is reduced, the albumen very often in a little time disappears entirely. The explanation of the albuminuria in these cases may be found in the irritation of the kidneys by the excessive flow of urine loaded with sugar. It is not certain, though it is probable, that at the time of the albuminuria there are real pathological changes in the kidney, as, for instance, slight inflammation and hyperæmia. After death, the kidneys of diabetics are often found enlarged, hard, and hyperæmic. Maguire found that the kidneys of persons who had had diabetes of long duration were often hard with much interstitial connective tissue round the vessels and glomeruli, and between the tubuli. He thinks that this increased growth of connective tissue is a result of the continuous overwork, and a preliminary stage of chronic granular kidney.

5. *Albuminuria produced by Chronic Disease of the Kidney*.—Although Frerichs has denied that chronic inflammation of the kidneys could be caused by diabetes, I agree with Senator⁹ who mentions diabetes mellitus as one of the causes of chronic nephritis. Which form of chronic inflammation is chiefly produced is a question not yet settled, for while some authors, for instance Ingleissis¹⁰ and Seegen,¹¹ say that parenchymatous nephritis is the form of chronic kidney disease chiefly to be observed, others, for example v. Noorden,⁴ declare it to be almost always granular kidney. My own observation agrees with that of the former writers, for I have often seen the symptoms of parenchymatous nephritis, and but rarely those of granular atrophy. In one case I had the opportunity of making a *post-mortem* examination, and there I found pathological changes of the parenchyma as well as an increase of the interstitial tissue. Of my 191 cases of diabetic albuminurics, 44 suffered from chronic Bright's disease. Of these 5 had gouty kidney, while in 17 cases chronic alcoholism, and not the diabetes, was the probable cause of the affection of the kidneys. There remain 22 cases for which no other cause than the diabetes could be detected. I cannot here give an analysis of those 22 cases, but I may quote one instance to show how the nephritis develops in these cases.

A lady, aged 56. I first saw her in 1893, and each year since then she has been under my observation for six weeks or two months in the summer. Her diabetes dates from 1839, when much sugar and a trace of albumen were passed. Under treatment there was great improvement, and in 1893 sugar only appeared in the urine when the diet was not properly attended to. There was no albuminuria. This condition lasted till 1895, when the diabetes became more obstinate, and albumen reappeared for the first time, but again disappeared when the glycosuria was reduced by a strict diet. In 1896 the albumen had returned, and could not be got rid of; there were no casts, and the quantity of albumen was small. In the spring of 1897 there were the symptoms of chronic kidney disease—much albumen, casts, blood corpuscles, etc., in the urine, slight œdema, and spots of gangrene of the skin on both feet. Now (April, 1898) the condition is again worse. There are attacks of nausea and vomiting, probably of a uræmic nature, general debility, and œdema of both legs. The urine contains 0.6 per cent. of albumen, traces of peptone, casts, etc. Only a very little sugar is excreted on a diet rich in carbohydrates, while with a fair amount the urine remains sugar free.

The development of chronic nephritis in diabetic patients is generally to be explained as follows: The constant and excessive flow of sugar-containing urine through the kidneys leads first to irritation of the epithelium of the tubuli uriniferi, to hyperæmia of the kidneys, and functional albuminuria. If the irritation continues it leads to inflammation of the parenchyma, and increased growth of the interstitial connective tissue. Diabetic nephritis is much more common in men than in women, the proportion in my cases being 16 to 6. Pathologically and clinically very different from the diabetic nephritis is gouty nephritis with glycosuria. It is not always easy to distinguish at once between the two, but it can be done with some degree of certainty when the whole history of

the case is known. The chief difference between the two affections is afforded by the glycosuria; in gouty nephritis with glycosuria the latter is and has been almost always slight, while persons with diabetic nephritis almost invariably have had for some time a considerable glycosuria. Pathologically, gouty nephritis means almost always chronic granular degeneration of the kidneys.

TREATMENT.

As regards the treatment of the functional albuminuria the chief object we have to attain is the reduction of the irritation of the kidneys. This we accomplish by diminishing the glycosuria, which, as everyone who has had any experience of the disease knows, can only be done satisfactorily by dieting. In cases of real chronic nephritis it is often difficult to treat both affections satisfactorily, and one has to try not to do harm in one way by doing good in the other. If, as sometimes occurs, the glycosuria ceases altogether, or at least becomes quite inconsiderable, as the nephritis develops, the treatment is somewhat easier, as the dieting gives less trouble. On the whole, the prognosis in these cases is very grave. The cessation of the glycosuria seems to show that the tissue of the kidneys is more changed than in those cases in which there is still some excretion of sugar.

If milk is well tolerated I prescribe it freely. Mineral waters can be beneficial, especially those of Neuenahr, after the use of which I have sometimes seen decided improvement. If the patient has sufficient strength warm baths may be made use of with advantage. They relieve the kidneys by increasing the action of the skin. That it is essential to keep these patients quiet, or even in bed for some time, and to avoid everything that could increase the irritation of the kidneys need scarcely be pointed out.

REFERENCES.

¹ Senator, Die Erkrankungen der Nieren, in *Specielle Pathologie u. Therapie*. Herausgegeben von Nothnagel. ² Schmitz, *Berl. klin. Wochenschr.*, 1894, No. 5. ³ Quoted by Jacobson, *Gazette des Hôpitaux*, 1894, No. 94. ⁴ v. Noorden, *Die Zuckerkrankheit*, 2te Aufl., 1898. ⁵ Bussiére, *Thèse de Paris*, 1893. ⁶ Maguire, *BRITISH MEDICAL JOURNAL*, vol. ii, 1886, p. 543. ⁷ Ebstein, *Deutsches Arch. f. klin. Med.*, 1881, Bd. xxviii, s. 183. ⁸ Fichtner, *Virchow's Archiv*, Bd. 114. ⁹ Senator, *loc. cit.* ¹⁰ Ingleissis, *Le Rein dans ses Rapports avec le Diabète*, *Thèse de Paris*, 1885. ¹¹ Seegen, *Der Diabetes Mellitus*, Berlin, 1893.

NOTE ON A CASE OF RECURRENT SARCOMA OF THE FASCIA OF THE BACK, TREATED BY COLEY'S FLUID.

By A. MARMADUKE SHEILD, M.B., F.R.C.S.,
Assistant Surgeon to St. George's Hospital.

IN the *BRITISH MEDICAL JOURNAL* for January 23rd, 1897, I published a case of recurrent sarcoma of the mammary gland treated by Coley's fluid (the mixed toxins of streptococcus and micrococcus prodigiosus), where the patient died of pyæmia, apparently excited by the necrosis of the growth. I have now to report a case which shows many features of interest, since rapid improvement ensued after injection; but after a time the growths again increased, the remedy seeming to have little influence over them. The injection method in this instance was thus carried out. On each occasion the surface of the skin was carefully purified, the needle was sterilised, and the puncture made by it at once covered with iodoform and collodion. Pure carbolic acid was kept in the barrel of the syringe between the intervals of use. The fluid was tested and found sterile, and the syringe was filled by plunging the needle through a septum of fine gutta-percha tissue stretched over the bottle which contained it. The puncture thus made was at once closed with wax. The fluid was kindly provided by Professor MacFadyen. During the intervals between use it was kept in a cool chamber. The following is a very abbreviated account of the case. The leading features and circumstances are alone reported. I am indebted to Mr. Roland Whipham, the Assistant Surgical Registrar, for careful notes of the case.

The patient was a well-nourished, healthy-looking woman of about 60 years of age. She first came under Mr. Warrington Haward's care in October, 1895. There was then a livid-coloured hard tumour in the posterior lumbar region on the right side, which was covered by arborescent vessels, presenting the usual appearance of sarcoma. There was a history of eight years, with rapid growth of late. The tumour was freely re-

moved and the wound soon healed. The microscope showed a fibroma; the deeper parts were undergoing change into spindle-celled sarcoma.

On April 21st, 1896, the woman was admitted for several recurrent nodules in the scars, which were again removed.

In November, 1896, another removal. The tumour was large and formidable this time, very soft and gelatinous, and full of blood. The operation was very extensive. The mass was freely cut away until the lumbar fasciæ were exposed. On each of these occasions spindle-celled sarcoma was declared by the microscope, the cells being large and plump, and the growth of a very malignant aspect.

She was again admitted into St. George's under my own care on September 15th, 1897, with extensive recurrence, the tumours extending from the tenth rib to the iliac crest. Another free removal was done right down to the subperitoneal tissue. Dr. Rolleston reported that the microscopical sections were becoming more and more malignant in appearance.

In February, 1898, she was again admitted under my care. A tumour the size and shape of a very large potato reached from the right lower ribs to the crest of the ileum. The skin was stretched and thin at one spot superiorly, as though the growth would soon fungate. The colour was a livid blue, with arborescent vessels. The growth was so soft as to fluctuate like a hæmatoma or abscess, for which, apart from the history, it might readily have been taken. Below the crest of the ileum was another soft elastic tumour the size of an orange. This, the patient said, had rapidly formed. The accompanying photograph by Mr. Drake-

Brockman well shows the extent of the larger tumours. The general health and condition of the patient were good. Seeing the hopeless nature of this case for operation, I suggested treatment by Coley's fluid, and an account of the injections is now given. The larger tumour was alone injected, and a part was selected where the skin was thick, so as to obviate any tendency to sloughing near the punctures. The fluid was thrown into the centre of the tumour by a long needle.

On February 11th, at 4 P.M., the first injection of miiij was given. The effects were very alarming. Within an hour a severe rigor occurred, the temperature rising to 102°. She was faint and felt very ill, and the bowels acted involuntarily. Under warmth and stimulants she improved, and the next day the temperature fell, and the patient seemed better.

The tumours became intensely painful, and it was singular to observe

that the lower tumour, which was not injected, seemed the more affected of the two. The skin over the tumours was of that roseate hue so often described. For some days the temperature remained above the normal, and the tumours were inflamed and exquisitely tender. There was some slight broncho-pneumonia, which cleared up, and herpes formed on the lips and face.

On February 21st it was noted that the larger tumour, at the site where the skin was so thinned and prominent, threatened to slough. An area the size of a florin was blackened and obviously dead. The lower tumour was distinctly smaller.

On March 4th the black slough separated from the upper tumour, and some necrotic tissue could be seen beneath. The cavity was painted daily with liquefied carbolic acid, and dusted with iodoform gauze.

By March 9th it was noticed that the lower tumour was very decidedly smaller. Slough was discharged daily from the upper tumour, which was somewhat flattened and reduced in size. The latter was again injected in its lower part, m of the fluid being employed. There were no constitutional symptoms after this, but the tumour became more red and painful.

By March 15th the smaller tumour had shrunk to the size of a walnut. The larger tumour was also smaller, and strikingly soft and diffident.

On March 17th m of the fluid was injected. There was slight constitutional disturbance, and trivial local reaction.

On March 19th m of the fluid was injected. There was practically no reaction.

On March 30th a large dose (miv) was injected. There was no reaction. On April 1st it was noted that the lower tumour remained as before; it had shrunk very materially. The upper tumour was also much flatter.

On April 6th the large dose of miv of the fluid was injected. There was no reaction.

At the end of April it was obvious that the growth was again extending. The lower tumour began to progressively increase, and the upper tumour to rapidly extend both laterally and below. The patient became dissatisfied with the treatment, and went out. I have since heard that symptoms are developing indicative of implication of the spinal cord. The lower tumour in ten days increased to quite its former size, and the upper one also extended with rapidity.

In any novel method of treatment it is most important that the true results should be noted. The injection of sarcomatous tumours by Coley's fluid has now been carried out in a considerable number of cases, yet comparatively few really successful instances of disappearance of indubitable sarcoma have been recorded, leading to the inevitable conclusion that the remedy has given disappointing results. It may be remarked that it is not enough that tumours should wither away and disappear under its influence. Some sort of hope of permanency of cure must also be promised, or the remedy affords little advantage over the ordinary methods of operation. Many of the reputed successful cases have not been watched afterwards for long enough periods to note the length of time that freedom from recurrence may last. It may also be pointed out that cases have recently been reported that are calculated to somewhat shake our faith in our clinical knowledge of sarcomatous growths. The diagnosis, especially the microscopical diagnosis, of sarcoma from inflammatory formations is distinctly dubious, and growths which present all the microscopical features of sarcoma, have undergone spontaneous resolution from sloughing or atrophy. This has been from time to time noted about the jaws or naso-pharynx. All this is mentioned to show the various sources of error which may occur. Perhaps the close simulation of sarcoma by inflammatory formations is hardly enough guarded against, by those who employ the injection method extensively, and report disappearance under its use. We are all apt to fall into the too common error of attributing ill-understood pathological changes to whatever remedy, the patient happens to be employing at the time.

The case here shortly reported was, however, certainly sarcomatous. The clinical history, and the repeated microscopical examinations, left no room for doubt. It was of the spindle-celled variety, situated on the fascia, and seemed a peculiarly suitable case for the method. There could be no doubt that the fluid powerfully influenced the tumours, yet I cannot but think that the phenomena were largely inflammatory, and the actual increase of the growths which later occurred, while the patient was receiving large injections of the fluid, showed the futility of proceeding further. Had the growths in this case been small, say of the size of a pigeon's egg, one can easily understand they may have disappeared altogether. How far the disappearance would have been more permanent than that affected by operation it is difficult to define. It would be of great importance if a collective investigation could be made of the results of injecting these tumours, say two years after the treatment. The reports of cases are most contradictory, and the unfavourable results in my own two reported cases of undoubted sarcoma, have not impressed me with the value of the method. Nevertheless, in the face of the experience of others, I would not reject the

remedy on such small evidence, but when occasion offers I would give it a further trial, selecting only cases of undoubted sarcoma for its employment.

EXCISION OF THE UPPER END OF THE TIBIA FOR MYELOID SARCOMA AS A SUBSTITUTE FOR AMPUTATION :

WITH A RECORD OF TWO CASES.

By CHARLES A. MORTON, F.R.C.S.,

Professor of Surgery in University College, Bristol; and Surgeon to the Bristol General Hospital.

It has been a well-recognised fact that myeloid sarcoma is the least malignant form of sarcoma of bone. Several years ago Gross estimated that it recurred after resection of the affected bone, or amputation through or above it, in 1 in 12½ cases, and that it gave rise to metastatic deposit in 23 per cent. of cases.¹ Quite lately Mr. Bland Sutton has stated² that if these tumours have not perforated their capsule they may be safely removed without fear of recurrence, and that they do not disseminate, and he suggests that they should no longer be called myeloid sarcoma, but myeloma. Mr. Clutton also considers that they so seldom return after removal that they may almost be looked upon as benign growths.³ In several cases, instead of amputating the limb, the portion of bone in which the myeloid growth occurred has been excised, and there has been no recurrence. Mr. Morris has recorded a case of resection of the lower end of the radius for myeloid sarcoma, and the patient was perfectly well thirteen years later,⁴ and Mr. Lucas one of resection of the lower end of the ulna for the same form of growth, and the patient was also well ten years after operation. A patient of Mr. Sutton's, from whom a myeloid sarcoma of the sternal end of the clavicle had been excised, was shown at the Clinical Society free from recurrence nearly four years later.⁵ Mr. Clutton has recorded⁶ one case of excision of the lower end of the radius for myeloid sarcoma, with no recurrence at the time of the patient's death from Bright's disease eighteen months later, and another of the same kind with freedom from recurrence nearly two years after operation.

Excision of the affected portion of bone has now been accepted by most surgeons as the recognised method of treating such cases if the removal of bone can be carried out so as to leave a useful limb. I can find no record of a case in which this has been successfully done in the femur or tibia, though it has in the upper end of the humerus.⁷ I can only find the records of two cases in which it has been attempted. Billroth⁸ excised the lower end of the femur in one case of central sarcoma, but the patient died of pyæmia. The excision was performed because the patient refused amputation. In another case he excised the lower 3 inches of the tibia for "pulsating central osteosarcoma," but in this case gangrene occurred, and this patient also died of pyæmia.⁹

I have now excised the upper end of the tibia for myeloid sarcoma in two cases. In the case first operated on the patient has a most useful limb, and in the other is beginning to walk without support. In the first case I determined to try to save the limb, but the patient consented to amputation if I thought during the operation that it would be the best method of dealing with the disease. In the second the patient absolutely refused amputation under any circumstances. It seemed to me that the shortening could easily be made up for by a high-soled boot, but what I feared was that after removal of so large an area of bone, when I approximated the tibia to the femur, the tissues in the popliteal space, instead of retracting to a sufficient extent, might kink, and thus, by obstruction of the popliteal vessels, cause gangrene in the leg. That the operation can be performed without danger of kinking of the vessels is now evident, and that we may get perfect osseous union and a very useful limb is shown by my first case. That even in cases in which the union permits a little movement the limb may be fairly useful is also evident, and I am inclined to think that in the course of a few months the limb will have become much more useful; if not a second operation for refreshing and screwing the ends of the bones could be undertaken.

But all endosteal sarcomas are not myeloid, and the question will arise as to the possibility of the diagnosis of myeloid growth before operation. Fortunately, most central sarcomas are myeloid. Gross estimated their frequency at nearly 70 per cent. of all endosteal sarcomas of the long bones. Out of 8 cases of central sarcoma of the upper end of the tibia in St. Bartholomew's Hospital referred to by Mr. Butlin and Mr. Colby,¹⁰ only 1 was not myeloid. Slow growth and pulsation would be conditions in favour of myeloid tumour, and so would evidence of cyst formation if we could get it. I do not think we should do wrong in making resection the rule in endosteal sarcoma at the articular ends; and if on examination of the growth after removal it was found not to be myeloid, the patient should be very carefully watched, and amputation performed at the earliest sign of local recurrence.

We must recognise the fact that some cases of myeloid sarcoma have been successfully treated by scraping out the growth from the interior of the bone, but this method does not seem to me a satisfactory way of removing a tumour. The results in several cases after this method of treatment have, however, been satisfactory. Of the 8 cases already referred to of myeloid sarcoma in the head of the tibia at St. Bartholomew's Hospital, 4 were treated by scraping, and it is stated¹¹ that "two of the patients were well two years later, but one had a tiny sinus leading into the cavity from which the tumour had been removed; the third was well at the end of eight years, and the fourth patient suffered from recurrence at the end of two years." Dr. Hinds¹² has recorded a case of myeloid sarcoma of the lower end of the femur treated by scraping, with no recurrence two and a half years later; a small sinus persisted.

CASE I.—G. T., aged 40, came under my care at the Bristol General Hospital, January 10th, 1897. Six months before, he had suffered from a little pain just below the left knee, and then a swelling appeared there. He had no pain at the time I saw him, but only a feeling of weakness in the limb, and was able to continue his employment, which necessitated a good deal of work for the lower limbs. There was a smooth, round, tense, fluctuating swelling, the size of a walnut, projecting from the head of the tibia, just on the inner side of the ligamentum patellæ, and slightly pushing the latter outwards. There was distinct "eggshell crackling" over the lower part of the swelling. The needle of the exploring syringe entered for two inches, and then struck bone, and straw-coloured fluid was withdrawn. The whole of the head of the tibia was not expanded. As the fluid was not like that of hydatid cyst it seemed more likely that the cyst was only part of a sarcomatous growth, and on January 21st, when an exploratory incision was made, this suspicion was confirmed. A portion of tumour tissue removed and examined microscopically showed the structure of myeloid sarcoma. This exploratory incision healed by primary union.

On February 1st I operated by the method which I will now describe, and which I planned so as to save the popliteal vessels and external popliteal nerve from damage. I did not expect to be able to preserve the anterior tibial vessels from injury, but was successful in doing so, so that no important vessel or nerve was divided. After applying Esmarch's bandage and tourniquet I first made an incision across the front of the knee, just below the patella, from a point midway between the front and back of the limb on one side, to a similar point on the other side. This was an incision equal in length to half the circumference of the limb. The skin and subcutaneous tissues were then separated from the patella, and retracted upwards, and the joint opened by an incision around the upper border of that bone. It is most important that the patella shall be left, as this forms a useful handle for levering the head of the tibia forwards when clearing its posterior surface. The lateral ligaments and the crucial ligaments were then divided, and the knee-joint strongly flexed, with the sole of the foot resting on the table to steady the limb. By drawing forcibly forwards the head of the tibia by means of the patella and ligamentum patellæ, I was now able to clear the upper end of the tibia and fibula from the tissues lying behind them for the extent of three inches. In this mass of tissue lay the popliteal vessels, but they were not seen. The popliteus, the biceps tendon, and the tendons of the inner hamstrings were separated from the bones. This separation was carried out mainly with the scalpel, but in parts by a blunt instrument.

I now carried two incisions vertically down the leg, one from each end of the previous transverse incision, and joined the lower ends of these by another transverse cut extending across the leg 3 inches below the first. These four incisions enclosed a quadrilateral area of skin, which included the portion over the tumour, and corresponded in its vertical extent to the amount of bone to be removed. The limb was then turned on its side, and the upper ends of the tibia and fibula (now detached from the soft tissues on their posterior aspect) made to project from the lower end of the femur by manipulation of the leg. I then sawed through the tibia at what I thought was a safe level below the growth, and then easily dissected the upper 3 inches of the bone off the muscles in the interosseous space, and, after division of the ligaments of the superior tibio-fibular joint, removed it. I then cleared a corresponding length of fibula, taking great care not to cut the anterior tibial nerve which was not seen, and the bone was divided with the saw at the same level as the tibia. The anterior tibial artery was then seen passing between the bones uninjured, and the external popliteal nerve was also seen to be uninjured. A slice of bone, one-third of an inch thick, was then sawn off the lower end of the femur. As many small cut vessels as could be recognised were then secured in forceps, and the tourniquet removed. No large vessels bled, but numerous small points of oozing had to be dealt with. The cut sur-