nation suggested; a similar inquiry was undertaken in connection with the dorsal dullness normally found in childhood, and an analogous experiment was conducted in the body of an infant aged 3 months, the subject of general miliary tuberculosis. Precisely the same results were obtained in every point, with the exception of the relative position of the pancreas, which was found to be apparently lower, namely, below the level of the twelfth rib, probably in connection with the more elevated position of the ribs, peculiar to the chest of infants. In this case, also, a thin wedge of lung tissue extended downwards beyond the level of the tenth rib; but with that exception no other organ shared with the liver the horizontal section of space defined by the four pegs.

It may thus be definitely stated that, where it occurs in the adult as the result of extreme wasting of the body and of the lungs, the dullness is directly produced by the liver, by the pancreas, and by the intervening solid parts (perhaps slightly also by the kidney); and that it is favoured by the imperfect expansion downwards of the lung, and, in most cases, by a low stomach. In healthy children these conditions probably obtain as normal anatomical features. In pericardial effusion we may provisionally accept an analogous explanation for the dorsal dull patch, the left lung being considerably compressed, the liver much depressed anteriorly, the thoracic portion of the stomach partly displaced into the abdomen, and the dullness special to the liver greatly intensified by the mass of fluid with which that organ is capped.

PRACTICAL CONCLUSIONS AS TO THE VALUE OF THE TEST.

The occurrence of the dullness normally in children, and pathologically in some cases of extreme wasting in the adult, does not detract from the practical value of the test, since in both instances, when pericardial effusion occurs, its detection by the usual methods of percussion is favoured by the smallness of the chest and the thinness of its walls; whilst its absence can be readily made out. Difficulties in diagnosis more commonly arise in non-emaciated adults. But among the many healthy adults whom I have examined for this sign, I have not hitherto succeeded in finding a single instance of the presence of the dorsal dull patch as a personal peculiarity. I therefore regard its occurrence in them as always suggesting a strong suspicion of pericardial effusion.

In cardiac enlargement, inasmuch as the heart may depress the liver and compress the lung, the dullness in question might be expected; I have not, however, found it in any of the adult cases which I have examined. In young subjects this is a source of fallacy to be carefully borne in mind, as was shown in the case of a young girl the subject of chronic heart disease and of recent pericarditis, in whom paracentesis was performed and no fluid obtained. In this case I had diagnosed the presence of fluid on the strength of the anterior percussion of the chest, which owing to an unsuspected complication simulated that of effusion, and I persevered in the diagnosis in spite of the circumstance that the dorsal patch, which was present, did not yield absolute, but partial dullness only.

With the reservations which have been indicated, the practical value of the sign remains unimpaired, and whenever other signs point to the probability of effusion, the presence or the absence of an absolute dullness of the exact shape and situation described becomes a special test which may decide the question conclusively, and which I regard as trustworthy.

AN ADDRESS

ON THE

USE OF ANTISEPTICS IN MIDWIFERY,

WITH

SPECIAL REFERENCE TO CORROSIVE

SUBLIMATE.

Delivered before the British Gynaecological Society.

By CLEMENT GODSON, M.D.,

President of the Society, and Consulting Physician to the City of London Lying-in Hospital.

[AFTER some preliminary observations, in which he referred to an opinion which had been expressed as to the danger of mercurial poisoning in consequence of the use of solution of corrosive sublimate in obstetric cases, Dr. Godson said that he proposed to devote his address to an examination of the results obtained at the City of London Lying-in Hospital,

which would be found to corroborate the testimony given respecting the General Lying-in Hospital since corrosive sublimate was introduced there as an antiseptic by Sir John Williams and Dr. Champneys in May, 1884, who followed the example which had been set by Dr. Tarnier in the Maternity Hospital of Paris. He proceeded:]

It was the publication of the admirable results obtained by the physicians at the General Lying-in Hospital that decided us to discard carbolic acid in favour of corrosive sublimate as the antiseptic when the City of London Lying-in Hospital was reopened on July 1st, 1886, after having been closed for a short period for thorough cleansing, as was the annual custom. The strength of the solution used was 1 in 1,000 for the hands of the attendants, and 1 in 2,000 for vaginal irrigation. I gave a full account of how it was employed in a lecture delivered at the Midwives Institute and Trained Nurses Club on October 14th, 1887, which was subsequently published and distributed among the midwives and nurses in pamphlet form; and, except in small details, it has been used in the same manner up to the present time, December 31st, 1896, just ten years and a half, on 4,608 women, without a suspicion of mercurial poisoning in one; therefore I think we may at once dismiss the idea of any risk of poisoning in the way in which it is now used, and should be used, and it will probably be found that what Dr. Bantock refers to relates to when intrauterine injections were employed of a strength of 1 in 1,000, or stronger, such as those reported by Dr. Dakin in a paper on Mercurialism in Lying-in Women undergoing Sublimate Irrigation, read at the December (1886) meeting of the Obstetrical Society of London. Dr. Dakin brought the subject forward not with the view of advocating the discontinuance of its use, but to endeavour to elicit the best means of employing it with safety, for he says "the results at the General Lying-in Hospital have been such as to make the physicians loth to dispense with it."

In an admirable address² by Dr. Cullingworth on Puerperal Fever a Preventable Disease, he says: "Until the year 1877 this hospital" (the General Lying-in) "was scarcely ever free from puerperal fever, and the mortality, always high, occasionally became fearful. In 1838, of 71 women delivered, 19 died; in 1861, 14 died out of 195; and in 1877, 9 out of 63. On several occasions the hospital had to be closed for long periods, and thousands of pounds were spent on the sanitary improvement of the building. In October, 1879, this institution, having been closed for two years, was reopened, and has since been conducted on antiseptic principles, the details varying from time to time as increased knowledge and experience have dictated. Mark the result," and he gives the following table:

Period.	Deliveries.	Deaths.	Average Death-rate from All Causes.
1833 to 1860	5,833	180	1 in 32½ = 3.088 per cent.
1861 to 1877	3,773	64	1 in 58½ = 1.696 per cent.
1880 to 1887 (antiseptic period)	2,585	16	1 in 161½ = 0.618 per cent.

This last period of eight years takes in three varieties of antiseptics used, first, carbolic acid; next, permanganate of potash; and finally, corrosive sublimate.

Dr. Boxall, in a masterly paper read at the June and July 1890 meetings of the Obstetrical Society (which I recommend to the attention of you all) gives another period, from the time corrosive sublimate alone was used—May 1st, 1884, to June 30th, 1889—2,150 deliveries, 9 deaths, 1 in 239=0.418 per cent., a much greater diminution in the death-rate.

I propose to contrast the results at the City of London Lying-in Hospital from the time that I became connected with it, in March, 1870, up to the end of April, 1886, a period of 16 years, with that commencing July 1st, 1886, when corrosive sublimate as the antiseptic came into use there, up to December 31st, 1896, a period of 10½ years, and I think you will find that it thoroughly bears out what Dr. Cullingworth has said in his address, and what Stadfelt (of Copenhagen) has expressed (his opinion being emphatically endorsed by Dr. Boxall in his paper to which I have referred), the

belief that the hygiene of a maternity depends less upon its construction and its age than upon the hygienic principles upon which it is directed, and upon the perseverance with which these principles are carried out in the daily practice. For you will learn that while structural alterations failed to improve the condition at the City of London Lying-in Hospital, hygienic principles, and markedly the employment of corrosive sublimate, produced an extraordinary change for the better.

I will say a few words about the hospital itself, and my connection with it. The present building was erected in the City Road, at the corner of Old Street, in 1771, and was opened for the reception of patients on March 31st, 1773, so that it is 124 years old. There is not now, nor has there ever been, a house-surgeon; the chief resident is termed matron and midwife. There is only one medical officer in charge, who resides in the neighbourhood, and visits the hospital daily and whenever required. He is termed the surgeon-accoucheur. I took over the duties of this office in March, 1870, though I was not permanently appointed to fill it till the following July. There were two other medical officers, a consulting physician, and a consulting surgeon. These are not complimentary honorary appointments, as the term "consulting" often implies. The surgeon-accoucheur is expected to call to his aid the consulting physician whenever he is in difficulty in a delivery or in any case of serious illness, and he has the consulting surgeon to appeal to in any case that is surgical, outside midwifery and gynaecology. Both these medical officers have a seat on the Managing Committee. I resigned my appointment of surgeon-accoucheur in 1871, and in 1872 was elected from among the governors a member of the Committee of Management, the meetings of which I attended as regularly as possible up to July, 1881, when Dr. Greenhalgh resigned his office of consulting physician, and I was appointed in his place; so that it will be seen that for nearly twenty-seven years I have taken an active part in the affairs of the hospital.

In 1870, when I took over the medical charge, the patients were dying in the proportion of 1 in 19, 12 deaths occurring among the 227 women delivered, so that as soon as possible, seconded by Dr. Greenhalgh, I appealed to the Committee to close the hospital. Its doors were shut on June 15th for three months in response to our entreaty. I immediately set to work to endeavour to discover the defects and get them rectified as far as possible, and what was done is recorded in the annual report for 1870.

"Your Committee have had the building painted and coloured throughout, have done away with all the old bedsteads, and replaced them by new iron ones without curtains. Palliasses and hair mattresses have been provided in the place of the old feather beds, and all other furniture likely to retain infection has been removed; in fact, the wards have been almost entirely refurnished. Your Committee have also had all the drains opened, and double trapped at all the gratings and down spouts, and all cesspools filled with concrete. The wards have been thoroughly fumigated with chlorine gas, and every means has been used to get rid of all infection."

The result was that from the reopening on September 21st to the end of the year 103 women were confined with only 1 death, reducing the mortality for the whole year to 1 in 25. In 1873 it was 1 in 50; and in 1876 and 1877, the two years taken together, 1 in 45—so that on November 24th, 1877, the medical officers again appealed to the Committee to close the hospital. This was done on December 1st.

"It was resolved that Dr. de Chaumont, of Netley, be requested to inspect the hospital and report on the ventilation and drainage with a view to their improvement." The result of this was that great alterations were made, at an outlay of upwards of £4,500. The hospital, in its reconstructed form, was reopened on April 16th, 1879, after having been closed for sixteen months—and with what result? 287 women were delivered during the remainder of the year with only 1 death. But what a crushing disappointment came next year (1880)—12 deaths among 383 women delivered—1 in 32; and 11 of these were in the first half of the year; out of 196 deliveries, 1 in 19. Does not this bear out the opinions I have expressed in the earlier part of my address as to structural improvements? In the following year (1881) there was

again a large expenditure in this way, nearly £600—and with what result? At a meeting of the Committee held on April 28th, 1882, the medical staff reported that the sanitary condition of the hospital was very unsatisfactory, and recommended that it be closed at once. The Committee then resolved forthwith to act upon the advice of the medical officers, and appointed a subcommittee to inquire into the cause of the evil complained of. After several meetings the subcommittee presented a report to a special court of governors on June 2nd, recommending that the hospital be pulled down and rebuilt. Fancy this, after a recent expenditure of upwards of £5,000 on the building!

The special court, however, took exception to this report, and passed a resolution appointing a subcommittee from among the governors to co-operate with the Committee of Management "to report upon the present system of management adopted at the hospital." After several meetings the following report was presented:

It is recommended that the hospital be thoroughly inspected as to sanitary arrangements, and that it be at once cleansed, whitewashed, etc., and reopened as soon as possible, and that it be closed for thorough cleansing for a month every year. That the back of the hospital be pulled down, and a one-storeyed building be put up, together with other alterations at a cost not exceeding £1,000.

This was done, and the hospital was partially reopened on October 1st, after having been closed for five months. But little more than four months later, on February 17th, 1883, in consequence of the unhealthy state of the wards, the Committee, on the recommendation of the medical staff, again closed the hospital, and appointed a subcommittee to inquire into the sanitary condition of the hospital generally. Their report was to this effect:

That, as far as the construction of the hospital is concerned, no alteration is required. That certain improvements recommended by Dr. Favy, the Medical Officer of Health, be effected as regards the soiled linen shoots. That the wards be at once disinfected, cleaned, and whitewashed, and that the hospital be reopened as soon after April 1st as possible.

There were some other less important recommendations. The hospital was reopened on April 20th, and between that date and the end of the year 209 women were delivered with but 2 deaths, 1 in 104, and yet in the two years 1882-3, taken together, the mortality rose to no less than 1 in 23. In 1884-5, combined, it was 1 in 48. In 1886, up to the closure on May 14th, 3 women died out of the 133 delivered, 1 in 44; and now we come to the period when the sublimate treatment commenced.

From the reopening, under its beneficial influence, on July 1st, 1886, to September 30th, 1887, 420 confinements took place without any death. From the same date—July 1st, 1886—to December 31st, 1896 (ten and a half years), 4,608 women were delivered, with 11 deaths, mortality 1 in 419, or 2.387 per 1,000. From January 1st, 1892, to December 31st, 1896 (five years), 2,392 deliveries, with 3 deaths—1 in 797—or 1.250, that is, 1¼ per 1,000, and these deaths were:

First, in 1893, sudden death from pulmonary embolism, on the third day after delivery (patient had shown no sign of illness up to the occurrences, and the report of *post-mortem* by Dr. Tooth, Demonstrator of Morbid Pathology at St. Bartholomew's Hospital, was "embolism of the pulmonary artery; no septic condition present.")

Second death, in 1894, puerperal eclampsia, death third day after delivery.

The third death, in 1895, from puerperal eclampsia, the day after delivery.

So that we have:

1892		487	deliveries		no death
1893		430	"		1 death from pulmonary embolism
1894		492	"		1 death from eclampsia
1895		490	"		1
1896		493	"		no death.

Such a low rate of mortality as this one cannot expect to maintain, when it becomes a matter of fortune how many deaths from non-preventable causes may arise, such as the three, from eclampsia and pulmonary embolism, met with in the last five years. Only on December 26th last I was called to the hospital to a severe case of eclampsia; the patient has happily recovered, but, had she died, it would not have enabled me to say we had had no death in the year 1896, and it would have increased the rate of mortality in the last five years.

Thus it will be seen that in the last five years, among 2,392 deliveries, there had been no death in which septicæmia could have borne any part; and there had been also very little illness during this time. In the past year I have only been

asked to see one patient who had high temperature, and she was discharged quite well. I am assured that all the patients have been in a satisfactory state of health before they were allowed to leave the hospital.

A total mortality of $1\frac{1}{2}$ per 1,000 is, indeed, I think, something to be proud of, 5 in 4,000, and no death from septic causes. This seems to justify the title of Dr. Cullingworth's address—Puerperal Fever a Preventable Disease—rather different to the time of my introduction to the City of London Lying-in Hospital, when 1 in 19 was dying.

I do not dispute for one instant the great share that cleanliness plays in these results. My late colleague, Dr. Matthews Duncan, spoke of antiseptic precautions as transcendental cleanliness.³ "He had," he said, "always defended the good or the better hospitals, and he was proud to know that now, when thoroughly antiseptically managed, they needed no defence. Their mortality was equal to, if not less than, that of private practice. During his whole life he (Dr. Matthews Duncan) had watched the generous and persevering attempts of his brethren to improve midwifery hospitals. Long ago, in Dr. Rigby's time, he came to London, and saw a great ventilating scheme which was to work wonders. Architectural arrangements, drainage, ventilation, segregation, and drugs had all been tried, and all without securing success. The mortality in hospitals was terrible. He did not know what it was in private practice; probably it was as bad, or nearly as bad, as in big hospitals. Antiseptics, and antiseptics alone, had brought success, and had sustained success. It was, he knew, common in highly respected quarters in London to attribute this modern success or diminished mortality in general hospitals, as in lying-in hospitals, to many factors—to improved feeding, housing, cleanliness, nursing, and antiseptics; and no doubt all were important, but all had failed till the antiseptic era, and all would fail again were antiseptic precautions—transcendental cleanliness—neglected."

In his valuable book, *Mortality of Childbed and Maternity Hospitals*,⁴ Dr. Matthews Duncan says, "In a well-managed hospital they die at the rate of 1 in 100; all the country over the mortality is probably not much less; in the best private practice it appears to be greater." "In the years of my private practice of which I have preserved records I find 8 deaths in 736 cases, or 1 in 92." This was written in 1870, before antiseptic midwifery came in practice.

Why the sublimate treatment should not be adopted in private practice as well as in hospital I am at a loss to understand. In my lecture to the midwives that I have spoken of, given in 1887, I gave them a list of what I order to be ready in the house of every patient who expects me to attend her in her confinement, and among the things was what would make half a gallon of sublimate solution, 1 in 1,000, labelled "To be used with an equal part of hot water as a lotion"; and I have continued this, very much to my satisfaction, ever since, and I have used the irrigation just as in the hospital practice, and have never seen a sign of mercurial poisoning.

In July, 1881, I had the great misfortune to lose my first private patient, and from puerperal septicæmia. This was before the days of sublimate and sterilising instruments, but not before the introduction of other antiseptics, and every precaution had been taken by the use of carbolic lotion and carbolic cream. It was a forceps delivery; after forty hours' labour, a primipara—occipito-posterior presentation, perineum partially ruptured—became feverish after a rigor on the fourth day, and died on the thirteenth day after delivery. The most unremitting attention failed to save her life. I cannot tell you the distress the loss of this patient, a young wife, caused me. Only those of you who have had similar losses can realise what it was! I have mentioned on a former occasion a case of puerperal peritonitis in a patient delivered in December, 1890, who was attended by me on the most strict sublimate lines. It was a genuine case of acute septic peritonitis occurring a few hours after delivery, and, although no *post-mortem* examination was made, I feel convinced, from the reasons I gave you, that it was caused by rupture of a pyosalpinx, as in those cases narrated by Dr. Michie, of Nottingham, in his paper read before this Society in May, 1885.⁵ If this case be excepted, no case of septicæmia has occurred in my private practice since I commenced the use of corrosive sublimate.

It gives me a feeling of protection I cannot over-estimate. I feel sure if its employment in midwifery were generally adopted by practitioners for their hands and for vaginal irrigation, the Registrar-General's reports of the mortality from puerperal septicæmia would very soon show a difference, and in these days of portable compressed drugs there is no reason why it should not be almost universally used, even among the very poor; but it is only right to point out what may result if it be improperly used.

In the annual report of the General Lying-in Hospital for 1883, the medical report, signed by Drs. (now Sir John) Williams and Champneys, says that 342 women were confined, of whom 3 died (1 in 114), 2 from septicæmia, quite independent of one another, "a percentage which testifies to the healthy condition of the hospital." This was during the time that antiseptics were employed there, but, you will note, before the corrosive sublimate was commenced, and the Committee add in presenting this medical report to the governors: "The Committee beg to tender their congratulations to the eminent surgeon to the hospital, Sir Joseph Lister, Bart., to whom this and other institutions are so much indebted for the satisfactory medical results obtained through the system with which his name is associated."

Sir William Priestley concluded his paper read before the Obstetrical Society of London, July, 1885, entitled Notes of a Visit to Some of the Lying-in Hospitals in the North of Europe; and Particularly on the Advantages of the Antiseptic System in Obstetric Practice, with these words: "Ample funds and sufficient space are necessarily most potent aids in waging warfare with disease in hospitals, but, even with limited means and space, absolute cleanliness, supplemented by the antiseptic system, may in our own country have a large influence for good, and the labours of our illustrious Lister prove of equal value to the obstetrician as to the surgeon. If I am not mistaken some of our Fellows," says Sir William Priestley, "can tell us of triumphs of this kind in the lying-in hospitals of London." I hope you will think that I have this evening succeeded in doing so.

I feel I cannot conclude my address better than by saying that though numerous peerages have been conferred upon those "waging warfare," resulting in destruction of life and victory, never has one been more worthily bestowed than on "our illustrious Lister," for the victory gained by him in "waging warfare with disease in hospitals," resulting in such an enormous saving of life among our fellow creatures.

REFERENCES.

¹ A vaginal douche of 1 in 2,000, at a temperature of 115° , is given immediately after the delivery of the placenta, and three similar douches subsequently at intervals of night and morning, then iodine douches are given instead of sublimate. ² Delivered at St. Thomas's Hospital, London, and published by J. and A. Churchill, 1888. ³ *Obstetrical Society's Transactions*, vol. xxvii, p. 215. Discussion on Dr. (now Sir William) Priestley's paper, Antiseptics in Lying-in Hospitals. ⁴ Edinburgh: Adam and Charles Black, 1870, p. 116. ⁵ Pregnancy complicated by Suppuration in the Pelvis, *Brit. Gynaecol. Journal*, 1895, p. 160.

TWO CASES OF IDIOPATHIC ARTERIAL HYPERMYOTROPY;

WITH REMARKS ON THE SYMPTOMS, DIAGNOSIS, AND TREATMENT OF THE AFFECTION.

By THOMAS D. SAVILL, M.D. LOND., M.R.C.P.,

Physician to the West End Hospital for Diseases of the Nervous System; Examiner in Clinical Medicine in the University of Glasgow; formerly Medical Superintendent of the Paddington Infirmary.

By "arterial hypermyotropy" is meant a disease in which the leading anatomical feature is a hypertrophic increase in the middle or muscular coat of the smaller and medium-sized arteries throughout the body in varying degrees—those vessels, namely, in which the muscular tissue is normally most plentiful. The condition has for a long time been recognised—chiefly in the arterioles¹ only—in association with chronic renal affections; but it had not, so far as I am aware, been described as an idiopathic or separate affection, nor as one to which distinctive symptoms belonged, until 1891, when I presented a preliminary communication to the

¹ Dr. W. H. Dickinson has recognised a muscular thickening in the arteries of all sizes, large as well as small, in connection with chronic renal disease—The Baillie Lectures, *Lancet*, 1895, i, 137.