castration in the operative treatment of enlarged prostate, but I am by no means converted. The condition of some of my patients years after prostatectomy is such as to make me feel that in the operation suggested by my lamented colleague McGill and by Belfield, independently, but almost simultaneously, some years ago, we have to chronicle a real advance in the treatment of enlarged prostate which has failed to be relieved by catheterism and other milder measures. The details I gave in a paper at the International Medical Congress at Rome still holds good in my practice, both with regard to the preliminary preparation of the patient, the steps of the operation, and the subsequent treatment during convalescence.

A CASE OF AMPUTATION IN MIDDLE AGE FOR ALBUMINURIA, FOLLOWING TUBERCULOUS KNEE: RECOVERY.

By J. J. H. HOLT, M.B., CH.B. (VICTORIA),
Buckley, Chester.

THE question of operation in cases of albuminuria is one often difficult to decide, more especially when it occurs in patients who are advancing in years. The following report may therefore be of some interest and use in helping to arrive at a decision in some doubtful cases.

J. C., a business man, aged 52, bachelor, was first seen by me in May, 1896. He had been suffering from a deformed knee since childhood, but called me in because he was feeling generally unwell, and complained of obscure lumbar pains and restlessness at night. He had been passing a considerable amount of urine, which was acid, light in colour, specific gravity about 1012, and was distinctly albuminous. Some finely granular casts were seen under the microscope. Physical examination gave no sign of enlarged liver or spleen. The heart and lungs were normal. The left leg was flexed at right angles, and the knee greatly deformed, much swollen, and inflamed. On the outer side there was a large granulating surface, in the centre of which was a sinus leading down to the knee-joint. It was discharging a thin serous-like pus, and was very tender. It was injured when he was 11 years of age, and had been discharging since he was 12. The family history evidenced a tuberculous tendency, two of his sisters having died of consumption at about 27 years of age.

The question of operation had been considered when he was 12, but had been decided against. In 1891 he was in bed twelve weeks with an attack of rheumatic fever. He made a good recovery and enjoyed fairly good health until August, 1892, when the knee became very painful and discharged more freely. It improved in about four months, and he was again comparatively healthy until about May, 1896. He then began to be troubled with pain in the loins and could not rest well at night. Hoping that the condition might be temporary and that possibly he might improve by resting, he was advised to take a holiday and use boracic dressings for the knee. He went away for a month, and felt better in health whilst away and for some days after his return. Subsequent examination of the urine, however, gave even more distinct evidence of albumen than it did a month previously. Regarding this, therefore, as an indication that the kidneys were becoming secondarily affected, he was advised to consult Mr. W. Whitehead, of Manchester, as to the advisability of amputation. He was strongly advised by that gentleman to undergo the operation, whilst at the same time he was informed of its risk. Declining health and increasing pain, however, made him decide to have it done.

Amputation was performed by Mr. Whitehead on August 19th. Owing to the unhealthy nature of the tissues, it was found necessary to amputate between the upper and middle third of the thigh. Notwithstanding the increased risk from the high operation the patient rallied well and made an excellent recovery. He was kept on a milk diet and simple aperients given to regulate the bowels. He was able to leave his bed early in September, and has progressed most satisfactorily ever since. The albuminuria steadily diminished and has now disappeared. He has resumed work, is sleeping

well, is free from pain, and enjoys better health than he has done for years.

The case is interesting in showing the possibility of relieving secondary albuminuria even after long-standing suppuration. In this case the discharge had continued for forty years, but was followed by recovery, even after such a dangerous operation.

A NOTE ON A FEW SURGICAL USES OF PICRIC ACID.

By C. M. ALLAN, M.A., M.B., C.M.,
Surgeon, Longton Cottage Hospital, etc.

IN the BRITISH MEDICAL JOURNAL of December 26th, 1896, Dr. Maclellan drew attention to some uses of picric acid as a protective and healing agent in skin diseases.

When experimenting with various therapeutic agents, in a study—extending over fourteen years—of varicose ulceration of the leg and allied diseases, among others (including sulphurous, nitric, tannic, and oxalic acids and chlorine) I have used picric acid (a concentrated solution in rectified spirit diluted with two parts of distilled water, and applied with a spray), and found it the most successful in allaying the irritation and stopping the discharge, and irritation from the discharge, from eczematous and ulcerated surfaces.

In the case of ulceration or eczema, or of ulceration and eczema of the leg, the leg is, subsequent to the application of the acid, covered with a medicated adhesive plaster. The smarting and itching resulting from the disease and exposure to the air are immediately allayed, the inflammatory redness and irritation subdued, the parts kept aseptic or rendered so, and the fine coating of coagulated albumen and "epithelial debris" (which disappears before the application of the next dressing in a week) exerts an astringent and slightly compressive action on the dilated venous and lymph radicles in both ulcer and surrounding inflamed surfaces, and healing takes place rapidly, and with comfort to the patient.

In recent wounds, with or without great loss of substance, this treatment is equally beneficial, and is not so troublesome nor so unsuccessful as skin grafting, where the size of the wound would render that advisable. I have under my care a wound of the forearm with loss of skin which a hand could not cover, healing rapidly under this combined treatment after the Thiersch method had failed. In that most dangerous and most intractable form of hæmorrhage from wounds in hæmophilic subjects picric acid solution acts better than any form of treatment I have tried. Within the last few years I have had two such cases, one of which belonged to a distinctly hæmophilic family. The first, a cut several inches long, was sprayed with the acid while the blood was being mopped off with wool; and treated thus several times daily, did better than with iron and other astringents. The second, an abscess of the glands of the neck after scarlet fever, which necessitated incision, was packed with picricised gauze and did well.

In cases of severe cellulitis arising from the introduction of some irritant poison, and when incision or amputation is necessary, the bleeding from the paralysed and dilated radicles becomes almost uncontrollable. If the incision is packed with gauze wrung out of the picric solution the bleeding is soon controlled. For the oozing after the amputation the spray is much more successful than hot water, yet a combination of both may be used with great advantage, the water being used first (as it would decompose the acid if both were combined) and the spray applied immediately after. The acid acts as an astringent and antiseptic hæmostatic, and the parts heal *cito, tuto, et jucunde*.

OUR Correspondent in Hong Kong writes: The threatened outbreak of small-pox in Hong Kong has been successfully met by the vigorous measures adopted by the Government to stamp out the disease, as only an occasional case is now being reported. The number of cases since the beginning of December has been 25, 19 being in December, and 6 during the first thirteen days of January; and the number of deaths has been 11—7 in December and 4 in January to date. Ten of the cases and three deaths were in Europeans.