OBSERVATIONS

ON THE

ANTICIPATION OF POST-PARTUM HÆMORRHAGE,

WITH

REMARKS ON THE ACTION OF ERGOT
ON PREGNANT WOMEN.

BY LOMBE ATTHILL, M.D.,

Ex-Master of the Rotunda Hospital, Dublin.

A DISCUSSION on the subject of the Anticipation of *Post-partum* Hæmorrhage took place at the annual meeting of the British Medical Association held in London in 1873, after the reading of a paper on the subject by Dr. Ewing Whittle, and, in the number of the BRITISH MEDICAL JOURNAL for November 1st, 1873, a paper of mine appeared, in which I say: "Dr. Whittle's paper, though read just before the hour at which the Section adjourned, was listened to with great interest, and a short and animated discussion followed. The reason why this paper was listened to with such attention is obvious. Every practitioner present, from personal experience, dreaded *post-partum* hæmorrhage, and, while admitting the value of transfusion in extreme cases, rejoiced at the prospect of not being obliged to think of so difficult and uncertain a procedure; indeed, more than one member (during the debate which had previously taken place on transfusion) asked—Could no means be suggested for rendering this operation unnecessary? Dr. Whittle's paper professed to be a reply to this query, but I think the subject should be more fully discussed." This I then proceeded to do, and my example was followed by my lamented friend, the late Dr. Alfred McClintock, in an admirable paper read before the Obstetrical Society of Dublin, which will be found in the volume of the *Proceedings* of the Dublin Obstetrical Society for the session 1873-74.

I have no intention of wearying the reader with a summary of these papers. Anyone interested in the subject will find them, and a report of the discussion which followed, in the volume I have referred to; but the question then discussed was, how best to avert the anticipated danger in cases where labour was actually in progress, and though many of the suggestions were valuable, they too often fail to be efficient, because time does not exist to permit preventive treatment to take effect. Thus great stress was laid on the administration of ergot, but I have long since been convinced that, to be of real use, ergot requires not only to be administered some hours before its action as a uterine hæmostatic is needed, but also that the dose must be repeated, and that more than once, at intervals of about two hours, otherwise the effect is uncertain, and so transient, as to be of little value. I do not, therefore, propose to discuss this part of the subject, but to point out that in patients in whom a tendency to dangerous *post-partum* hæmorrhage is known to exist, or indeed is suspected to exist, the danger may in the majority of cases be averted, or at least greatly lessened, by treatment carried out antecedent to the expected date of the occurrence of labour.

I had not been long in practice before I became convinced that the theory then universally believed, that women should not take any drug, especially tonics such as quinine, iron, and the mineral acids, during the menstrual period was erroneous, and I took every possible opportunity of watching the effect of such medicines on women during menstruation, with the result I expected, namely, that no drug taken in its usual dose had any effect on the function in healthy women. This fact led me to administer drugs during pregnancy when necessary with equal freedom; and in the case of a lady whom I knew to have a tendency to *post-partum* hæmorrhage, I on one occasion administered quinine in 4 or 5-gr. doses up to the advent of labour for the cure of neuralgia, and on this occasion for the first time no hæmorrhage occurred with her. This struck me as remarkable, and it occurred to me that the unexpected freedom from hæmorrhage might have been due to the tonic effect of the quinine on the muscular fibres of the uterus; and, emboldened by the result in this case, I administered quinine to other patients in whose cases I feared the occurrence of hæmorrhage.

About this time the result of the action of ergot in a case of threatened abortion, to which I shall presently refer, induced me to prescribe ergot combined with strychnine for a patient, in whom from previous experience I feared *post-partum* hæmorrhage, for some days before labour set in, and I was gratified by finding that although there was a little hæmorrhage, it was as nothing compared with what had occurred in her previous confinements. From that time it has been my invariable rule to carry out this practice in all cases in which I had reason to fear the occurrence of hæmorrhage, and the results have been most satisfactory. I never once have had occasion to regret its adoption, and never observed the most trifling injurious effects on either mother or child to follow, and this although the ergot has been taken regularly for five or six weeks prior to the occurrence of labour. The following cases are typical ones, and very instructive.

About six years ago Dr. Wright, of Dalkey, sent for me to assist him in a dangerous case of hæmorrhage. More than two hours elapsed between his sending for me and my arrival, and before I did so his exertions had been successful. The hæmorrhage had stopped and the patient's life saved, but she was still in a very precarious condition; and Dr. Wright informed me that her previous confinements had also been attended with alarming hæmorrhage.

About eighteen months subsequently the lady, now residing in Dublin, requested me to attend her in her confinement which was again approaching, and I reluctantly consented to do so, on the condition that she would submit to a regular course of treatment prior to the expected date of her confinement. She had a great objection to medicine, and was unwilling to agree to this, but on explaining to her the *rationale* of the treatment she consented, and as soon as she had entered on the thirty-fifth week of utero-gestation I put her on the following mixture: Inf. ergotæ ad $\frac{3}{4}$ vj; ext. ergotæ liq., $\frac{3}{4}$ ij; liq. strychninæ, $\frac{3}{4}$ j; a tablespoonful to be taken three times a day. This mixture she continued to take regularly for three weeks. I then discontinued it for five days, when I directed it to be recommenced.

This lady had in all her previous confinements (five in number) been taken with labour six or seven days before the expiration of the normal term of 280 days from the cessation of her last menstrual period, but on this occasion labour did not set in till the 288th day, a delay which caused her some uneasiness, but as I had before then remarked that in the case of patients under the influence of ergot the term of gestation was likely to be prolonged, I considered it a good sign, and my prognosis was verified. Labour was in all respects normal; the first stage occupied about five hours, a longer period than usual with her; the second an hour and a half; the placenta was expelled in about twenty minutes, and there was not a wineglassful of blood lost; in fact, I think it was the most bloodless labour I ever witnessed. This lady was again confined eighteen months later; the same treatment was adopted. Labour did not set in till the 285th day; there was no hæmorrhage.

Very recently a similar case occurred to me. A lady of rank wished me to attend her in her confinement. This I refused to do, especially as she resided some miles from town, but I promised her if she arranged with a medical man to sleep at her house I would go to her if any cause of anxiety appeared. She was the mother of ten children, and in her case also labour usually set in at about a week before the expiration of the normal term of pregnancy. Her two last confinements had been followed by alarming hæmorrhages, and her husband was very anxious about her. I treated her in exactly the same way as the last case. On this occasion, instead of labour setting in too soon, it did not occur till the 295th day from the date of her last menstrual period; and Dr. Glenn, who attended her, can verify my statement that there was no hæmorrhage to speak of, that the child was vigorous, and that neither mother or child exhibited any ill effects, though ergot and strychnine had been taken with short intermissions for nearly seven weeks.

I might easily quote other cases; but these are sufficient to prove—at least, to my satisfaction: (1) That ergot, alone or in combination with strychnine, may be taken with absolute safety to both mother and child by pregnant women, in the usual doses, and for a considerable time. (2) That when

taken continuously for not less than three weeks prior to labour, it tends to delay the setting in of uterine action, and I may add that in no case have I ever seen it bring on labour in a case in which uterine action had not already been excited. (3) That its exhibition for some weeks prior to the commencement of labour arrests the tendency to *post-partum* hæmorrhage, and facilitates the involution of the uterus.

When I was a student we were cautioned not to administer ergot to women in labour till the os was fully dilated, it having been shown that, unless the child was born within the space of two hours after the specific action of ergot was manifested, that is by the occurrence of short, sharp pains with hardly any appreciable interval, the child was likely to suffer, and was not infrequently born dead. This statement I can verify as being perfectly true, but then the death of the child was not due to any poisonous property possessed by the ergot, but to the utero-placental circulation being interfered with in consequence of the rapid, almost incessant, contractions of the muscular fibres of the uterus. That this is so is proved by the fact that where the ergot failed to induce its specific effect the child did not suffer. True it is, that, as stated by McClintock, some children were born dead even when ergot pains had not been induced, but remember he wrote at a period when the forceps were very rarely used. During the six months I spent as a pupil at the Rotunda, the forceps were never, so far as I know, once applied. Labour was allowed to linger on sometimes for days, and when the fetal heart ceased to be audible the head was lessened and the woman delivered. Now, perhaps, we run to the other extreme, and apply the forceps too frequently, but that a child should be stillborn whether ergot was given or not, when labour was permitted to linger on for days, is not to be wondered at; that the death of the foetus was, however, due to the administration of ergot, when the true ergot pains did not occur, I disbelieve, and this conclusion I arrived at, while still a pupil, and its truth I am now convinced of.

The routine of giving a dose of ergot, when *post-partum* hæmorrhage has set in, is generally a harmless, and sometimes a useful practice. In perhaps one in twenty cases it may induce the uterus to contract, but to be of any permanent use, the drug should be administered some hours previous to the occurrence of the hæmorrhage, and the dose should have been repeated at intervals of a couple of hours or less. If then we meet with a patient in labour in whom a tendency to *post-partum* hæmorrhage has been previously observed, ergot should be administered at once, and if delivery does not take place for a few hours, and if the dose be repeated, the result is likely to be satisfactory.

It is to be borne in mind that I am not discussing the methods by which *post-partum* or any other form of hæmorrhage occurring unexpectedly during parturition should be treated, but that I confine myself to advocating the value of the exhibition of ergot prior to the end of the period of utero-gestation in a certain class of cases, and I am far from underestimating the advantage of rupturing the membranes, the use of friction over or compression of the uterus, cold externally, or hot-water douches when necessity arises, but I wish, if possible, to prevent the necessity for such.

Although foreign to this subject, I wish to state my experience of the use of ergot in cases of threatened abortion; in these hæmorrhage without pain is often the first symptom; and that indicates that the ovum is partially detached; when wholly detached hæmorrhage as a rule ceases and the contents of the uterus are sooner or latter expelled. But it is seldom that we can say with certainty when it has become hopeless to save the ovum, and therefore it is our duty to persevere in our attempts to do so as long as possible; and of all methods to this end, absolute rest in bed for a considerable time is without doubt the best. But it is most irksome to a patient, specially a mother with probably young children needing her care, to submit to the restraint imposed on her by keeping her bed for weeks, and many will refuse to do so; indeed, some women become so prone to abort that it seems useless to try to enforce prolonged confinement to their room, and then it is obviously better that the ovum should be quickly got rid of.

Such a case occurred to me some five and twenty years ago. About that time I had as a patient a delicate young married lady, who in the course of the preceding ten months had

twice aborted at about the tenth week of her pregnancy, and on each occasion alarming hæmorrhage had occurred. I attended her on the second of these. She became pregnant again within two months, and at the expiration of almost exactly the same time as on the previous occasions, hæmorrhage set in, which she knew was the forerunner of another miscarriage. I saw her a few hours later. There were no pains, but the os was patulous, and her state identical with what it had been in the early stage of her previous illness. I came to the conclusion that she must certainly abort, and I at once put her on ergot and strychnine, with the view of getting rid of the ovum rapidly. The dose of ergot was repeated every two hours, and I watched this lady all day, hourly expecting that she would have a recurrence of the hæmorrhage which had so alarmed her and me on the previous occasion. But instead of this the hæmorrhage lessened; night came on, and she slept. Next day the os was not the size of a split pea. This lady went to the full term and gave birth, to her great joy, to a son, now a strong man 6 feet high.

From that day on I invariably administered ergot to women threatened with abortion. In some it produced no effect whatever; in a few it induced uterine action, and the expulsion of the ovum followed. In the majority the threatening symptoms disappeared, and pregnancy proceeded normally; but in not one of them did I regret having administered ergot, and I am satisfied that if the ovum is not blighted, that is, ceased to be a living body, that ergot acts as a uterine tonic, and renders the organ in many cases fitted to undergo the further changes which take place in it during utero-gestation, but if the ovum is detached and blighted, then it becomes, as it were, a foreign body, and ergot is then likely to stimulate the uterus, and to expel its contents. This opinion is, of course, based only on the results of my personal observation, but of its correctness I have no doubt.

CONCLUSIONS.

To sum up the whole subject as to the action of ergot in combination with strychnine, which is the formula I usually adopt, I venture to lay down the following as the result of my experience:

1. That when administered previous to the termination of pregnancy in the case of women in whom a tendency to *post-partum* hæmorrhage is known to exist, it tends in a marked manner to prevent the occurrence of hæmorrhage.
2. That when so administered in ordinary doses, it does not produce any injurious effect on either mother or child, and that its exhibition seems to delay the commencement of labour in such case.
3. It tends to make the involution of the uterus more perfect, and lessens the chance of the occurrence of subsequent uterine troubles, many of which depend for their cause on imperfect involution of that organ.
4. It will not bring on premature labour or induce abortion unless uterine action has previously been set going.
5. In cases of threatened abortion its administration frequently seems to act as a uterine tonic, and in some cases tends to avert the danger of a miscarriage, provided the ovum be not blighted.
6. That if the ovum be blighted, and specially if it be detached, ergot usually hastens its expulsion.

Since writing the foregoing, I observe in the *EPITOME* which appears in the *BRITISH MEDICAL JOURNAL* of January 16th, an abstract of a communication made by Professor Schwab on the efficacy of quinine as an oxytocic. This drug, he maintains, "stimulates the uterine fibres when once they have begun to contract of their own accord. Like ergot, it does not set contractions going." I have never administered quinine as an oxytocic during labour, but it will be remembered that I have stated at the commencement of this paper that it was the unexpected immunity from *post-partum* hæmorrhage in a patient to whom I had administered quinine prior to labour, that induced me to carry out the preventive treatment I now advocate, and that it was quinine I first administered for this purpose. My observations, therefore, tend to confirm the views enunciated by Schwab as to the action of quinine on the uterine fibres, and further observations of the action of quinine as an oxytocic should be made. I do not think, however, it would prove as reliable a drug for

this purpose as ergot is. Schwab, too, confirms the opinion I have long held, that ergot does not originate uterine action.

A CASE OF SUPPURATIVE OTITIS MEDIA COMPLICATED WITH CEREBELLAR ABSCESS.¹

By H. SECKER WALKER, F.R.C.S.,

Assistant Ophthalmic and Aural Surgeon, Leeds General Infirmary;
Lecturer on Ophthalmology and Otology at the Yorkshire
College, Victoria University.

On November 22nd, 1895, I saw, in consultation with Mr. J. W. Hutton, C. J., aged 14. He had had right otorrhœa for some years with a polypus in this ear. I had removed the polypus two years earlier, and the otorrhœa ceased. Four months before the date on which I saw him again he was attacked by pain in the left ear, which was followed by a purulent discharge, and lately this had been tinged with blood; the pain had returned and spread over the head and become constant, being located chiefly in the occipital region. The boy looked pale and had wasted rapidly. Five weeks earlier he had several attacks of shivering. He had vomited occasionally in the morning, constipation had been marked, and there had been some delirium at times.

On examination of the left ear a polypus was seen deep in the meatus, the discharge was very offensive, the mastoid region looked healthy, but there was tenderness on deep pressure. He was very deaf in both ears. The cervical glands were enlarged, and one or two superficial ones in front of the neck had ulcerated. The temperature had been between 99° and 100°, the pulse from 110 to 130. There was no optic neuritis, no paralysis, no spasm of any of the ocular or facial muscles. The grasp of the hands was weak, but not markedly more on one side than the other. The knee-jerk was increased on the left, normal on the right side. When spoken to he answered intelligently, and his answers were not delayed. In the presence of the above symptoms there could be no doubt about the condition of the mastoid process, but I did not then feel justified in diagnosing intracranial disease, as most of the above symptoms may be present in a simple case of disease of the mastoid cells.

The following operation was performed:

FIRST OPERATION.

A curved incision, $2\frac{1}{2}$ inches long, immediately behind the attachment of the ear was made down to the bone, the cartilaginous meatus stripped from its bony attachment and the ear turned forwards, the periosteum was pushed backwards, so that a considerable area of the temporal bone, including the external auditory meatus and the greater part of the mastoid process, was exposed together with the surface landmarks which are of such great assistance in this operation. Immediately behind and slightly above the centre of the external auditory meatus lies the mastoid antrum. The bone covering this was of a dirty grey colour. The chisel easily broke this down and entered the antrum, which was filled with yellowish putty-like stuff with a very offensive odour. After clearing out this the posterior wall of the meatus and the external wall of the attic were chiselled away, and the carious ossicles and remains of the membrana tympani and polypus in the middle ear removed so that the middle ear and antrum formed one cavity partly divided into two by a transverse ridge of bone which contains the facial nerve. It is important that this ridge should not be chiselled away, otherwise the facial nerve will be divided, resulting in more or less permanent facial paralysis on this side. This accident will be avoided by not cutting away bone below the upper third of the posterior meatal wall.

The roof and posterior wall of the antrum were found carious and perforated so they were removed, exposing the dura mater lying against the temporo-sphenoidal convolutions and the lateral lobe of the cerebellum. The dura mater in each of these situations was firm, and, as I thought, probably limited the disease. After well washing out the cavity it was packed with iodoform gauze, and the superficial glandular abscess in the lower part of the neck was scraped.

For the next ten days the boy did well; he became brighter, the headache and vomiting ceased, the temperature fell a little, becoming subnormal, the pulse averaged 112, and he was only troubled by obstinate constipation. On the eleventh and twelfth days vomiting and headache again occurred; there was still no optic neuritis. The condition of the arm muscles and knee reflexes remained the same, but the boy was more prostrate. These symptoms continued; it appeared as if there must be some intracranial disease, and the symptoms pointed to cerebral or cerebellar abscess. From the amount of disease in the posterior and lower part of the

mastoid process lying near the lateral lobe of the cerebellum and the occipital headache, I thought this was the more likely place in the absence of any guiding symptoms. Dr. Churton and Mr. Teale, who kindly saw the case, agreed with me as to the probability of intracranial abscess, and were also in favour of the cerebellum being the seat of the disease.

SECOND OPERATION.

The opening into the posterior fossa of the skull made at the first operation was enlarged, the dura mater reflected, and a silver drain tube passed into the cerebellum for an inch. On the second trial about 3ij of very offensive pus welled slowly out of the tube. An unsatisfactory attempt was made to wash out the abscess cavity, two drain tubes were left in, and dressings applied. This was a fortnight after the first operation.

The patient improved and the headache ceased, but the vomiting still occurred occasionally. Two days later the symptoms all reappeared; the tube was pushed in a little further, more pus came away, and again an attempt was made to wash out the cavity.

The condition of the boy now underwent much improvement, and next morning he was sitting up in bed calling for his breakfast, and much interested in the *Yorkshire Post*. For three weeks the improvement was maintained. On December 30th he had several attacks of vomiting; the wound looked well and there was practically no discharge from the tube. Thinking that the presence of the drainage tube, acting as a foreign body, might be causing the vomiting, I removed it. For a fortnight again he did very well, and then all the old symptoms returned with redoubled force; in a few days he changed from being a comparatively healthy-looking boy to a mere skeleton.

THIRD OPERATION.

On January 3rd I again opened the cerebellum, and this time a little further back. Pus was found without difficulty, and two decalcified bone tubes were introduced and carbolic lotion syringed gently through them.

From this time the boy made a rapid and complete recovery, the only drawback being that now for the first time with all this disease and surgical interference he developed an attack of double optic neuritis, which, however, disappeared in a few weeks without permanently damaging the vision.

The patient now has a cavity of considerable size in the left mastoid process, lined throughout with epidermis, and communicating with the external air by an opening behind the ear and by the external auditory meatus.

REMARKS.

Recovery from cerebellar abscess is exceptional. As far as I have been able to learn there are only 10 previous cases on record. The difficulties in diagnosis, the absence of any special localising symptoms, or even in some cases of any symptoms at all, and the liability to a sudden fatal termination of the case before treatment can be carried out, are probably among the chief reasons why so few cases recover.

In reporting a successful case of cerebellar abscess, Dr. Acland and Mr. Ballance² have drawn attention to the occasional presence of certain paralytic symptoms which may be of great service in localising a cerebellar abscess, namely, muscular weakness of the limbs of the same side as the ear disease, affecting chiefly the arm, conjugate deviation of the eyes to the opposite side from weakness of the muscles, which draw the eye to the same side, and increased knee-jerk on the same side as the otorrhœa. That these symptoms have been present in some previously recorded and unsuccessful cases of cerebellar abscess a study of this interesting paper will prove, but their localising value had evidently not been understood; and, indeed, in one case the symptoms actually led the surgeon into the error of operating on the opposite and healthy side of the brain.

² *St. Thomas's Hospital Reports*, vol. xxiii.

MEDICAL STUDENTS AT AUSTRIAN UNIVERSITIES.—The following figures show the number of students at each of the universities of Austria during the current semester;—Vienna, 6,104; Graz, 1667; Prague (German), 1,391; Prague (Czech), 2,787; Innsbruck, 1004; Lemberg, 1,556; Cracow, 1,296; Czer-nowitz, 391. The total number of students belonging to the medical faculties of all these universities combined is 4,728, which is rather less than in the corresponding period of last year.