one has already said, you will see that I consider diuretics contra-indicated in the acute stage—yet I must observe that in our case there was from the first a large amount of albumen present, and that this has not apparently proved prejudicial to the use of the drug.

It is obviously unsafe to draw conclusions from one case, and it is no part of my purpose to-day, gentlemen, to attempt to reconcile these different statements. It must be a sufficient satisfaction to find that in the case before us great relief was obtained after we had used diuretin in place of other remedies, and that this relief of urgent symptoms coincided with the excretion of an increased quantity of fluid. In the early days of our employment of diuretin the asthmatic attacks first yielded, and afterwards we noticed that the amount of dropsy became considerably less. That the benefits obtained from this mode of treatment were not mere coincidences is shown by a slight return of asthma when, the patient having become accustomed to the use of the remedy, the amount of fluid excreted again became reduced; and you will have seen how the asthmatic seizures passed off at first with an increase in the amount of diuretin administered, and subsequently after the employment of digitalis in addition to the diuretin. I do not consider that the treatment is likely to produce any lasting beneficial effect. In all probability we shall have, sooner or later, to relieve the pleura again by aspiration, and perhaps to relieve the tension in the extremities by acupuncture or by Southey's capillary tubes. But these measures we endeavour to postpone as late as possible, since frequently the strength fails rapidly after draining large quantities of fluid from the extremities, and there is always some risk of exciting inflammatory changes in the neighbourhood of the punctures.

LECTURES

ON

DISEASES OF THE GALL BLADDER AND BILE DUCTS.

Delivered at the Royal College of Surgeons of England.

By A. W. MAYO ROBSON, F.R.C.S.,

Hunterian Professor of Surgery and Pathology, Royal College of Surgeons of England.

LECTURE II.

FISTULA.

FISTULÆ in connection with the bile passages are by no means uncommon, and their variety is considerable. They result from operation, or from disease, and in the latter case they are due to ulceration resulting from gall stones or cancer. The fistulous channel may either be direct or indirect. In the latter case the perforation occurs first into an adjoining parenchymatous organ or into a localised abscess, and then into the nearest hollow viscus or on to the surface of the body at some part. A fistula may also arise from a local abscess forming outside the biliary passages around the primary focus of inflammation, and then bursting into the adjoining cavities, which are thus made to communicate.

Although the establishment of a fistula is at times dangerous, and at others excessively annoying or uncomfortable, in many cases it forms one of Nature's methods of cure in certain forms of disease of the gall bladder or bile ducts, and the surgeon in forming a permanent biliary fistula in otherwise incurable jaundice, or in making an anastomosis between the bile passages and the intestine for the like purpose, is taking a leaf from Nature's book.

Many of the fistulæ are mere pathological curiosities, quite undiagnosable, and capable only of being discovered *post mortem*. Many must form and heal, leaving the patient cured, and thus never be discovered; for, contrary to what one might suppose, fistulæ between the bile passages and other hollow viscera in the majority of cases heal spontaneously, leaving only visceral adhesions, so that the fistulæ are comparatively rarely found *post mortem*.

Courvoisier and Naunyn¹ tabulate 384 of these fistulæ; Roth, Schroeder, and Schloth record 43 in 10,866 necropsies. My operations extend to 170 cases, in 12 of which there were

fistulæ following. Several of the fistulæ were inevitable, as the ducts were strictured; in others they were intentional, as in cancerous obstruction, producing jaundice.

Biliary Cutaneous Fistula.—Courvoisier's statistics gathered from reported cases would seem to prove that this is the commonest form of fistula. It may be pathological or post-operative. Pathological surface fistulæ usually open at the umbilicus, the abscess following the course of the remains of the umbilical vein, but they may form at any part of the abdominal wall, even near the pubes or on the left side of the abdomen. Calculi of various sizes—from a single one 3 inches in diameter, reported by Gutteridge,² to multiple small faceted ones the size of shotcorns—may be discharged in this way, leading to recovery and permanent cure, but until all the calculi are discharged the fistula is liable to remain open.

Post-operative fistulæ may be mucous or biliary. Mucous fistulæ are occasionally seen after the operation of cholecystotomy, where the obstruction in the cystic duct has not been overcome or where that duct is the seat of stricture. A mucous fistula, as a matter of fact, causes very little inconvenience, as only about 1 ounce of fluid is discharged daily; but if the opening be allowed to close, the accumulation produces pain, and it is therefore necessary for a patient under these circumstances either to wear a small tube and a pad of absorbent wool or to submit to operation; where there is stricture of the cystic duct the gall bladder should be removed, in other cases clearing out stones may suffice. The operation of cholecystotomy will not be followed by fistula if the bile ducts have been cleared and if the opening in the gall bladder be sutured to the aponeurosis and not to the skin.

Biliary fistula following on operation is quite a different matter from mucous fistula, as although in some cases it is compatible with good health, the inconvenience caused by 30 ounces of bile flowing from the fistula daily produces so much discomfort that in all the cases which have come under my notice the patients have preferred to accept the risks of operation rather than to retain their disability. If the ducts be clear and the fistula be small the application of the actual cautery to the margin of the fistula will frequently result in its closure. That failing, the abdomen may be opened, the gall bladder detached, and the opening sutured. Or the less severe method may be first tried of dissecting the fistula with its peritoneal covering intact away from the skin margin, afterwards doubling in the mucous edges, suturing them accurately, and over this applying one or two layers of buried sutures before bringing together the skin. Where, however, the ducts cannot be cleared, and the gall bladder is large enough to permit of it, the operation of cholecystenterostomy may be performed. If the fistula be dependent on gall stones or fragments in the ducts, the ducts may be syringed through a catheter daily with olive oil or with a 0.5 per cent. solution of *sapo animalis*, as recommended by Dr. Brockbank; or a solution of turpentine in ether may be used.

Biliary intestinal fistulæ, as might be expected from the contiguity of the gall bladder to the duodenum and colon, are the most common, and as a rule they are due to ulceration produced by gall stones. Usually the ulceration proceeds quietly and produces very few symptoms until the gall stone sets up obstruction in its passage down the intestinal canal. But the process is not always accomplished without symptoms, such as pain over the liver, more or less jaundice, and fever of an irregular type, to be subsequently followed by signs of more or less complete obstruction; the frequency of the latter event varies with the height of the fistulous opening into the bowel. Hæmorrhage into the stomach or intestines may occur in these cases.

When a gall stone is impacted in the common duct just before entering the duodenum, ulceration and perforation of the duct are apt to occur, the concretion thus escaping into the duodenum by an enlargement of the ostium of the common bile duct from ulceration or sloughing. Roth, who has paid special attention to this condition, found it 5 times in 25 cases of biliary fistula. These gall stones are usually smaller than those causing gall bladder intestinal fistula, are seldom larger than filberts, and do not often cause intestinal trouble.

I have only one case to advance from my own experience, and in that instance the vomiting of gall stones made the

diagnosis probable, though the complete recovery of the patient, a woman of 50, rendered it impossible to be absolutely certain that the surmise was correct.

INTESTINAL OBSTRUCTION.

Intestinal obstruction from gall stones is such a distinct complication of cholelithiasis calling for special treatment that I think it will not be beyond my province to consider it; and as the chief variety of obstruction is necessarily associated with fistula, it seems to me convenient to consider it here.

So much has been written about obstruction for gall stones that at first sight it might seem to be a somewhat common ailment; such, however, is not the case, as may be gathered from the fact that, on inquiring of the registrars and pathologists, I found only four such cases had been treated during a period of twelve months in some of the largest hospitals in the kingdom, representing 80,000 in-patients, and several hundred thousand out-patients attended to during the same time.

Again, only one case, according to Dr. Brockbank, had occurred in the Manchester Royal Infirmary between 1883 and 1896, during which time 50,000 in-patients had been treated. There are clearly four classes of obstruction of the intestines depending primarily on gall stones. The varieties are: 1. That, due to local peritonitis in the gall bladder region, leading to paralysis of the bowels. The symptoms may be so severe as to resemble strangulation by a band, or acute intussusception. The diagnosis will not, as a rule, be difficult, as the history of the occurrence of previous attacks of spasms, though not of necessity followed by jaundice; the similarity to these of the commencement of the attack in question; the severe and persistent pain, at first localised to the right side of the abdomen; the absence of distension at the commencement, and then the occurrence of distension on the right side only, becoming general later; the lateness of the onset of faecal vomiting, and only after continued retching; the existence of collapse at an early stage owing to the severity of the pain, which is usually relieved by a morphine injection; the usual absence of visible peristalsis; and, lastly, the onset of jaundice, if the concretions have reached the common duct, afford so much guidance that error will not often occur, especially if the patient be a woman of middle or old age. But difficulties may arise, as I have found in two cases of my own.

These cases will, as a rule, yield to general and medical treatment, and it will only occasionally be necessary to resort to operation during the seizure if the symptoms are not subsiding, though subsequent surgical treatment may be required.

The second variety, volvulus of the small intestine, dependent on the violence of the colic caused by an attack of gall stones, or on the contortions induced by the passage of a large concretion through the small intestine, is probably uncommon, though two cases have come under my personal observation.

In this class of cases a positive diagnosis is probably, for the most part, out of the question, except after the abdomen is opened, as volvulus of the small intestine is an extremely rare event, and we know that a large gall stone may quietly ulcerate its way into the gut without any preliminary warning, the symptoms only arising when the concretion is passing through the small bowel; but in my own cases, in addition to the signs of acute obstruction, there was a well-marked localised swelling near the umbilicus, becoming hard during the paroxysms, pointing to the site of the obstruction; and in the second case there was not only the previous history of cholelithiasis, but the characteristic onset of a gall-stone attack followed by acute symptoms.

Treatment.—In this class operation holds out the only hope of success, as, the obstruction being mechanical, nothing short of remedying the cause can be of use.

The fourth class is characterised by obstruction coming on after the original cause has disappeared, and depends on adhesions left by local peritonitis due to gall-stone attacks, or on narrowing caused by the healing of a fistulous opening through which a large gall stone has made its way into the intestinal tract.

The third class is the most important variety of obstruction dependent on gall stones, and is the one furnishing not

only the greatest number of cases, but a considerable number of museum specimens. It is dependent on the mechanical obstruction and damage to the bowel produced by the passage of a large concretion through the intestine.

In a paper read before the Royal Medical and Chirurgical Society in 1894, I related notes of cases illustrating this condition; and since then I have successfully operated on another case, and similar ones have been reported by Mr. J. Hutchinson jun., Mr. Lund, and Dr. Everley Taylor.

The following, among other cases which I could relate, will serve to illustrate the passage of gall stones without operation:

On August 26th, 1895, I was asked by Dr. Lever, of Harrogate, to see a lady, aged 73, suffering from acute intestinal obstruction of three days, with faecal vomiting of twenty-four hours' duration. As there was a previous history of gall-stone attacks years before, and as the pain of the present attack started over the hepatic region, we decided to wait and treat the case medically, with the result that a large gall stone was passed naturally after two days, and the patient made a good recovery.

On April 27th, 1896, I received the following notes of a case from a former Leeds house-surgeon, Dr. Wilkinson, of Anerley Hill. My patient is a lady, aged 65, and the facts are briefly an attack of biliary colic followed by symptoms of acute intestinal obstruction, stercoraceous vomiting, etc.

Obstruction lasted three weeks, giving way finally under rest, opium, and copious enemata, and three weeks later a gall stone was passed *per vias naturales* about the size of a pigeon's egg, and weighing 5 drachms, 4½ grains. There was a well-marked facet on the stone, and the patient had only the faintest tinge of jaundice.

In making a study of the reported cases, and especially of museum specimens, one cannot help feeling astonished at finding fatal obstruction depending on quite small concretions, and the comparatively easy passage of very large gall stones. For instance, side by side in Guy's Museum are specimens Nos. 1456, 1457, 1458, and 1459, showing by no means large calculi—one, in fact, only weighing 55 gr.—all of which nevertheless caused fatal obstruction; and specimens Nos. 1449, 1450, and 1451, showing large stones safely passed *per anum*, though in one case the stone was 3¾ inches in circumference, and in another 3 in. long and 1¼ in. in diameter. A case of Dr. J. Blackburn's, 3¾ in. long by 1½ in. broad, actually produced no symptoms except when at the anus; it is now in the Hunterian Museum (No. 143A).

Schuller (Strassburg, 1891), in reviewing 139 published cases, found that the subjects were women in 74.3 per cent., and out of these 75 per cent. of the cases occurred in women over 50, though instances were found from 18 to 94. Lobstein³ (Heidelberg) gives the most common age as between 40 and 60.

It is a curious fact that, although the calculi usually produce intestinal trouble within a few days of reaching the intestine, in some cases they may remain in the bowel for long periods—for example, in a case of Mr. Eve's⁴ ten years and in one of Mr. Smith's⁵ probably fifteen years. In Courvoisier's elaborate statistics, out of 53 cases examined, he gives the site of obstruction as 21.4 per cent. in the duodenum and jejunum, 65.4 per cent. in the ileum, 10 per cent. at the ileo-caecal valve, and 2.4 per cent. in the sigmoid flexure.

Museum specimens amply demonstrate that the gall stones producing obstruction of the intestine in nearly every case enter the bowel through a gall bladder—duodenal fistula. They rarely enter through the colon; in fact, I can only find two specimens of the latter in the London museums.

The disease is a peculiarly fatal one. Out of 280 cases collected by Schuller, Dufort, and Courvoisier, 156 died, or 52 per cent. Kirmisson and Rochard,⁶ out of 105 collected cases, gave the mortality as 50 per cent. The cases that recovered lasted on the average eight days; those that died ten days, but the duration of obstruction may vary from one to twenty-eight days. A case reported by Sargent in the BRITISH MEDICAL JOURNAL, 1879, actually died, apparently from the intensity of the pain, after symptoms lasting only half an hour. Lobstein⁷ collected 92 cases; of the 61 not operated on, 32 recovered, the remaining 29 died of peritonitis or exhaustion; of the 31 operated on 12 recovered, but as many of the 19 which died were moribund when operated on their death cannot be charged to operation.

The symptoms are those of acute intestinal obstruction from other causes, with early faecal vomiting and severe abdominal pain. The higher in the gut the impaction the more violent as a rule will be the symptoms. The obstruction can only very rarely be felt through the abdominal walls. If there is no previous history to guide one it is quite im-

possible to say whether or not the attack is one dependent on the cause in question or on a volvulus or band, or internal hernia, which, if left, must inevitably lead to death, and that speedily. The age and sex, together with the history of chronic dyspepsia and pain in the hepatic region, are, however, well worth bearing in mind, as well as the early and persistent vomiting and visible peristalsis limited to the small intestines.

Treatment.—If the diagnosis could always be made with certainty I think this a condition in which medical and expectant treatment might be fairly given a trial, since we have ample evidence of large gall stones having safely passed without other treatment. But we must not forget that 52 per cent. of cases treated on medical and expectant lines are fatal, and although surgery has not yet shown a much greater percentage of recoveries, it is because surgical means are frequently only resorted to when the case is hopeless and after all other means have been tried.

When it is borne in mind that there are no symptoms peculiar to this form of obstruction, and that the course pursued by an obstruction by a band or by an internal hernia may be exactly the same as in gall-stone obstruction, the surgeon who waits beyond the period when an operation may be undertaken with every hope of success is incurring a very serious responsibility.

With regard to the method of treatment after the abdomen is opened and the cause found, if the gall stone can be easily crushed through the intestinal coats, without too much force being required, so much the better, but that failing enterotomy should be performed, as it can be done very quickly and with very little damage to the bowel. Should the patient be too ill to bear a search being made for the obstruction, enterotomy, or perhaps short circuiting, might be performed in order to give temporary relief, the cause of the obstruction being afterwards removed, if this is not effected naturally.

As to when operation should be performed, no definite rule can possibly be formulated which will apply to all cases. The surgeon will as a rule not be called in before decided symptoms of intestinal obstruction have manifested themselves, and until medical means have been fully tried. In such cases it would seem to me to be idle waste of time to delay surgical intervention until the patient is so exhausted that operation is only undertaken as a *dernier ressort* when the subject is almost moribund. If, however, the case be seen at an earlier stage, morphine will have to be given to relieve the pain, and I have usually recommended ext. belladonnæ in $\frac{1}{4}$ -grain doses every four hours, the stopping of all food by the mouth, and the administration of one or more large siphon enemas, given slowly with the buttocks elevated. If relief does not speedily follow, and the diagnosis is not clear, chloroform anæsthesia may assist in two ways. In the first place it enables a thorough examination of the abdomen, and at times a diagnosis of the cause to be made; and, secondly, the manipulation, if made methodically, may reduce a hernia or volvulus, or may possibly help onwards an obstruction. This failing and the symptoms persisting I should operate, and at this comparatively early stage, I feel sure, with every prospect of success.

TUMOURS OF THE GALL BLADDER AND BILE DUCTS.

If by tumour be understood new growth, then tumours of the gall bladder and bile ducts are not common; but if we accept the usual interpretation of the term, and include all enlargements as tumours, we shall find them by no means rare.

I propose to consider the subject under the following classification, which appears to me to include all the chief varieties:

I. Tumours of the Gall Bladder

(A) Distension:

- (a) With bile.
- (b) With concretions.
- (c) With pus "empyema."
- (d) With mucus "hydrops."

(B) New growth:

- (a) Malignant.
- (b) Simple.

II. Tumours of the Bile Ducts.

(A) Distension.

(B) New growths:

- (a) Malignant.
- (b) Simple.

Distension of the Gall Bladder.—A tumour of the gall bladder though distension with bile must be a rare phenomenon, though it is sometimes described as an accompaniment of a gall-stone attack, where the concretion is impacted in the common duct; even in such a case it must be a symptom of short duration, since if the impaction be complete the bile speedily becomes absorbed, and gives place to distension by mucus. A perceptible tumour formed by distension with gall stones is also rare, unless it happens that some have become impacted in the cystic duct, when a gradual enlargement from the retained mucus will follow. I have removed as many as 750 gall stones from a gall bladder during an operation, and yet that gall bladder could not be felt as a distinct tumour. Occasionally a large single gall stone may form a hard perceptible swelling below the liver, but such is very rare. Calcified gall bladder, which is due to cholelithic catarrh, may lead to the formation of a hard rounded painless tumour, and this is evidently not very uncommon, if we may judge of its frequency by specimens in the museums.

Hydrops and dropsy of the gall bladder are terms used to denote distension of the gall bladder by mucus. It may result from any obstruction in the cystic or common ducts, provided that the gall bladder has not atrophied as the result of previous gall-stone irritation. It is due to the gradual accumulation of the natural secretion of the mucous lining, and may attain such a size as to be mistaken for an ovarian cyst, as in cases reported by Mr. Lawson Tait and by Professor Kocher, though it is uncommon to find the tumour of greater size than 15 to 20 ounces capacity. As an extremely rare event the walls of the gall bladder may form an adipose tumour, as in a specimen from Guy's Museum, No. 1403; the walls of the gall bladder, infiltrated with fat, being one-third of an inch thick. It was removed from a man, aged 66, suffering from kidney disease and cirrhosis of the liver. If the obstruction be associated with inflammation the contents of the gall bladder may become purulent, and an empyema of the gall bladder result.

Signs.—Enlargements of the gall bladder may vary from a tumour just perceptible to the touch to one of such a size as almost to fill the abdomen, though one of greater size than a large pear is exceptional. The same tumour may also vary in size at different times, this variation being frequently found in gall-stone obstruction. The direction in which the gall bladder enlarges is well known, but the position of the tumour will vary with the size of the liver. If uncomplicated, it will have a smooth, rounded, and pear-shaped outline, the larger end being below, quite free, and movable from side to side, the upper end being fixed and passing under the lower margin of the liver at the fissure of the gall bladder. A distinct sulcus between the liver and gall bladder is nearly always perceptible to the touch if the warmed flat hand be laid over the right side of the abdomen and the patient be told to take a deep breath, when the tumour and the liver will descend together and pass under the fingers.

In some cases additional information may be obtained by placing the patient in the genupectoral position, and passing the flat hands round the abdomen from behind, when a tumour of the gall bladder will rest distinctly on the hand, and on deep inspiration it can be felt to move just beneath the abdominal walls. In some of the larger swellings a thrill, almost like the hydatid fremitus, may be felt on gently flicking the tumour with the finger nail. Percussion by no means always elicits dulness coextensive with the tumour, especially if the surrounding intestines be distended, so that dulness on percussion is a very variable sign, and palpation will be found more reliable. Tenderness on palpation is a variable symptom depending on the presence or absence of local peritonitis.

Jaundice may complicate tumours of the gall bladder, both being dependent on the same cause, a blocking of the common bile duct. Although not absolutely pathognomonic of malignant disease, the combination should always raise a suspicion of cancer of the head of the pancreas or of the liver or bile ducts, especially if it be associated with great loss of flesh and strength, and with absence of characteristic gall-stone pain.

I have observed in a considerable number of cases distension of the gall bladder with jaundice to be associated with malignant disease, but much less often the combination

of tumour, jaundice, and gall stones, the explanation of this apparent anomaly being that the gall bladder frequently becomes diminished in size and adherent, as the result of gall-stone irritation, or that when the common duct becomes blocked by a calculus, jaundice occurs without distension of the gall bladder, as it is unable to expand. If, however, the common duct becomes blocked by gall stones before the gall bladder has contracted and formed adhesions, there may be the combination of jaundice and tumour. Gall-bladder tumours usually contain mucus, occasionally pus, rarely bile.

In obstruction of the common duct by gall stones, the gall bladder, though usually contracted, may be found distended by bile at first and mucus later, though, as a rule, the swelling subsides more or less rapidly and no tumour persists, the gall bladder shrinking. When the obstruction becomes absolute, as in malignant disease of the head of the pancreas, the tumour formed is persistent; and although the block is in the common duct, bile soon ceases to reach the gall bladder, and the tumour is always found to contain mucus.

Diagnosis.—Tumours of the gall bladder may have to be diagnosed from: (1) Movable right kidney; (2) tumour of the right kidney or of the suprarenal capsule; (3) tumour of intestine or faecal impaction; (4) tumour of liver; (5) pyloric tumour; and (6) abnormal projection of liver.

The diagnosis of enlargement of the gall bladder from movable right kidney is, as a rule, easy in thin persons. They resemble one another in that each forms a moderate-sized, distinctly defined, rounded and movable tumour on the right side of the abdomen, which is found to descend on inspiration. The previous history may throw light on the case, especially if jaundice has been present. By inspection of the abdomen, a gall-bladder tumour is often apparent, moving rhythmically with the respiratory movements when the patient is recumbent, but a floating kidney can rarely be so detected.

The general outline of the tumour as detected by palpation may afford valuable assistance. The gall-bladder tumour can usually be moved to a limited extent inwards and outwards by manipulation, but under no circumstance can it be depressed into the pelvis. On relieving it of pressure it tends to resume its old position under the liver. Floating kidney has a generally wider movement, can be depressed into the pelvis, and when relieved of pressure tends to pass towards the right loin, especially when the patient is recumbent. A valuable diagnostic sign is the sulcus often felt between the lower margin of the liver and the gall-bladder tumour. In the case of renal tumour as well as in movable kidney, by distending the intestine with gas the kidney will be pressed back into the loin, but the gall bladder will be pushed up towards the liver, and made more prominent. The last test is usually also sufficient to enable a diagnosis to be made between a distended gall bladder and a tumour of the right suprarenal body; but this is not always reliable.

In tumour of the intestine or of the pylorus the associated stomach or bowel symptoms are usually sufficient to enable a diagnosis to be made, but when in doubt distension of the stomach or bowel with gas will help to clear it up, or examination under an anæsthetic will afford assistance.

Tumour of the liver itself, either cancer or hydatid disease, may be almost indistinguishable from one of the gall bladder, though the presence of nodules in the liver, with the history and other symptoms of malignant disease will usually be sufficiently distinctive in cancer, while the less localised and more generally fluctuating swelling, together with the longer history and absence of pain, will distinguish hydatid tumour.

Puncture with an exploring syringe would of course give valuable information, but this should not be lightly undertaken, as it is not devoid of risk, death having occurred on more than one occasion as a direct result of this apparently slight operative procedure. If it is decided to employ an exploring needle the aspirator should always be used, in order that the tense cyst may be completely emptied. In case of doubt, especially where the symptoms demand interference, exploration of the tumour through a small abdominal incision can be undertaken with very little risk, and at the same time further treatment, where called for, can be carried out.

Of the tumours dependent on new growth, cancer of the gall bladder is the most important, innocent growth, excep-

of inflammatory origin, being extremely rare. Cancer of the gall bladder is by no means frequent, and as a primary affection is somewhat rare. It is usually secondary to gall stones or to cancer of adjoining organs, and in the latter case is not amenable to surgical treatment.

Malignant tumour of the gall bladder may probably assume one of four varieties: (a) Columnar-celled carcinoma originating in the mucous glands (the common form); (b) spheroidal-celled carcinoma starting in the surface epithelium; (c) sarcoma starting in the connective tissue; and (d) squamous epithelioma. The disease may occur as a uniform thickening of the walls of the gall bladder, and in the centre of the mass is often formed a cavity containing gall stones.

The very frequent association of cancer of the gall bladder with gall stones is an undoubted fact, and in all probability there is a connection between the two diseases. Zenkert⁹ found gall stones in 85 per cent. of cancers of the gall bladder; and Masser, from an analysis of 100 cases, gives the proportion associated with gall stones as 69 per cent.; while Courvoisier found gall stones present in 74 out of 84 cases of primary cancer of the gall bladder; and other observers give an even higher proportion.

The two theories which have been current to explain the coexistence of gall stone with cancer of the liver, are first, the "irritation" theory, that gall stones are formed first, and by acting as foreign bodies set up irritation which leads to malignant growth; and secondly, the "concentration" theory, that gall stones arise as a secondary result from stagnation of the bile in the ducts, caused by their obstruction from malignant growth.

Mr. C. Beadles, in a paper before the Pathological Society of London,⁹ stated that out of 100 *post-mortem* examinations at the Cancer Hospital, 4 were cases of primary carcinoma of the liver, and all had calculi in the gall bladder; 36 had secondary carcinomatous growths in the liver, but there were no gall stones present in any of them.

Symptoms and Signs.—If the growth be primary, there will be the history of a more or less rapidly-growing tumour developing under the right costal margin, accompanied at first by a sense of discomfort, shortly changing to pain which, though at first localised to the right hypochondrium and epigastrium usually before great advance has been made, extends round the side to the right infrascapular region. When the enlargement is first noticed, it is felt as an egg-shaped swelling beneath the liver, descending on inspiration. The tumour is hard to the touch, and very slightly or not at all tender to pressure. At a later stage, it becomes more fixed and more diffused, and nodules may develop and be felt on its superficial surface.

According to the invasion or not of the hepatic or common bile ducts, so will be the presence or absence of jaundice, but in nearly half of the cases some degree of icterus will be found as the disease advances, owing to the presence of catarrh of the bile ducts.

Interference with the action of the bowels even to partial or complete obstruction at times occurs. General failure of health, continued wasting with loss of strength, ascites, and marked cachexia characterise the later stages.

Perforation may occur and hasten the end by the onset of general peritonitis. Where gall stones with jaundice complicate cancer of the gall bladder exacerbations of pain will usually be accompanied by rigors and fever—"ague-like attacks," with an intensification of the icterus, and in such cases petechiæ in the skin, with hæmorrhage from the nose and rectum usually supervene.

Diagnosis.—Cancer of the gall bladder may usually be diagnosed by the progressive character of the disease, and by the presence of the characteristic hard tumour; but it is by no means always easy to diagnose cancer from a tumour formed by matted intestines due to local peritonitis in the neighbourhood of the gall bladder.

When in doubt exploration is probably the best method of settling it, as at the same time treatment may be carried out. Cancer of the right suprarenal body may afford a difficulty in diagnosis, and the same applies to cancer of the pylorus, which, however, is accompanied, for the most part, by characteristic stomach symptoms.

Treatment.—The alleviation of symptoms, especially of

pain, by sedatives is practically all that can be done, except in those rare cases where the disease is limited to the gall bladder, when cholecystectomy may be performed.

TUMOURS OF BILE DUCTS.

Tumours of the bile ducts *per se* only occasionally form a projection so large as to be distinguished through the abdominal walls. The common duct has, however, been found dilated to such a size as to form a cystic tumour presenting all the characteristics of a distended gall bladder, the gall bladder itself being atrophied.

Terrier has recorded four cases in which an external fistula led into the common bile duct, and I have had two of a similar nature, in one of which I performed choledochostomy after choledochenterostomy, and in the other choledochenterostomy after cholecystectomy.

Although hitherto the results of choledochostomy have not always been favourable, probably in consequence of the fact that extreme distension of the bile duct is often accompanied by infection of the biliary passages, it would be well to reserve our opinion as to the prospects of the operation until we have more experience of it.

The new growths found in the bile ducts may be simple or malignant. Of the malignant we must take into consideration the two classes—primary and secondary; the former arising most frequently as the result of gall-stone irritation, the latter by extension from neighbouring organs.

Primary malignant growth may be: (a) Cylindrical-celled carcinoma; (b) spheroidal-celled carcinoma; (c) sarcoma; (d) squamous epithelioma. In the Hunterian Museum, specimen No. 2809 is a melanotic sarcoma of the gall bladder.

Out of the 11 cases of malignant disease of the bile ducts recorded by Dr. Rolleston, only 4 were associated with gall stones. He thinks that calculi are less frequently associated with primary cancer of the bile ducts than with cancer of the gall bladder. In Dr. Rolleston's paper a case is referred to, which makes it probable that an adenoma preceded cancer. Papilloma is probably an earlier stage of cancer, and is rare. Cancer is usually secondary to gall stones, though they may not always be found, having passed into the bowel before operation. Although these tumours are usually seen in the common duct, they may occur in the cystic or in the hepatic ducts. In Musser's 18 cases the hepatic ducts were alone involved three times, the cystic and hepatic ducts once, and the common duct fourteen times. In Rolleston's 11 cases the common duct was always the seat of the tumour. These growths are usually found between 50 and 60, and, unlike cancer of the gall bladder, where 75 per cent. of cases occurred among women, this disease attacks both sexes equally, the symptoms of course vary with the ducts affected. Suppurative cholangitis is apt to supervene, the case then taking on a more acute course, and being accompanied by fever, ague-like attacks, and rapid loss of flesh and strength. Needless to say, the disease is uniformly fatal, though operation may delay the final catastrophe.

Cancer of the ampulla of Vater has received some attention from M. Hanot,¹⁰ M. Durand-Fardel,¹¹ M. Rendu, and Dr. Rolleston.¹² Two forms at least are described: one arising from the intestinal walls of the ampulla, and the other from the orifice of the common bile duct and the pancreatic duct. They are characterised by intermittent jaundice, wasting, and cachexia, but as a rule pain is absent.

Diagnosis.—The diagnosis of new growth in the bile ducts from gall stones is practically impossible, as the symptoms are the same, and in fact the two frequently coexist. The absence of pain in some cases and the rapid deterioration of health may afford a little help, but in some cases the pain may be as acute as in cholelithiasis. Cystic dilatations of the bile ducts are often indistinguishable from enlargements of the gall bladder, for which they are usually mistaken, but they may resemble a cyst of the pancreas or a hydatid tumour of the liver; but as the treatment in all these conditions is abdominal section, no harm will be done if the diagnosis is only completed when the abdomen is opened.

Treatment.—The operative treatment of these tumours is in its infancy, and has thus far not proved uniformly satisfactory. Any growth should be removed if possible, but where that is impracticable the dilated gall bladder or ducts may be opened and drained, or better still drained into the

duodenum or jejunum by means of one of Murphy's buttons. If the cause be a removable one, such as a gall stone, it should be taken away. Choledochostomy has not yielded good results in cystic dilatation of the bile ducts, whereas the results of choledochenterostomy establish its claim to be considered the best method of treatment.

REFERENCES.

- ¹ *Beiträge zur Pathologie und Chirurgie des Gallenwege.* ² *Lancet*, 1878, 1, p. 851. ³ *Annals of Surgery*, January, 1896. ⁴ *Trans. Clin. Soc.*, 1895. ⁵ *Lancet*, December 3rd, 1887. ⁶ *Arch. Gén. de Méd.*, February, 1892. ⁷ *Loc. cit.* ⁸ *Deut. Arch. f. klin. Med.*, 1889. ⁹ *Lancet*, March 9th, 1895. ¹⁰ *Arch. Gén. de Méd.*, November, 1896. ¹¹ *La Presse Médicale*, 1896. ¹² *Med. Chir. Trans.*, 1895.

THE MILROY LECTURES ON QUARANTINE.

*Delivered before the Royal College of Physicians of London,
March, 1897.*

By W. COLLINGRIDGE, M.A., LL.M., M.D.,
Medical Officer of Health for the Port of London.

[ABSTRACT.]

THE CAUSES OF THE FAILURE OF QUARANTINE.

HAVING thus sketched the history of quarantine and its final suppression by medical inspection, the chief causes which have gradually led to its abandonment may be summarised as follows:

1. That the quarantine system was based upon a theory which is no longer tenable.
2. That it has never been found possible to maintain it absolutely, and that consequently a country has been subjected to a barbarous, unnecessary, and costly system, without even the satisfaction of knowing that the experiment was complete.
3. That, where carried out as strictly as possible, it has not had the effect of arresting the importation of disease.
4. That sanitation and a rational system of medical inspection have proved more efficacious, and at the same time less burdensome.

The whole history of quarantine is a series of illustrations of its futility, and proves it to be, as Sir John Simon so graphically described it, an elaborate system of leakiness. Theory must give way to practice, and the real test of quarantine as a system to be relied upon by the Government of a country for its protection must eventually be that of actual experience.

As soon as investigation was made into the facts it was extraordinary how little evidence there was of the value of the system for the purpose which it was intended to serve.

An effective quarantine is practically impossible in a commercial country. In early days, when trade was relatively of small volume, this was apparent, and the enormous growth of commerce has simply emphasised the fact. The question could only be considered at the present day with reference to the period of incubation, and it would obviously be accepted that any vessel was safe if there had been no case of disease during a period at least equal to that of the possible latency of the disease.

THE INEFFICIENCY OF QUARANTINE.

It is scarcely possible for those not actually interested in shipping to realise how serious a matter quarantine has been for shipowners, not only in the remote past, but even up to a very recent date. First in importance probably would come without doubt the loss occasioned by delay and uncertainty of detention, and even in some cases actual prohibition of intercourse with infected port or ports supposed to be infected.

Dr. Bowring, after extended travel and experience of quarantine, says that "the pecuniary loss inflicted by the system may be estimated by millions of pounds sterling in delays, demurrage, loss of interest, deterioration of merchandise, increased expenses, fluctuations of markets, and other calculable elements; but the sacrifice of happiness, the weariness, the wasted time, the annoyance, the sufferings inflicted by quarantine legislation—these admit of no calculation, they exceed all measure."