trix had a raised and glassy look. In a few days it thinned down to a mere blister, which gave way, allowing the escape of about 3i of foetid pus. There was nothing faecal about it, however, and I never heard that such was the case at any period of the patient's illness. Mr. Mitchell Banks now took up the case with me, and on passing a probe he found it passed $4\frac{1}{2}$ inches into a tortuous channel running downwards and towards the middle line in the direction of the bladder; it was clear the original wound had never healed soundly but had left behind it this long sinus, the orifice of which had closed over imperfectly, and then had given rise to the belief that everything was sound. Some pus having accumulated in the sinus had caused the patient to feel very ill, a condition from which he at once recovered as soon as the cicatrix gave way and the pus escaped.

To lay this extensive sinus open in its entire length was a distinctly serious proceeding, not devoid of danger, and with the possibility of leaving behind it a weakened abdominal wall, which would be a serious thing for an officer. It was resolved, therefore, to try to get the sinus to heal if possible by various stimulating and other liquid applications, nitrate of silver, etc., all to no effect. At last Mr. R.'s patience became exhausted, and he resolved to have the sinus completely laid open, which was done by Mr. Banks on August 19th. The steps of the operation are as described by himself. As the process of laying open went on the incision necessarily became deeper and deeper. The rectus, and with it the epigastric artery, were completely divided, the incision finally ending in the suprapubic fat and cellular tissue. It was impossible, moreover, to help opening the peritoneum. Through the rent the finger introduced into the abdomen found no evidence of adherent intestine; all seemed perfectly right.

The point of interest, however, is that in the course of laying open the sinus something was cut across which exactly resembled a very small intestine; it was recognised to be the appendix vermiformis, and was stitched up. The sinus was thoroughly scraped in its whole extent, and having been put into the condition of a fresh raw wound was stitched up in its entire length, except for a small portion at its lower end, in which a tube was inserted. By this means primary union was obtained of the peritoneum, and through a portion of the long gash, thereby greatly diminishing the liability of rupture which would otherwise have resulted. The wound finally closed in December, and, having gone away for a change, the patient returned to duty in the beginning of March, the part being protected by an aluminium plate. He has remained well up to the present date.

The points of interest are:

- (1) The occurrence after typhoid fever of an abscess probably perityphlitic in character.
- (2) The occurrence of a long and deep sinus after this abscess, which had closed over unsoundly, only to give rise to a collection of retained pus.
- (3) The discovery and division of the vermiform appendix imbedded in the hard tissue of the sinus.
- (4) The satisfactory result of the operation, and the complete recovery, after so prolonged and serious illness.

POISONING BY THE "PRIMULA OBCONICA."

By J. HEADLEY NEALE, M.B., M.R.C.P.LOND.,

Physician to the Leicester Infirmary.

ON April 9th, 1896, Mrs. W., a young married lady, consulted me for a rash which had suddenly appeared the day before on her face and hands, "which itched and burned intensely." On the face the eruption was urticarial, but the blotches were too confluent; while on the hands, especially between the fingers, it was suggestive of scabies. I suspected some local irritant, and finding that the soap used was not irritating I described as carefully as I could the flowers of the *P. obconica*, but she denied having handled any. Under a topical soothing application the face was soon better, and desquamating, but the hands became intensely painful. On the dorsal aspect the spots had turned to purple blotches, while the whole palmar surface of hands and fingers had become stiff and unusable. Two days later the face was comparatively well, the blotches on the dorsal aspect of hands

and fingers fading, but there was evidence of deep-seated blisters forming at the tip of each finger and above and below each phalangeal flexure. To make sure I smartly pricked the tip of one finger and a bead of clear fluid escaped. I ordered the patient to macerate her hands thrice daily for ten minutes at a time in rectified spirit (to which was added a fair proportion of tincture of belladonna), and then to swathe them in lint soaked in glycerine and covered with oil silk, relying upon the hygroscopic properties of the spirit and glycerine and the possible action of the belladonna as an anodyne. The result fully justified the experiment. A few days later she told me that "her hands were nearly well," and she "could bend her fingers." The first thing that attracted my attention as I entered her room was the presence of three fine blooms of the *P. obconica* among other flowers on the table. On pointing out that these were the very flowers I had been looking for and asking about, she told me she had gathered them the day before in the conservatory, and on going there I found eight fine plants, some in full bloom, which she had been in the habit of watering, etc. That they had not bloomed on April 9th, when the rash appeared, opens out a fair field for research as to the properties of the several parts of this pretty but obnoxious plant.

This case recalls to my mind the result of the enthusiasm of a young assistant at the Botanical Gardens in Edinburgh. Having in his menial capacity been present at a lecture when the late Professor Dickson was decanting upon the poisonous properties of the "*Rhus toxicodendron*," this inquiring youth straightway broke a twig and smeared some of the exuding juice on his left arm, appearing at the infirmary a few days later. Never shall I forget that arm with its gangrenous-like line of demarcation, midway between the shoulder and elbow-joint, while below it was intensely red and swollen; one huge pemphigoid eruption, some of the "bullæ" being as large as pigeon's eggs. Nor shall I easily forget the enthusiasm displayed upon the faces of Professor Fraser and Dr. Murrell as, between the "clinics" which were then in progress, they watched me making a water-colour sketch of the arm. The drawing is no doubt still in Dr. Fraser's possession, and is possibly the only one extant of the result of poisoning by "*Rhus toxicodendron*."

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

AN ANENCEPHALOUS MONSTER.

ON the afternoon of April 10th I was called to attend a lady, aged 28, a primipara, eight months pregnant. She had suffered greatly from pain in the abdomen for some days previously, and had not slept the preceding night owing to it. She was a rather thin woman of average height. The abdomen was of large size, the abdominal walls very tense, rendering palpation difficult. The fetal heart was audible at the left side, above the level of the umbilicus; the movements of the fetus could be felt low down on the right side, and were, indeed, a source of discomfort to the patient. I could not feel the head at the fundus. The os was dilated to the size of a shilling.

I ordered a simple enema, and then, having administered a dose of chloral and opium, left the house. On the following day the patient felt much better, and was able to be out of bed and came downstairs.

About 7 A.M. on April 13th I was again sent for. Throughout the night the patient had severe and frequent labour pains; the membranes ruptured before my arrival, and a foot presented. The child was delivered in about twenty minutes; when placed on the bed it did not cry, but moved its arms and legs freely for about a minute, during which time the chord pulsated distinctly; it soon, however, ceased to move.

The child, a female, was 15 inches long, the arms and legs were very thin, but of average length, the abdomen normal. It had no neck whatever, the chin being continuous with the thorax. The face was very small and the eyes prominent. The head at the level of the supraorbital ridges was perfectly flat, covered in by pretty hard bone and skin, through