tact and resolution may be requisite in pressing for an inquiry into this important matter; indeed, some individuals seem to resent almost as a personal affront a hint that the sanitary arrangements of a house which they have made their abode can be otherwise than perfect. Latterly, I have seen a great many cases from India, more especially from certain parts of Bombay and Burmah, in which acute inflammation of the tympanum has gone on to perforation and discharge, and I have been unable to attribute this mischief to anything else than foulness of atmosphere from unhealthy surroundings.

THE SAFE ADMINISTRATION OF ANÆSTHETICS : WITH SPECIAL REFERENCE TO CHLORO- FORM AND METHYLENE.

A REPORT OF SIX HUNDRED CASES.

By CHARLES O'NEILL, M.D., MCh., D.P.H.,
Belfast.

THE high death-rate caused during the administration of anæsthetics is a sufficient reason for every medical man so to study the subject and perfect the system of giving anæsthetics as to reduce the mortality to the lowest possible amount.

THE PREPARATION OF THE PATIENT.

The patient should be prepared by taking a hot bath the night before, and at bedtime a brisk purgative, and the following morning a light breakfast about three hours before the anæsthetic is administered. Giving an anæsthetic on a fasting stomach tends to retching, and occasionally great depression and weakness. Should the patient have taken a full meal shortly before having an anæsthetic, it often causes vomiting during the administration, allowing them to recover and causing delay. When vomiting occurs during its use, the patient should be turned on the left side, so as to prevent any of the vomited matter from entering the trachea. Should vomiting occur after the anæsthetic is stopped, it is merely a sign of returning consciousness.

Before giving an anæsthetic you should examine carefully for any signs of mitral disease, fatty degeneration or dilatation of the heart, or calcareous degeneration in the vessels, or chronic bronchitis, asthma, or pleurisy. Should abdominal dropsy be present, sufficient to cause thoracic pressure, any of these conditions should warn you to use great care in the administration. See that there is no tight clothing on the chest or abdomen; everything must be loose to allow of free breathing. If the patient has got artificial teeth, they must be removed. If possible the urine should be examined for albumen and sugar.

THE METHODS OF ADMINISTRATION.

In administering anæsthetics, one of two ways may be employed. First the so-called "open" or "Syme's" method—why so called is a mystery, as it is really a method where the fresh air is almost entirely excluded from the patient while taking the anæsthetic. In this method you pour a certain quantity of the anæsthetic on a towel, sponge, or lint, and hold it over the patient's mouth and nose. The distance it should be kept off the face cannot be ascertained in any reliable manner. On the other hand, distance makes little difference whether it is held 2, 4, or 6 inches from the face. If the atmosphere of the room is quiet, the vapour drops down in a more or less undiluted form, being heavier than air; while if the room is draughty it is more uncertain still, as the patient may only get an occasional breath of the anæsthetic. Unless in midwifery practice and in some exceptional cases, I consider this a most unscientific way of giving it. It is idle to talk of graduating the dose of the anæsthetic to the age and requirements of the patient, so long as this rough and ready way is employed. We should dilute an anæsthetic with air as carefully before it is breathed as we would dilute hydrocyanic acid with water before administering it to the stomach. In this method the dose is uncertain, and depends, not on what you put on the towel, but on what quantity reaches the lungs, and the patient is compelled at first to inhale air saturated with the vapour of

the anæsthetic until little is left on the towel. On adding more you keep the patient oscillating between an overdose and an underdose. This is a very wasteful way of giving an anæsthetic, a case being recorded in the BRITISH MEDICAL JOURNAL of July 14th, 1894, where a female, aged 25, had 12 drachms administered in thirteen minutes before becoming unconscious. The operation lasted fifty minutes, and the chloroform used was 31 drachms.

The second method, which I consider the safe one for administering methylene or chloroform, is by giving it through a Junker's inhaler, which has been much improved by Messrs. Krohne and Sesemann, London. The improvements on the older inhaler consist in the addition of a respi-

Junker's inhaler.

ration indicator and a soft inflating indiarubber cushion fitted on the facepiece. When this inhaler is used you can regulate the dose, and the patient at each inspiration takes in the normal supply of air properly mixed with the anæsthetic. It may be used with equal safety for persons of any age. In giving an anæsthetic a most important point is carefully to observe the breathing; see that it keeps regular, and stop giving the anæsthetic whenever it shows signs of irregularity, and allow free access of air until it becomes normal. Since the breathing is of much more importance to observe than the pulse, the pulse is of importance where the patient has had severe hæmorrhage previous to or during the administration. In these cases the pulse is very important to watch carefully for showing signs of collapse. In ordinary cases it is sufficient to watch the respiration, as there is great danger in watching the pulse to overlook the breathing, since the heart is only affected secondarily in consequence of the failure of the respiration. If an anæsthetic is pushed so far as to cause the pulse to show signs of heart failure the limit of safety has been so far exceeded that a fatal result is likely to ensue. One great source of danger in administering anæsthetics is when the person giving it watches some interesting operation and takes his attention completely off his own duties. Should the patient cease breathing for thirty seconds it is almost impossible to restore animation. Another danger is when the patient struggles while taking the anæsthetic and requires holding, you should see that no pressure is applied to the chest or abdomen, as the breathing obstructed in this way causes collapse. If the tongue falls back it is also a source of danger; or the patient may require to be turned on the side for an operation to be performed; this may also cause dangerous symptoms.

Death from sudden stoppage of the heart without any previous warning from irregular breathing is most frequently caused by giving too much of the anæsthetic, since in an overdose the supply of air is cut off, and in its place a narcotic is given. The risk is increased by the patient struggling, or by anything which causes them to take gasping or irregular inspirations. Sir James Simpson advised the administration of a large dose at first as he considered the state of semianæsthesia the critical time. Of course, fatal cases have occurred at the beginning of the administration, but it is more likely they occur from an overdose than an underdose, as the more of the anæsthetic used the less air is inhaled. In using a Junker's inhaler, you add the vapour of the anæsthetic to the normal quantity of air used in breathing. In giving an anæsthetic to grown-up persons, you should instruct them to breathe in the ordinary regular way,

and try if possible to reassure them, should they show signs of nervousness, and bring the facepiece gradually forward, so as to avoid the suffocating sensation which is caused by bringing it suddenly over their mouth and nose. If you succeed in this, the patient, generally speaking, gives very little trouble; but, on the other hand, without giving any instructions to the patient, you place the facepiece suddenly over their mouth and nose. This causes a feeling of suffocation, and so frightens them that it is with great difficulty they are quieted, and all through the administration they show signs of nervousness. The only exception to this is in the case of children; you should put them over as quickly as possible, as they nearly always shout and struggle.

THE ADVANTAGES OF METHYLENE AND CHLOROFORM.

Methylene or bichloride of methylene, discovered by Sir Benjamin Richardson in 1867 (CH_2Cl_2), contains one volume less chlorine than chloroform. Some consider it only diluted chloroform, but this is an error, as it is a distinct preparation. Its advantages as an anæsthetic when given through a Junker's inhaler are: 1. It is quick in action; you can have complete anæsthesia in an adult from five till ten minutes. 2. It may be prolonged up till two hours or more with safety. 3. It seldom causes sickness or vomiting. 4. The patient recovers consciousness from three till five minutes after its administration is stopped. Sir Spencer Wells in 1,200 cases used this anæsthetic, to the exclusion of all others. But there are certain cases and certain classes of patients where this preparation is not strong enough to produce full anæsthesia. I shall specify some of these in speaking of chloroform. I do not wish it to be understood that any anæsthetic is absolutely safe, since there are always a certain number of persons which constitute a class where a comparatively slight shock may prove fatal. In these cases death is caused by syncope, or sudden arrest of the heart's action, and no previous warning is given. In using methylene through a Junker's inhaler we have the safest anæsthetic at present known, provided the precautions I have mentioned are strictly attended to.

Chloroform (CaHCl_3) contains one volume more chlorine than methylene. It can be administered the same way—that is, through a Junker's inhaler. As it is a stronger preparation than methylene, the danger of giving an overdose is greater. There are four unfavourable symptoms I shall give in their order of importance, which should be carefully attended to: (1) coughing, (2) gasping, (3) choking, (4) struggling. We should pay attention to the first symptom, "coughing," which is a sure sign that the patient is getting the anæsthetic too concentrated. If this warning is neglected, the patient may become worse, and a fatal issue may result. By attending to these symptoms and remedying them in time, we find that prevention is better than cure.

When chloroform is administered the patient loses consciousness sooner than in methylene; but they are more liable to be sick, and do not recover so soon after its administration is stopped. But on the other hand there are, as I mentioned before, certain operations and certain classes of patients which require this stronger preparation, such as persons of intemperate habits and those of a nervous temperament, and also in operations of a painful nature, such as excision of piles, etc.

RESUSCITATION IN COLLAPSE.

Should collapse occur during the administration of an anæsthetic, we should always have a number of suitable remedies beside us to be prepared for any emergency—such as nitrite of amyl, strong solution of ammonia, ether, liquor strychnine, and a hypodermic syringe, tongue forceps, and an electric battery—as prompt application of remedies is a very important factor in restoring a patient. Should a patient show signs of collapse—that is, should the breathing get irregular, slow, sighing, or stop, or the face suddenly get pale, or the lips cyanosed—we should stop the administration immediately, lower the patient's head, draw forward the tongue, and use artificial respiration according to Sylvester's method, apply nitrite of amyl or strong solution of ammonia to the nostrils, inject ether hypodermically, or liquor strychnine (from three to five drops for an adult), apply the poles of the battery over the diaphragm; pure oxygen may also be

administered. Inversion of the patient may also be tried. But if a patient suddenly stops breathing, and the heart collapses at the same time, any restorative we apply is almost useless. At the present time we have at least fifty different ways for restoring consciousness after an overdose of an anæsthetic. A great number of these hail from America. But we require little explanation how to treat collapse if we administer the anæsthetic carefully; for if you give air too much impregnated with an anæsthetic to maintain life, it is difficult to restore breathing when life is almost extinct.

I shall now refer to a series of 600 cases extending over the six years from 1890 till 1896, where a note was taken of the age of the patient, the quantity of the anæsthetic used, and the average time taken to produce anæsthesia, as well as remarks on sickness and vomiting. These include both methylene and chloroform cases. The ages ranged from 1 hour (in this case it was given to aid in the reduction of an umbilical hernia) up till 80 years. The quantity given every five minutes to an adult until complete anæsthesia was about 40 minims; complete anæsthesia usually occurred from five till ten minutes, and about half this quantity was required to keep up the anæsthesia, which is at first in the proportion of $\frac{2}{3}$ of a minim during each inspiration, taking 20 inspirations per minute as normal, and afterwards $\frac{1}{2}$ of a minim each inspiration to keep the patient unconscious. Three drachms was the usual quantity used in half an hour, and about 4 drachms was sufficient for an hour. Two per cent. of the vapour is enough to use until the patient gets accustomed to the anæsthetic, and afterwards increase it to about 4 per cent. Had the open method been used this quantity would have been altogether insufficient. The average length of time until complete anæsthesia occurred was in infants two minutes, children three minutes, and in adults from five till ten minutes. Great care should be taken not to produce anæsthesia too quickly, as this is a great source of danger. In 12 per cent. of the cases sickness and vomiting occurred either during or after the administration. Up to the present time I have never had a death from the effects of an anæsthetic.

Regarding fatal cases occurring we should never treat lightly the giving of an anæsthetic, no matter how short a time the administration is required, since fatal cases have occurred during any stage, even in the first few inspirations; therefore we should keep before us the fact that while giving an anæsthetic the patient's life is in our hands. My reasons for placing methylene before chloroform are it is safer, and adapts itself for use in minor operations. In these cases we would hesitate before giving a stronger preparation, and so deprive the patient of the benefit of a painless operation. Therefore we have for all practical purposes in these two valuable anæsthetics all that is required to allow of any operation being safely performed without pain.

THE report presented by the Water Committee to the London County Council for the six months ending April 27th states that there is a remarkable disparity between the quality of the water delivered by the different companies, notwithstanding the fact that five of them draw their water from the same source. The Southwark and Vauxhall Company is reported to have allowed to pass twice as much suspended matter as the other Thames companies; and in the case of that company the samples containing objectionable matter were nearly thrice as numerous as in the case of the West Middlesex Company. Great variations occurred also in the number of occasions on which the proportion of microbe present exceeded 100 per cubic centimetre. This was the case with the Kent Company on 14 days out of 100, but on 58 out of 100 days with the Southwark and Vauxhall Company.

THE foundation stone of the Acton Cottage Hospital was laid by Lady Rothschild on June 9th. The site has been given by Lord Rothschild and Mr. Leopold Rothschild, while Mr. Passmore Edwards has subscribed £2,500 towards the building expenses. The two main wards will each contain four beds, but these will be in addition convalescent wards.

THE Drapers' Company of London has offered £15,000 to the University of Oxford for the erection of a building to provide additional accommodation for the Radcliffe Library. This munificent gift has been gratefully accepted.