smoothly keeping both eyes open. No one who has seen a patient work his way through these several stages can doubt that bar-reading-aids in the recovery of binocular vision. It is chiefly effective when practised in conjunction with the use of the shade. Its weak side is that it is inapplicable in very early life. To older patients it affords interesting occupation.

Fig. 5.—Bar-reading.

[Cases of long-standing strabismus were cited in which treatment by glasses and operation left a slight deviation, with no diplopia and no tendency to fuse, and in which the subsequent use of the shade and reading bar restored binocular vision in the course of three to six weeks.]

The fusion tubes were used, together with other treatment, in 21 cases. The plan was to let the patient hold them in such a way as to blend the two white holes; then to separate and join these many times in succession, so as to get the idea of fusing double images. Then to overlap them partly, and try to make them go together by looking at them. Lastly, to practise this at home. Seeing that this exercise grows easier with a little practice, I presume that it serves to stimulate the sense of fusion. It is irksome if long continued, but interesting for a while, and can sometimes be managed by children who are too young for the reading bar. With the stereoscope proper I have done nearly nothing; time failed for any sufficient trial.

The old idea that the squinting child needs a little skilful cutting and nothing more is dying out. Most people know that glasses are often necessary; but that the child may require teaching, at some trouble, to use the squinting eye is a new idea to many. But it is easily grasped, and must be grasped if our efforts are to be effective. I have used the word "educative" as a help in that direction. Of course there are parents and there are children on whom all time and trouble spent in this way are entirely thrown away, but on the whole I have been surprised at the care and patience with which directions have been carried out even in homes where one might least expect it. In this lecture I have stated only my own experience; it has been impossible to compare it with that of others. I have referred but little even to Javal, but those who know his work, especially his recent *Manuel du Strabisme*, will know how much I am indebted to it. On some points of etiology his views are, I think, open to question, but as regards the education of the visual function which he has so ably advocated, I am convinced that the principle is true and the method practically useful. Used rarely by itself, usually in conjunction with other treatment, it promotes real as distinguished from apparent cures of squint, and gives the maximum security against relapse.

REFERENCES.

¹ *Manuel du Strabisme*, Paris, G. Masson, 1896. ² and ³ The fusion tubes and heteroscope were made by R. Bailey, Bennett's Hill, Birmingham. ⁴ *BRITISH MEDICAL JOURNAL*, June 20th, 1896. ⁵ W. Preyer, *Die Seele des Kindes*, Leipzig, Grieben, 1884. ⁶ See Automatism and Control in Lloyd Morgan's *Comparative Psychology*, London, Walter Scott, 1894. ⁷ B. A.

Randall, A Study of Refraction in School Children, *American Journal of Medical Sciences*, July, 1885 (p. 25 in reprint). ⁸ *Ibid.* ⁹ See also Barratt and Lang, *Roy. Lond. Ophth. Hosp. Reports*.

THE CAVENDISH LECTURE

ON

GROWTH AS AN AGENT IN (1) PRODUCTION AND (2) THE REMOVAL OF DEFORMITY.

Delivered before the West London Medico-Chirurgical Society

By HOWARD MARSH, F.R.C.S.,

Surgeon to, and Lecturer on Surgery at, St. Bartholomew's Hospital.

GROWTH has been defined as the increase of a part by the addition of material like that already existing in it. In its simplest form growth may be observed in the enlargement of a crystal. If a crystal of alum sulphate, for example, is placed in a saturated solution of that salt it gradually increases in size as more and more alum sulphate is incorporated in it. Another equally familiar example is the growth of a bone in a young animal. The tibia, for instance, becomes larger and larger by the addition of more and more bone to its length and thickness. When normal growth is examined it at once becomes apparent that it takes place in conformity to type. This is illustrated in a remarkable manner in the case of crystals. A crystal of alum sulphate, when it is perfect, is, whatever its size and at all stages of its progress from small to large, always octohedral. But, further, if a crystal of alum sulphate is broken, and is then placed in its mother solution, the defect is repaired, and at length it is restored to its typical shape, and becomes once more a perfect octohedron.

Here are casts of some crystals (No. 42B in the Museum, St. Bartholomew's Hospital) prepared many years ago by Sir James Paget to illustrate an experiment recorded by Jordan in *Müller's Archiv*, 1842, page 46. A broken crystal of alum was placed in a solution of this salt, and in a few days "while the whole crystal had increased in size, the increase on its broken surface was proportionately so much greater than on any other, that the perfect octohedral form was nearly regained." A few weeks ago, at my suggestion, Dr. Chattaway, in the Chemical Laboratory of St. Bartholomew's Hospital, selected two crystals of alum sulphate of the same size (Fig. 1 (A) and (B)), and then, breaking off a large piece

Fig. 1.—Repair of a crystal. A crystal of alum sulphate (b) was fractured at the level of the dark line shown on the other crystal (a). It was then placed in a solution of alum, and the part which had been broken off is seen to have been, in the course of a month, nearly reformed.

from one, marked the other to show the line of fracture. When placed in a solution of alum the broken crystal, growing in correspondence with its type, had within a month, to a great extent, as a glance at the two figures shows, regained its original shape.

In studying growth in the vegetable and animal kingdoms

respectively, it is found that while its ultimate object is in both the same—namely, adaptation of the individual to its environment—an important difference may be observed. In plants the influence of type is comparatively limited, for there are numerous instances in which a plant, when its environment is changed, is able, as it grows, so far to adapt itself to the new conditions under which it is placed that it becomes modified in a remarkable degree. This is well seen when the land forms are compared with the water forms of the same amphibious species,¹ or when the habitat of a plant is changed from a cold to a hot climate. The castor-oil, for instance, which is an annual in England, becomes a tall tree in Malta; while M. Freyn states that *Beta maritima* on the cliffs of Istria grows in shrub-like form, with erect stems, bearing hardly any resemblance to our cultivated beet.² Again this capacity for adaptation to novel circumstances was graphically demonstrated by an experiment recorded by Dr. Lindley. A young willow tree had its crown bent down to the ground; this was covered with earth, and soon emitted an abundance of roots. The true roots were then carefully removed from the soil, and the stem inverted. The roots now became branches and emitted buds, and the tree grew ever afterwards upside down.³

In animals, on the other hand, the law of conformity to type exercises a much stronger influence, so that, although adaptive variation takes place in a series of generations, yet it is scarcely appreciable in the life of a single individual.

When at length the individual has attained its complete form and structure, and is thus in correspondence with the type which best equips it for the functions of its life, general growth, its purpose having been fulfilled, comes to an end. In man this stage has, generally speaking, been reached by about the twenty-first year when, as the phrase is, the individual "has done growing." Variations, however, in the directions both of deficiency and of excess may occur so that the normal stature is either exceeded or not reached. Hence giants and dwarfs. Or the variation from the normal may be limited to a single part. In such cases what has been termed formative power or germinal capacity is in local excess, with the result that such specimens as these of so-called congenital hypertrophy of fingers and toes are met with. The middle finger of this hand is nearly five times its proper bulk; and this is a cast of the foot of a boy aged 13. The foot was 12 inches long, and the instep $11\frac{1}{2}$ inches in circumference. The great toe was $4\frac{1}{2}$ inches long, and had a circumference of $6\frac{1}{2}$ inches.

Yet on closer examination it is observed that, after general growth has ceased, the capacity for further growth in many parts still exists in a latent form or in reserve, but ready, should the interest of the individual require it, to be brought into activity. This provisional capacity for the growth of individual parts or organs after general growth has ceased is one of the most striking and valuable resources with which the individual is endowed. General growth adapts the individual to its environment under normal conditions, while latent growth serves a different purpose; it provides means by which a part is enabled to keep pace with the increased demands which are made upon it, when either its own proper work is increased or when some other part has become defective, so that, by its aid, the struggle for existence can still be maintained under circumstances which would otherwise in many cases soon lead to a fatal result. It is therefore in the highest degree protective or conservative. Thus such muscular organs as the heart, the large intestine, and the bladder grow stronger when they encounter abnormal resistance in the expulsion of their contents; and, to give minor instances, the epidermis thickens over points of increased pressure, and the cremaster grows into large fleshy bundles in cases in which heavy tumours, such as hydrocele or irreducible hernia, are present in the scrotum. And here is a photograph of one of Professor Sandow's pupils, whose muscles in the discharge of increased functions have grown to this extraordinary size (Fig. 2).

The growth of the uterus during pregnancy is yet another remarkable example. And, in the second place, organs grow when vicarious functions devolve on them. For instance, one kidney, the other being absent or useless, may acquire twice its normal bulk. When one limb is paralysed or lost in a young subject, the opposite grows to an abnormal size, and

thus acquires abnormal strength. Note, again, the growth till then latent, which takes place in the arteries of a limb when, after occlusion of the main trunk, a collateral circulation is established; or in superficial veins when the deep veins have become obstructed. And it is chiefly those organs and parts which are most essential to the welfare of the individual which manifest the largest share of this reserve power of growth; for instance, in such indispensable organs as the heart or a remaining kidney the capacity for latent growth is much more considerable than it is in parts which are not essential to the maintenance of life, such as a hand or a foot; while the capacity for growth which the blood vessels hold in reserve, in order to maintain the vitality of a limb under very exceptional conditions was recently illustrated by a case⁴ recorded by Sir Thomas Smith, in which, when the femoral artery and vein, the profunda artery and vein, and the popliteal artery and vein were all simultaneously ligatured during the removal of a tumour in which they were involved, the enlargement by growth of collateral channels, both arterial and venous, was so considerable and so rapid that no serious embarrassment of the circulation occurred. Even in bone this reserve power of growth is very remarkable. Here is a skull the holding capacity of which is at least three times the normal amount. It is from a case of chronic hydrocephalus. Growth took place in order to avert intracranial pressure, which, without it, must soon have led to a fatal issue.

Fig. 2.—The shoulder and arm muscles of one of Professor Sandow's pupils.

I have entered upon these general, and I am afraid too fragmentary, remarks in respect to growth in order to introduce the immediate subject of the present lecture. In what follows I propose to direct attention to the two forms of growth which I have mentioned—that by which the stature of the individual is originally attained, and that which, till the occasion arises, remains in reserve—in their relation to surgery, with the object of showing that they are agents of considerable potency, which, according to the conditions under which they act, may either lead to deformity or conduce in an important degree to its removal.

Growth producing deformity is met with when the influence of the law of conformity of type, which ought to regulate the process, is interrupted or inhibited by some other influence of a stronger kind. This state of things may occur under a variety of circumstances, and many examples can be easily named: I have already referred to cases of what has been termed congenital hypertrophy, in which parts grow in excess because they are endowed, not with the usual, but with an abnormal, amount of germinal power or formative capacity. Deformity from overgrowth, depending on an abnormal blood supply, is seen in those not very rare cases in which, owing to tuberculous disease of the knee, the growing ends of the femur and tibia are rendered more than normally vascular, and the limb becomes an inch, or, as I have seen

it, two inches, longer than its fellow. The same result may follow inflammation of the shaft of a long bone, for instance of the tibia. Thus in chronic tuberculous osteitis, or in acute infective inflammation followed by necrosis, and attended with abnormal vascularity, continued over a long period, the bone may become as much as an inch and a-half longer than the opposite. But the most remarkable example of deformity due to abnormal growth depending on increased blood supply is furnished by Hunter's admirably ingenious experiment, in which the spur of a cock, which he transplanted from the leg into the much more vascular comb, grew to be six inches long (Fig. 3). The

Fig. 3.—The head of a Maltese cock, into the comb of which a spur had been transplanted from the foot, and had grown into the spiral horn six inches long, shown in the drawing. Preparation in the Museum of the Royal College of Surgeons.

production of deformity when normal growth takes place under abnormal circumstances, is further illustrated in cases in which an injury of the lower end of the radius⁵ has been followed by arrested growth of the bone, and as normal growth has continued in the ulna, the hand has been pushed over to the radial side. A similar result is met with in the lower limb, and many examples have been recorded. Two years ago I saw a young married lady whose left tibia was $\frac{3}{4}$ inch shorter than the right, and curved inwards just above the internal malleolus. The fibula was disproportionately long, so that its lower end, as the patient stood, was nearly 1 inch nearer the ground than its fellow. The external malleolus was prominent, and the foot was considerably turned in. The history of this case was that when the patient was 6 years old she injured her ankle by a fall—no doubt, as the result showed, separating the lower epiphyses of the tibia. Splints were applied, and in a few weeks she was walking naturally. Deformity, however, was slowly developed, owing to continued growth of the fibula, while growth of the lower end of the tibia had been arrested by the injury, which had involved the epiphysal line. I removed $\frac{3}{4}$ inch of the fibula just above the malleolus and divided the tibia, and was then able to place the foot in its normal position. Some years ago I met with a case of deformity of the lower jaw in a boy aged 9. The teeth, instead of being vertical were nearly horizontal, for during growth the jaw had been constantly drawn upon by a strong scar following a burn in the neck, and in which the bone was firmly imbedded. Further examples of growth resulting in deformity with which all are familiar are met with in the spine and in the tarsal bones. In the spine in scoliosis developing in early life growth is checked by increased pressure in the concavity, and favoured by diminished pressure in the convexity of the curves, so that the bodies of the vertebræ become wedge-shaped. In this specimen the vertical measurement of the bodies is in the concavity $\frac{1}{2}$ inch against $1\frac{1}{2}$ inch in the convexity of the curve. In Pott's disease, when

several of the vertebral bodies have been destroyed, the thorax as growth proceeds becomes deformed, for the ribs, jammed together at their posterior ends and so becoming fixed, grow almost straight forwards, and lead to extreme protrusion of the sternum. In talipes equino-varus, when this deformity is left without treatment, the tarsal bones are altered widely from their original shape. In the knee, again, in tuberculous disease, if the joint has been long considerably flexed, or, as I have seen in a case of septic arthritis followed by fibrous ankylosis in a position of flexion, the front part of the condyles of the femur and of the head of the tibia grows in excess; and this change of shape may be so marked that it so far locks the bones as to become a considerable element of difficulty in the reduction of the deformity; while even if the deformity could be reduced, the articular surfaces would be found so changed in shape that they would no longer admit of accurate coaptation in the extended position of the limb.

Growth, although it is in itself and in its surroundings perfectly normal, may produce a gradually-increasing deformity. This is seen in the instance of scars. Thus, vaccination scars produced in infancy grow to scale with the limb on which they are placed, and become three or four times their original size. The growth of a scar I lately saw illustrated in a distressing manner in a case in which a nævus on the face had during infancy been treated by vaccinating its margin at four different points. The inoculations had all "taken," with the result that when the patient was a girl of 18 she presented a group of four scars on her cheek, each of which was nearly as large as a threepenny-piece. As a further illustration of the growth of scars I may mention a case in which, some twenty years ago, I treated a cutaneous nævus on the face of an infant 2 months old with strong nitric acid. The nævus was nearly round, and not more than a quarter of an inch in diameter. As this patient grew the scar grew also, and, a few weeks ago, I found, on examining it closely, that—though being quite flat and smooth and of the same colour as the surrounding skin, it hardly shows—it is seven-eighths of an inch in diameter, so that, growing to scale with the face, it has become nearly four times its original size.

I now pass on to consider growth as an agent not in the production, but in the removal of deformity. I have shown that in many cases when growth produces deformity, it does so under the influence of some condition which has supplanted or inhibited the action of the law of conformity to type. For instance, when growth continues in the ulna, after it has been arrested in the radius, in consequence of a fracture or other injury involving the epiphysal line at the lower end of this bone; or when the lower jaw has during growth been diverted from its typical shape by the dragging action of a scar in the neck. In such cases, however, although the law of conformity to type is prevented from asserting itself by some more powerful adverse condition, it is merely suspended, but not abrogated; and is ready to resume its influence as soon as the agency which has opposed it is removed. It thus becomes a powerful corrective agent which surgery may often turn to very useful account.

RICKETS.

The following case, which occurred in 1874, affords a good illustration. The bones of the lower extremities of a rachitic child, aged 7, had during infancy yielded under the weight of the body, and became so deformed that I determined to operate on the tibiæ, and then on the femora. To straighten the tibiæ, which were sharply curved forward in their lowermost third, I performed osteotomy, first on one side, and then, after an interval of some months, on the other, and removed a wedge-shaped piece of bone from each. In those days aseptic surgery had not been fully established. Both wounds suppurated, and from both tibiæ small scales of necrosed bone slowly separated. Under these circumstances twelve months passed, and during the greater part of this period the child was kept off his feet. Then when the femora were examined, it was found that in the course of these twelve months during which the weight of the body had been removed they had, under the undisturbed influence of the law of conformity to type, regained their normal shape, and that any resort to osteotomy was entirely unnecessary.

Again, it is frequently observed that the limbs of rickety

children which for a time are markedly deformed subsequently "grow straight." I have seen a considerable bend disappear in the course of nine months. In such instances the original deformity takes place because, when the bones are deficient in rigidity, the weight of the body is a stronger influence than the law of conformity to type. But when, a year or two later, the rickety state has become less marked or has entirely ceased, and when the bones have regained their natural strength, they gradually, without surgical interference, grow straight. But sometimes the rickety condition is present in a more aggravated degree. In such cases the bones under the weight of the body bend easily, and continue to do so for an extended period. And not only so, but growth itself is now an agent to increase their distortion; the bones, in fact, "grow bent." Thus thwarted in its normal course, the law of type is found to assert itself in another direction. Unable, under the circumstances, to restore a curved tibia to its normal shape by ordinary growth, it calls forth the other form of growth I have mentioned—that which is latent or in reserve—and by its means develops a buttress in the concavity of the curve, which effectually prevents any further bend—in other words, any further divergence from the normal type.

Fig. 4.—A rickety femur, showing a strong buttress in the concavity of the curve.

This reserve power of growth is present, it is interesting to notice, in a remarkable degree in plants.⁶ Thus Hegler found that "the hypocotyl of a seedling sunflower, which would have been ruptured by a weight of 160 g., bore a weight of 250 g. after having been subjected for two days to a strain of a weight of 150 g. The weight was subsequently increased to 400 g. without injury. Leaf stalks of *helleborus niger*, which broke with a weight of 400 g., were able to resist one of 3.5 kilog. after having been subjected to a strain for about five days." This increased strength was acquired by the rapid growth of the mechanical tissues of the plant in the form of supporting buttresses.

It must, however, be observed that although such a buttress (Fig. 4) prevents further deformity, it also prevents, or materially limits, the correction of the deformity by future growth, for

it acts as a strong tie beam, and often renders the curvature to a great extent permanent. These considerations have an important bearing on the treatment of deformities due to rickets. In their early stage these deformities can be easily limited or removed by treatment, by which the rickety state is corrected and by keeping the patient much off his feet, so that instead of allowing the weight of the body to supersede the influence of the law of conformity to type, this latter force shall exercise its normal influence on the result of growth. In the case I have mentioned above the thigh bones under such favourable circumstances, although the deformity was very marked and tie-beam buttresses were present, grew straight within the year. A similar result was met with in the following instance. A child, aged 3, presented rickety deformity to an extreme degree, involving both the upper and lower extremities, all the bones of which were short, large at their articular ends, and so curved that a single, or perhaps a double, osteotomy of each would have failed to put them straight. The patient was placed in the country, fed on milk and fresh meat, ordered cod-liver oil, and kept entirely off his feet for eighteen months. At the end of this time his bones were practically straight, and he was allowed to be on his feet. The patient is now 21; the deformity has disappeared. He is a groom, and, I am informed, is "very well and active."

These cases will suffice to show that in the early stage before a buttress, acting as a tie-beam, has been developed, rickety deformity will be corrected by growth, if its normal course is secured by favourable circumstances. They show also that deformity is removed much more rapidly than is, I think, generally recognised. On the other hand, however, when buttresses have once been developed, though even then growth will usually tend to straighten the bones, its effect in this direction is greatly curtailed.

FRACTURES.

Fractures in the young not rarely escape detection, chiefly because many of them are of the green-stick—that is incomplete—variety. Under such circumstances union is apt to be attended with considerable deformity. This is the case especially in the clavicle, but the same result is met with in green-stick fractures in the forearm, the thigh, and other parts. Now if these cases are kept under observation for twelve or eighteen months it is found that as growth proceeds deformity, even though it was at first very considerable, gradually becomes less, and at length entirely disappears. Although fracture of the clavicle is a frequent accident in early childhood, and although union often occurs with the fragments in a bad position, it will be allowed, I think, that it would be difficult to find in children of six or eight, one in whom any evidence remained to show that the accident had ever taken place. At the Children's Hospital, where I was formerly surgeon, in dealing with green-stick fractures of the forearm united with deformity, I used at first to straighten the bones under chloroform, but afterwards I found that this method, unless the distortion was severe, was unnecessary; for the deformity became imperceptible in the course of six or nine months. But I am able to relate a case which offers conclusive evidence that after fracture in young subjects attended with deformity, as growth proceeds under the influence of conformity to type, the bone concerned may completely regain its normal shape. Some years ago I performed Macewen's osteotomy in both femora for genu valgum in a child 7 years old. I have no record of the case at the time of the operation, but as I never resort to this proceeding in any but somewhat severe examples of knock-knee, the bones must after the operation have been put in a position of considerable deformity. The child died three years later of measles, and this is one of the thigh bones (Fig. 5). It is of normal shape, and bears no trace whatever of the deformity which was produced by the osteotomy. Its longitudinal section, however, shows at the situation indicated by a small cross, a projection of the wall of the shaft into the medullary canal, and this no doubts marks the point at which the bone is divided. This point, it is interesting to observe, is situated an inch and a half above the proper level for an osteotomy for knock-knee, a circumstance readily explained by the growth which took place in the lower end of the shaft

between the operation and the death of the patient three years afterwards.

Fig. 5.—The left femur of a child aged 7, on whom double osteotomy (Macewen's) for genu valgum was performed three years previously. The bone is markedly rickety, with an antero-posterior curve in the middle of the shaft. Before a section was made the seat of the fracture could not be ascertained. Looked at in front the shaft is perfectly straight. At a spot indicated by a small cross on the longitudinal sections, a faint trace of the fracture is still visible, namely, slight thickening of the cancellous tissue, especially towards the posterior aspect of the bone (Museum, St. Bartholomew's Hospital, No. 807 G).

This complete disappearance of the deformity is very remarkable, and at first sight may seem scarcely credible. Yet is it not a result that we might, on reflection, be almost led to expect when we remember what takes place in the repair of fractures in adult bones in which the plastic agency of growth is absent? Here, for example (Fig. 6), is an adult femur in which a fracture has united with the fragments considerably overlapping each other. The specimen shows how strictly the process of repair has been governed by the law of conformity to type. The ends of the fracture, which were at first open, have been closed by the formation of bone exactly similar to that which forms the external layer of the normal shaft. All the interior callus has been removed so that the medullary canal has been restored, and the exterior callus, which was at first present, has also been absorbed, and all projections and irregularities of the surface have been removed, while the callus which remains to connect the overlapping ends, and which is about 2 inches in length, has acquired compact walls and a medullary cavity, so that it precisely resembles the type on which the original shaft was modelled. The imitation of type is throughout complete. Need we wonder, when we see how closely, even in the adult, bone repair conforms to the original structure concerned, that, when the plastic element of growth is also present, conformity to type is well-nigh absolute?

THE SPINE.

The influence of growth in the removal of deformity is displayed in the spine in a degree which deserves particular notice. Scoliosis, so common in its slighter degrees in the young, usually begins between the ages of 7 or 8 and 12 or 14, but it may commence in quite early infancy as the result of

rickets, and, needless to say, unless proper measures are adopted, the condition tends to increase and become permanent; for here, as in the case of the long bones, the weight of the trunk acting on parts diminished in rigidity becomes a stronger influence in controlling the results of growth than the influence of conformity to type, so that the normal shape of the column is exchanged for one of deformity. But those who have paid much attention to the subject are aware that in a large proportion of cases of scoliosis in children between 8 and 14 the curvature in the course of two or three years will completely disappear under favourable conditions when good air and diet are secured, when overwork and constrained positions at school are avoided, and when an increased period of horizontal rest is provided (11 or 12 hours at night, with an hour in the morning and an hour in the afternoon). When, with the accessories I have mentioned, the spine is relieved by posture of the trunk for about half of the 24 hours, the result of growth is soon apparent—the spine, like a rickety femur or tibia, “grows straight.” The spine, however, differs from a long bone in an important respect. The shaft of a long bone being all in one piece is possessed of inherent stability, and is not, in its normal condition, dependent on external support; while the spine, formed as it is of separate elements superimposed on each other, depends for support on the muscles and ligaments by which it is surrounded. While, therefore, the spine tends under the influence of conformity to type to grow straight when it is relieved of the weight of the trunk, yet as the surrounding muscles are weak the column yields as soon as the patient assumes the upright position.

Fig. 6.—Overlapped fracture of a femur, in which the ends have become closed by compact bone, and the callus by which the fragments are united has been modelled on the type of the original shaft (Museum St. Bartholomew's Hospital, Series III, No. 98).

Another element, therefore, in treatment must consist in the development and strengthening of the muscles. This at the present day is secured by massage and exercises judiciously employed. The idea entertained by the public and by some practitioners that scoliosis is always serious and intractable is to a great degree a misconception, which has

arisen from the traditions of former years, when the affection was either neglected or treated with heavy steel instruments in which the trunk was encased, and which were screwed up twice a week, or on ordinary commercial principles every other day. At the present day few surgeons so far ignore the rudiments of physiology as to apply steel instruments to push a scoliosis straight, as if they were pushing up the dead weight of a leaning wall, for it is clearly seen that the attempt to correct not only lateral deviation but rotation of the vertebrae by pressing in a horizontal direction upon the series of elastic arches formed by the ribs, always has been and always must remain a mechanical absurdity. Formerly the result of this treatment was to aggravate scoliosis to a very grave extent; and now that it has been discarded, the proportion of cases which become severe is greatly diminished. Indeed, clinical experience shows that in a large majority of instances scoliosis is not prone to attain any serious degree; and that when the treatment I have so briefly mentioned is adopted early, and is efficiently used, a very large proportion (70 or 80 per cent.) end in complete recovery. In those cases in which infants, as the result of rickets, develop scoliosis during the first or second year, the prognosis is very unfavourable, for during the long period in which growth will continue, deformity is likely by degrees to become extreme, and there can, I think, be little doubt that the grossly distorted skeletons which are to be found in every large museum are from subjects whose curvature began during early childhood when they were suffering from rickets. No cases, I believe, more graphically illustrate the agency of growth in the production of deformity. The treatment of infantile scoliosis is attended with great difficulty. Instrumental support cannot, it must be allowed even by those who approve of it in older patients, be applied when the abdomen is protuberant, the pelvis much smaller than the thorax, and the skin unfit to bear even slight pressure; nor can exercises be employed with good effect. The case, however, is one in which growth as an agent in the removal of deformity may be turned to very good account. The spine must be placed under conditions in which it shall be able to "grow straight." The infant must, therefore, be kept very much in the horizontal position, and appropriately gentle massage must be persistently employed to develop and promote nutrition of the spinal and other muscles. A case in illustration may be briefly related.

A rickety infant, aged 13 months, a patient of Dr. McCaskie, of Brompton, had lateral curvature with very marked rotation. The child's muscles were so weak that when he was placed in a sitting position the curvature was considerably increased. The deformity, which had been noticed for three months, was rapidly getting worse. It was evident that, if matters were allowed to go on, in the course of a year or two, or even a few months, the deformity would become extremely grave, for the bodies of the vertebrae were abnormally soft, and by being compressed in the concavity of the curves would soon become wedge-shaped and otherwise distorted. The patient was therefore kept for nine months almost constantly in the horizontal position in a small wicker-work tray, in which he could be carried about, and gentle massage for the trunk and limb muscles was regularly employed. During this time the child remained in very good health and grew quickly, and at the end of the time the deformity, which had grown less and less marked, had so nearly disappeared that scarcely a trace of it remained. He was now allowed to sit up a little and be on his feet each day for a gradually extended period. He soon learned to walk, and when he was 3 years old he was running about in perfectly good health and strength, and with no return of his curvature. He is now a strong active boy of 11, and his spine is quite normal. The long period during which the horizontal posture was maintained was, of course, a troublesome ordeal for the child's parents. For himself, he always took it in very good part. But I would ask, By what other means could the development of an aggravated scoliosis have been prevented?

JOINTS.

In tuberculous disease of the knee the joint is often considerably flexed, the head of the tibia is displaced backwards and outwards, and the leg is rotated outwards. This three-fold deformity is, as every surgeon knows, one which

it is very difficult to correct. An attempt to straighten the limb by orthopaedic apparatus involves, [as I have pointed out elsewhere,] the use of leverage, and two objectionable results unless great care is taken are apt to follow. The head of the tibia may be forced still further backwards, and owing to the production of interosseous pressure the disease may be aggravated, and the joint rendered painful.

In cases of this description, provided the displacement is only moderate in degree, if the joint is enclosed in accurately-modelled leather splints, and if the patient is not allowed to walk on the limb, as time goes on the joint will "grow straight." In 1886 I saw a child, aged 14 months, a patient of Dr. Buck, of Clapton, with acute tuberculous disease of the knee, attended with considerable swelling, heat, pain, and relaxation of the ligaments. The joint was flexed at an angle of about 100 degrees, and the head of the tibia was displaced backwards and outwards to an extent that I have never seen corrected by any kind of apparatus. In so young a child it would have been a very difficult matter to use any apparatus for extension. The knee was therefore fitted with leather splints and kept at rest. These splints were uninterruptedly worn for the next four years. The joint gradually became more and more extended, the splints being remodelled about every six months to correspond with the improved position of the limb, and by the end of four years all trace of the displacement had disappeared, and the joint was in a position of complete extension, nor were its movements in the least degree impaired. At the present time the joint is still sound, quite freely movable, and of normal shape. The only remaining defects are some muscular wasting and two inches of shortening due no doubt to damage done in the course of the disease to the growing ends of the femur and tibia. In the later stages of a disease of this kind the limb may with great advantage be placed in a Thomas's knee splint. In this excellent appliance a broad webbing strap may be passed over the front of the thigh just over the knee, and another over the upper part of the tibia, and be so connected with the lateral bars of the splint that they assist in straightening the limb. By this means, without the production of any injurious interosseous pressure, the process of extension of the knee may be materially promoted, while the patient can move about on crutches and a patten on the other foot. It must be clearly understood that I do not suggest this method for the removal of deformity of the knee in the usual run of cases, when more active measures can be successfully employed. I have, however, from time to time met with cases in which it has led to results that could not, so far as I am aware, have been secured in any other way, and it is in exceptional cases that exceptional methods are often of the greatest value.

One further example may be related. A boy, aged 3, had tuberculous disease of the knee-joint, with an abscess in the tibia which reduced the outer part of the head of the bone to little more than a shell, with the result that not only was the knee flexed, but the tibia was deflected so that a severe degree of knock-knee was developed. The distortion, when I first saw the case, was such that there seemed no probability of straightening the limb except by excision—a very undesirable proceeding in so young a child. The joint was, however, with the double purpose of arresting the tuberculous process and preventing further deformity, put up in leather splints in anticipation that at some future period excision would be done. In the next year so great an improvement in the position of the limb occurred that splints of modified shape were applied on two occasions to keep pace with the altered shape of the joint, and a year later still the deformity had so far disappeared that the joint was almost fully extended and only a trifling degree of genu valgum remained. The boy, after wearing the splints for another year, had a movable joint and a limb which was practically free from deformity, except that it was three-quarters of an inch shorter than its fellow. On this limb he could walk with scarcely any appreciable defect. I feel sure that had excision been performed when the case first came under observation, or even at a later period, the result would have been very much less favourable.

I hope the observations which I have laid before you will be sufficient to show, as I proposed at the outset, that growth

is an agent, the tendencies and results of which must be fully taken into account in the clinical surgery of the young. This necessity arises from the fact that in the instances I have adduced, and in others that could be mentioned, various deformities are developed because conditions have arisen which have interfered with or superseded the law of conformity to type, under the regulating influence of which growth normally takes place. When this is recognised it at once becomes apparent that treatment, if it is to be successful, must be guided by the principle of removing as far as possible the disturbing influence, whatever it may be, that is at work, and substituting for it those conditions by which normal growth shall be promoted to the greatest available extent. Nor, I think, can any principle be named which rests on a sounder foundation, or which in the true sense of the term is more directly conservative. Many forms of treatment which are properly named conservative are conservative in only a limited or qualified degree, for, although under the circumstances they undoubtedly secure the safety or promote the advantage of the individual concerned, they do so only by the sacrifice of parts or organs, the loss of which involves more or less serious disability or inconvenience; while in the cases to which I have been alluding the surgeon, recognising in growth one of the most strongly operative of all the manifestations of life, and observing the marvellous results which are produced by its agency, makes it his endeavour to ascertain the conditions which are most favourable to it, and does his best to secure their full effect. It would not be difficult to mention plans of treatment which have done violence to Nature and her methods, and which have inevitably led to failure and disappointment, for *Nature non imperatur nisi parendo*. But the subject I have discussed may suffice to remind us of one of the chief rules of practice, that under which the surgeon makes it his object to act as the strict disciple of Nature, and renders himself ancillary to her work.

NOTES AND REFERENCES.

¹ *Origin of Plant Structures*, by the Rev. George Henslow. 1895, p. 136.
² *Loc. cit.*, p. 190. ³ *Loc. cit.*, p. 180. ⁴ *St. Bartholomew's Hospital Reports*, vol. xxx, p. 223. ⁵ This subject has been fully discussed by Mr. Jonathan Hutchinson, jun., in his able and instructive lectures at the College of Surgeons on Injuries to the Epiphyses and their Results. ⁶ Henslow, *loc. cit.*, p. 204. ⁷ *Diseases of the Joints and Spine*, 1895, pp. 272 et seq.

THE CROONIAN LECTURES

ON THE

CHEMICAL PRODUCTS OF PATHOGENIC BACTERIA
CONSIDERED WITH SPECIAL REFERENCE
TO ENTERIC FEVER.*Delivered before the Royal College of Physicians of London,*

By SIDNEY MARTIN, M.D., F.R.S., F.R.C.P.,

Professor of Pathology, University College; Assistant-Physician to University College Hospital and to the Hospital for Consumption, Brompton.

LECTURE III.

THE PRODUCTION OF THE POISONS OF THE TYPHOID BACILLUS
IN ARTIFICIAL CULTURE OF THE MEDIA.

Two different kinds of media were employed in which to grow the bacillus. Ordinary peptone broth was used to determine the presence or not of secretory products of the bacillus; and broth containing no peptone but mixed with some proteid solution—either blood serum, alkali albumin made from the spleen, or an extract of lymphatic gland—in order to determine the presence of digested products of the bacillus. In some cases, liquid blood serum sterilised at 60° C. was used without the addition of broth. The virulent bacillus grows well in ordinary peptone broth, but the growth in solutions containing much proteid is very much less exuberant. This is especially the case with sterile blood serum, and most commonly the bacilli, after growing for a time, sink to the bottom in clumps and cease growing, although they are still alive and active. It grows much better in a solution containing a small proportion of alkali albumin, but still not nearly so well as it does in broth. The object of growing them in the presence of proteids was to test whether there

was any digestive action of the bacillus. In one experiment, two test tubes were filled about 3 inches deep with liquid blood serum, which was rendered semi-solid by heating at a temperature of 65° to 70° C. The medium was translucent, and one of the tubes was inoculated with the bacillus, the other being kept as a control. It was observed that the bacillus rapidly grew in the medium, spreading through all parts of it, but that no liquefaction indicative of digestion occurred during incubation, which lasted 102 days. When grown in a liquid medium, however, containing a small proportion of alkali albumin, and especially alkali albumin made from splenic pulp, the virulent bacillus had a slight digestive action, although a very insignificant one. Thus in one experiment, after the bacillus had grown thirty-two days in such a medium, it was found that the solution gave a distinct biuret reaction with copper sulphate and potash, and that the alkali albumin was beginning to be changed into albumoses. This digestive action, although slight, is explanatory of the albumoses found in the spleen in cases of typhoid fever.

POISON FORMED IN PROTEID SOLUTIONS.

If the non-virulent bacillus is grown in a solution containing digestible proteid instead of peptone, a poison is formed which has an action like that of the poison formed in broth. Thus, in one experiment the bacillus was grown in a 1 per cent. solution of serum alkali albumin in broth for twenty-nine days, at the end of which time it was filtered and the filtrate concentrated, and the proteid separated by alcohol. Some amount of digestion had taken place, as the solution gave a biuret reaction. The solid product obtained was injected in two doses (about 0.3 g.) on successive days into the marginal vein of the ear of a rabbit. Following each injection there was a marked lowering of the temperature and a subsequent rise, but not to the average normal temperature of the animal for some days. Some bodily depression was observed on the first two days of the experiment, and there was a loss of body weight, but there was no diarrhoea; and the animal subsequently completely recovered.

In another experiment the non-virulent bacillus was grown in a solution of alkali albumin made from the splenic pulp, but without adding broth. One flask was filtered after forty-five days, and 10 c.cm. of the filtrate were injected in two doses on successive days into the marginal vein. After the first injection there was only a very slight rise of temperature, which, however, was more marked after the second injection. The animal showed no particular symptoms except a gradual loss of weight, which amounted to 235 g. at death on the twenty-eighth day of experiment. The *post-mortem* examination revealed no naked-eye change in any of the organs or tissues of the body. No cultivations were obtained from the heart's blood; the heart muscle showed no fatty degeneration, and there was no degeneration of the peripheral nerves. Another culture was filtered after fifty-nine days' incubation, and 12 c.cm. were given in two doses on successive days, also into the marginal vein. There was a well-marked rise of temperature after each injection but no diarrhoea was produced, and there was only a slight loss of weight, which was recovered from. The animal died in twenty-four days, and no cultivations were obtained from the heart's blood, liver, spleen, or peritoneal cavity. There was subcutaneous hæmorrhage in patches in front of the thighs and the caudal region, and the peritoneum contained about 1 cm. of clear fluid. There was no peritonitis, and the organ showed no naked-eye change.

The results of growing the non-virulent bacillus in a proteid solution show that the poisons excreted are of the same nature as that formed by the virulent bacillus in broth. The digestion which takes place in the proteid solution is extremely slight, and the toxic body is thus mainly an excretion of the micro-organism.

Although the toxic products are developed in the presence of digestible proteids, yet they are most evident when the bacillus is grown in peptone broth. In one experiment the broth was inoculated with the typhoid bacillus obtained from the spleen of a case of enteric fever, but before it had been made virulent by passage through animals. The growth was allowed to continue for eleven days, and the broth was filtered through a Chamberland filter. Two injections into the marginal vein were made on successive days.