few were larger, being more than twice the size of lymphocytes. Whilst most of the cells were round, or nearly so, some, especially the larger ones, were oval, or slightly irregular in shape. In no place were they densely packed together, and in many parts of the tumour they were very scanty indeed. There was an ill-defined granular matrix, and the blood vessels were numerous, relatively large, and very rudimentary. The photomicrograph shows one of these vessels surrounded by sarcomatous tissue. In no part of the tumour were the cells more numerous than is here represented, and in most parts they were present in much smaller numbers. Tissue of this typical sarcomatous type formed a comparatively small part of the tumour as a whole. A great part was formed by a loose sarcomatous tissue infiltrated with blood and blood clot. In many places it was mainly composed of red blood corpuscles with little or no fibrin, but many young organising blood vessels shooting through it, and only a scanty sprinkling here and there of sarcomatous cells. Other parts, forming a considerable bulk of the tumour, were composed of inflammatory tissue and thrombi, the inflammatory tissue being of a comparatively recent date, and in many places, at any rate, of a septic nature. Much of the inflammatory tissue was rich in fibrin, and the disposition of many leucocytes within its meshes presented in places a superficial resemblance to a myxomatous tissue. Careful search, however, failed to find a single myxomatous cell throughout the tumour.

The growth was, therefore, a round-celled sarcoma of great vascularity and rapid growth, in which frequent and repeated hæmorrhages had occurred, and in and around which low forms of septic inflammatory changes had arisen.

In this case we have the interesting phenomenon of the occurrence of benign polypi in one nostril and malignant in the other, and not only so, but with benignity and malignancy co-mingled in one and the same nostril. The results of the microscopical examinations, which were made at intervals of about a month, afford reasons for the opinion that the polypi in the left nostril were originally benign, and that the sarcomatous invasion or degeneration was a comparatively recent occurrence, and probably commenced in the deeper part at or about the time of the first hæmorrhage. For undoubtedly polypi must have been present for many months at least before that occurrence.

Cases have already been recorded in which a sarcomatous has been preceded by a benign growth, as in that by Onodi,¹ which differed from the present one in that the transformation occurred only after ablation of the polypi. Sendziak² met with a case of malignant sarcoma with benign polypus of the nose, the malignant being situated low down on the septum, and the benign in the middle meatus. Haring³ mentions the case of a woman, aged 26, in whom four months after removal of a polypus apparently of ordinarily benign character, the nasal cavity was blocked by a round-celled fibro-sarcomatous growth. Not only simple and sarcomatous growths coexist, but more than one variety of malignancy may be present. Plicque⁴ quotes Ricard regarding two cases where, along with ordinary mucoid polypi, not only sarcoma but also epithelioma was present, and the same writer mentions that similar cases have been recorded by Schaeffer, Hopmann, Terrier, and Voltolini.

Apart from the etiology of malignant disease generally, the point which seems worthy of some consideration is the question of (surgical) traumatism in determining or inducing malignancy in ordinary mucous polypi. The view is held by some, and certain of the cases now quoted would appear to bear this out, that instrumental or operative interference and irritation are necessary to induce malignant transformation in an ordinary simple polypus. I do not hold that opinion, for, as already stated, in the case now submitted, no operative treatment had been practised before I saw the patient, and at the first interview I expressed my opinion to the medical attendant that malignant disease was present.

In deciding upon the precise nature of these cases, we have seen, in the earlier stages of the disease at least, that the microscope cannot be trusted to reveal the full and true nature of the lesion. There seem to be two features of value in the diagnosis of malignant polypus of the nose in its earlier stage—namely, hæmorrhage, and the locality of the growth. The former, which may be spontaneous or easily provoked, appears to be far and away the most important symptom; it is usually severe and recurrent, and, as I have seen in other cases, generally occurs for the first time at an early period, but may be somewhat postponed, as in the present instance.

In estimating the diagnostic value of this symptom, it ought not to be forgotten that women, more rarely men, sometimes suffer from a benign form of bleeding growth of the septum, usually in the left nostril, and known as "bleeding polypus of the nose." This is of the nature of a lymphangioma, is easily removed, and does not usually recur; but with this exception we may safely assume that, given an elderly individual with

a nasal polypus, or a sessile polypoidal growth, which bleeds frequently and profusely, the probability is that it is malignant, more especially if the growth can be seen attached to the septum, for this appears to be the malignant area of the nose. One writer, indeed, goes so far as to affirm that "a polypus inserted on the septum must always be regarded as of a malignant character" (Plicque).⁵ In the later stages of the disease it may be impossible to say whence the growths have sprung, or whether they are or are not pedicled, and hence any diagnostic value which may be attached to these features is lost. On the whole, I am inclined to think hæmorrhage the most reliable test we have, and I would say that sessile polypoidal septal growths, or blood-stained or hæmorrhagic spots or areas on the surfaces of what appear to be ordinary mucous polypi, along with frequent and profuse bleedings, are almost conclusive proof of malignancy.

Whilst sarcoma of the nose is usually met with in elderly individuals, it is not exclusively so. I have not seen it occur under 40, but Bliss⁶ has recorded two cases of sarcoma of the nasal chambers in children, one aged 4 and the other 9 years. Nichols⁷ mention its occurrence in a boy (age not stated), and in another patient aged 17 years. A few similar cases have been recorded by other writers.

The prognosis as regards the results of operative treatment depends upon the degree of malignancy and the stage and locality of the disease. Carcinoma and epithelioma, which are fortunately of rare occurrence in the nose, are highly unfavourable. Fibro-sarcomata are much more favourable than pure sarcomata, and several cases have been recorded in which the former have not recurred after operation. Pröbsting and Cramer⁸ have even noted spontaneous shrinkage and healing of a naso-pharyngeal sarcoma, and Onodi is of opinion that the internal administration of arsenic tends to cause their disappearance. Long periods of immunity after operation have also been noticed by other observers in cases of fibro-sarcoma of the nose and naso-pharynx. Early cases of sarcoma which present as small sessile polypoidal growths on the septum are more favourable than those less frequent ones in which the ethmoid is primarily affected; and if the case be seen and the disease detected at a sufficiently early period, an intranasal operation by means of the curette and electro-cautery may suffice to permanently cure the patient. I have seen one such result in the case of a gentleman, aged 70, which I did not diagnose at the time, but which, in the light of an extended experience, I now believe to have been sarcoma of the septum.

I have to express my indebtedness to my friend, Dr. R. F. C. Leith, for his careful microscopical examinations of this and of other cases of malignant polypi of the nose which have been under my care.

REFERENCES.

- ¹ *Internat. Centralt. für Laryngologie*, vol. ix, p. 325. ² *Journal of Laryngology*, vol. x, p. 103. ³ *BRITISH MEDICAL JOURNAL*, February 20th, 1896. ⁴ *Journal of Laryngology*, vol. iv, p. 105. ⁵ *Annales des Mal. du Larynx*, March, 1890. ⁶ *Internat. Centralt. für Laryngologie*, vol. xi, p. 80. ⁷ *Trans. of the American Laryng. Assn.*, May, 1897. ⁸ *Internat. Centralt. für Laryngologie*, vol. xiii, p. 45.

CASE OF CEPHALIC TETANUS TREATED WITH ANTITETANIC SERUM: RECOVERY.

By G. E. HALE, M.B., B.C.,
Eton.

THE following case of tetanus treated with antitoxin, and followed by recovery, seems worthy of record. It was undoubtedly subacute, as thirteen days elapsed between the date of the injury and the onset of the earliest symptoms; but the comparatively favourable prognosis suggested thereby was discounted by the fact recorded by Gowers¹ that "all the cases (of cephalic tetanus) met with over 25 have been fatal." Moreover, he adds, "In all cases that have recovered, the tetanic spasm and facial palsy have very slowly passed away, the symptoms continuing altogether several weeks." In the following case the spasms were probably mitigated in intensity and lessened in number by the chloral and bromide given simultaneously; but in view of the rapid improvement commenced after the second large injection on the fifth day of the disease, and the complete disappearance of all symptoms on the twenty-first day, it seems only right to attribute

the successful issue of the case chiefly to the antitetanic serum.

H. R. T., male, aged 30, fell on the evening of January 12th last, and struck the bridge of the nose against the kerb of the pavement, causing an abrasion, after which the face is said to have become greatly swollen. He first sought advice on January 27th, complaining of an uncontrollable impulse to laugh, and of a "drawn" sensation of the left side of the face since the 25th. He had laughed, quite against his will, on the receipt of the news of the death of a relative. No limitation of the movements of the jaw could be made out, nor any objective symptom. The abrasion had healed. When next seen on the evening of Sunday, January 30th, eighteen days after the injury, there was slight but distinct trismus, slight ptosis of the left eye, and inability to close the eyelids completely; some right facial paralysis, and a painful feeling of stiffness round the waist. No cramps nor stiffness of the neck were complained of.

January 31st. Symptoms somewhat more marked; some difficulty in swallowing; slight wrinkling of forehead; a few cramps in legs; towards evening the neck became stiff; 10 c.cm. of antitoxin injected. A full dose could not be given, as through a misunderstanding only a prophylactic dose had been sent.

February 1st. Slept well. Trismus increased, and body become stiff: towards evening back of neck became quite rigid; 60 c.cm. of antitoxin injected.

February 2nd. Slept well, except for a few spasms of a not very severe type. By night the teeth could only just be separated, the abdomen became rigid, and there was distinct opisthotonos.

February 3rd. Sleep more disturbed; spasms more frequent and severe, the tongue being frequently bitten. Opisthotonos well marked, but the legs being unaffected, the arch was between the buttocks and the occiput. Teeth firmly clenched; fed by siphoning liquids from a baby's feeding bottle, the india-rubber tube being attached to a straw which was insinuated through the gap in the teeth in which he held his pipe. Spasms frequent throughout the day; patient in a very despondent state. In the afternoon 40 c.cm. of antitoxin were injected, and a mixture of chloral hydrate gr. xv with sod. brom. ʒj ordered to be given every four hours. By night there was slight relaxation of the trismus, and the spasms were noticed to be less frequent.

February 4th. Slept eight hours: abdomen and back less rigid. Patient in much better spirits; head can be moved slightly forwards; facial paralysis less marked. Fewer spasms (twenty-five in twelve hours).

February 5th. Trismus again more intense. Spasms fewer (nine in twelve hours), some severe, tongue being frequently bitten; 20 c.cm. of antitoxin injected. It was hoped to inject more, but Messrs. Burroughs and Wellcome's supply was exhausted on that day. One hour after injection the trismus relaxed sufficiently to admit the little finger as far as the middle joint.

February 6th. Slept well first part of night, but had frequent and sharp spasms from 4 A.M. till 8 A.M. Teeth could be separated 1/2 inch. Fifteen spasms during day; 40 c.cm. of antitoxin injected at 2 P.M. Abdomen in the evening less rigid, and opisthotonos less marked. Hypodermic injection of 1 gr. of morphine with 1/2 gr. of atropine at night.

February 7th. Slept seven hours in night with only two spasms. Spasm sufficiently relaxed to allow patient to rotate his head at will. Opisthotonos quite slight, but abdomen not greatly relaxed. Teeth could be separated sufficiently to allow patient to drink from a cup. Five spasms during the day; 20 c.cm. of antitoxin injected.

February 8th. Three spasms only throughout the day; very drowsy. Chloral mixture discontinued; catarrhal conjunctivitis of both eyes.

February 9th. No spasms; opisthotonos disappeared, movements of head free; 20 c.cm. of antitoxin injected.

February 10th. No spasms; able to eat minced chicken and mashed potatoes.

February 11th. Slept well, no spasms. Neck and back quite supple. Still some rigidity of abdomen, and trismus. Ptosis of left eye persists, and a slight degree of right facial paralysis.

February 12th. Sits up in a chair and eats solid food well, though the movements of the jaw are not yet absolutely free.

February 13th. All trace of trismus and rigidity of the jaw gone; facial paralysis and ptosis disappeared; able to walk about well.

On March 5th the patient went to the seaside, and on the 21st was able to return to work.

It is interesting to note the presence of ptosis, which is said to be very rare in cephalic tetanus, and the emotional disturbance accompanying the onset. Altogether 210 c.cm. of

serum were injected. The injections were made into the cellular tissue of the back by one or more punctures, 20 c.cm. being injected at each puncture. The usual antiseptic precautions were observed, no inflammation occurred, and but little pain was caused. The serum was obtained through Messrs. Burroughs and Wellcome from the Pasteur Institute at Paris.

REFERENCE.

¹ *Diseases of the Nervous System*, vol. ii, p. 631.

A CASE OF TRAUMATIC TETANUS TREATED WITH ANTITOXIN.

By G. BLACKER MORGAN, JUN., M.B.,
Sunderland.

H. M., aged 20, a farmer, was kicked over the right eye by a young unshod horse. A lacerated wound below and parallel to the right eyebrow was thus produced. The wound was seen some twelve hours later and thoroughly cleansed with carbolic lotion, and two sutures of carbolised horsehair put in. By the sixth day the wound was quite healed and the oedema of the eyelid had disappeared. On the seventh day he complained of some stiffness of the jaw. There was puffiness of the right side of the face simulating facial paralysis. On the morning of the eighth day, after a very restless night, he developed the typical sardonic smile, with tonic spasm of the muscles of the face and neck. He was given chloral hydrate 20 gr. every two hours, and was kept in bed in a darkened room; he was very restless all day, and had increasing difficulty in swallowing. Injected 8 c.cm. of antitoxic serum into the abdominal wall at 6 P.M. The temperature was 97.6° F., and the pulse 56. He was also given an enema of potassium bromide 3ss and chloral 25 gr., in milk 3iv, and this was directed to be repeated every two hours. At 10.30 P.M. it was reported that he had been quieter. There was no reaction from the antitoxin, and the temperature was still subnormal, pulse 60. He had a restless night, but slept some three hours. On the ninth day there was very little twitching, and grinding of the teeth occurred twice. He retained all the enemata but one, and took an ounce and a half of milk by mouth. The temperature was 98.4°, and the pulse 62. Injected 8 c.cm. of antitoxin at 11 A.M., passed a catheter and emptied the bladder, drawing off a copious secretion loaded with phosphates. At 6.30 P.M. the third injection of antitoxin, 8 c.cm., was given, and the bladder emptied. He did not retain the enemata. The temperature was 99.6° F. At 11 P.M. the spasms were very frequent and opisthotonos well marked. Gave him a hypodermic injection of chloral, gr.10, and restrained the spasms with chloroform.

On the tenth day, at 6.30 A.M., after a quiet night, he was much more conscious and answered distinctly. He was given another hypodermic injection of chloral during the night (gr.10). At 1.30 P.M. found he had got steadily worse since the early morning, and was quite unconscious; the conjunctivæ were insensitive, and tetanic spasms were nearly continuous, but could be controlled by chloroform. The pulse was 100, and the temperature 99.8° F. At 3 P.M. repeated the hypodermic injection of chloral (gr.10), and at 4 P.M. he was quieter. At 11 P.M. the pulse was 120, and the temperature 102° F. Cheyne-Stokes respiration was observed. Injected 16 c.cm. of antitoxin, and emptied the bladder with a catheter. He died at 2.30 A.M. on the eleventh day.

Remarks.—The colt which caused the injury was kept in a loose box in which a horse had died from tetanus three months previously. The loose box had not been disinfected in any way. The wound healed quite normally, and on the sixth day all swelling had subsided. The eyelid became swollen again on the ninth day, but there was no fluctuation to be made out over the wound. He had three injections of antitoxin, 8 c.cm. in each, in the first twenty-four hours, but, owing to the difficulty in obtaining a further supply, there was a delay of twenty-four hours before the fourth injection of 16 c.cm. could be administered. It is possible that the general improvement in his condition in the early morning of the tenth day might have been maintained had there been a further supply of antitoxin at hand. The injections caused no local disturbance nor pain.