REFERENCES.

¹ Koch, Some New Preparations of Tuberculin, *The Medical Week*, April 16th, 1897. ² Hirschfelder, Die Behandlung der Tuberculose und anderer infectiösen Krankheiten mit Oxytoxinen, *Deutsch. med. Woch.*, Therap., Beil., No. 4, 6 Mai, 1897. ³ Smirnow, Ueber das künstlich dargestellte Diphtherie Antitoxin, *Berl. klin. Woch.*, 1896, p. 597. ⁴ Bolton and Pease, The Production of Antitoxin by the Passage of Electricity through Diphtheria Cultures, *Journal of Experimental Medicine*, vol. i, No. 3, p. 537.

THE CROONIAN LECTURES

ON

THE MEANS BY WHICH THE TEMPERATURE
OF THE BODY IS MAINTAINED IN
HEALTH AND DISEASE.

Delivered before the Royal College of Physicians of London,

By W. HALE WHITE, M.D., F.R.C.P.LOND.,

Physician to and Lecturer on Pharmacology and Therapeutics at
Guy's Hospital.

LECTURE I.

[ABSTRACT.]

BEFORE studying pyrexia experimentally there are some pitfalls of which we must beware. In the first place, the normal range of temperature in the lower animals is wider than in man. I have very often taken the rectal temperature of rabbits, dogs, and monkeys, and in these animals the range is greater than in man. I have compiled from Dr. Pembrey's article on Animal Heat in Professor Schäfer's forthcoming *Physiology* the appended table (Table I.), which shows very well the range of temperature in different animals.

TABLE I.

Animal.	Average rectal temperature.	Extremes of observations.	Number of observations.
	Degrees C.	Degrees C.	
Horse	37.8	36.1—38.6	962
Cow	38.7	37.5—39.4	116
Sheep	40.4	38.5—41.8	304
Dog	39.0	37.1—39.9	250
Cat	38.7	37.9—39.7	41
Pig	39.2	38.7—40.8	269
Rabbit	38.9	37.0—40.8	110
Guinea-pig	37.9	37.0—39.2	19
Rat... ..	37.5	37.0—38.5	16
Mouse	37.4	37.1—38.6	27
Monkey (Rhesus) ...	38.4	36.9—39.7	22

The prolonged administration of an anæsthetic causes a fall of temperature. Kappeler found that after the administration of ether to human beings the bodily temperature sinks from 0.3° to 1.5° C. (the mean of twenty etherisations was 0.68° C.); but it rises again in a few hours to normal. Twenty-three observations were made on the effect of chloroform on human beings, and it was found that on the average the temperature fell 0.53° C. The fall did not begin till at least twenty minutes after commencing inhalation and was generally most marked after narcosis had passed off. In from twenty minutes to five hours the normal temperature was regained.

The performance of artificial respiration will by itself cause a considerable alteration of temperature. Dr. Fawcett and I in the course of some experiments we were conducting on the physiological action of β -tetrahydronaphthylamine made some observations upon the combined effect upon the temperature of etherisation and artificial respiration. Rabbits were used, the animal was kept fully under ether for two hours, and during this time it was fixed to a board in the usual way, and artificial respiration was carried on through an opening in the trachea.

The depression of temperature was very considerable; in one case 4.8° C. in $\frac{3}{4}$ hours, in another 2.2° C. in $\frac{2}{3}$ hours, and in a third 3.4° C. in $\frac{3}{4}$ hours. It could not be prevented by wrapping the animal in cotton-wool.

The fundamental problem in a study of pyrexia is to

discover whether or not the production of heat is increased, for it is clear that the temperature of the body may rise either because the loss of heat from it is less, while the production remains the same, or because the loss of heat remains the same while the production of heat is increased, or, if both vary, because the variations in the production have not been exactly counterbalanced by variations in the loss. I hope I shall, in these lectures, be able to bring forward evidence to show that sometimes pyrexia is due to an increased production of heat and sometimes to a diminution of the loss.

I should like to call attention to a point I have never seen discussed—namely, whether there is any evidence that pyrexia may be due to a variation in the specific heat of the body. It takes more heat units to warm to a given temperature a body with a high specific heat than it does to warm a body with a low specific heat to the same temperature, so that if the specific heat of the human body diminished while its production and loss of heat remained the same its temperature would rise. Most observers have estimated the specific heat of the body by calculation after death from that of the various tissues of the body. The lowest estimate given by this method is 0.75, the highest 0.9; but this is what might have been expected when we remember that a large proportion of the body is water. The properties of dead and living bodies are so unlike, and the specific heat of complex substances is often so difficult to foretell, that I thought it would be worth while to see if some means could be devised of estimating the specific heat of living animals. This would be easy enough in the case of cold-blooded animals, but the thermic mechanism of warm and cold-blooded animals is so different that it would be rash to assume that the specific heat of warm-blooded animals is the same as that of cold-blooded animals.

It struck me, however, that by using hibernating mammals it would be possible to calculate the specific heat of living animals not far removed from man. I employed dormice, for they are easy to obtain and to observe. When hibernating they lie tightly curled up in a ball, with their feet tucked away in the centre of the ball, and it is easy to take their temperature by placing the bulb of a thermometer in the centre of the ball. When hibernating their temperature rises and falls with that of the surrounding medium. Experiment showed, for instance, that when the temperature of the room was 37° F. that of the dormouse was 37.5°, and that when it was brought into a room at 52° its temperature slowly rose till it was 49.25°; it was then put back into the cold room and its temperature slowly fell. A large number of observations gave similar results.¹

It is clear that as the temperature of the dormouse rose and fell with that of the air, the number of degrees an equal weight of distilled water rose or fell divided by the number of degrees the temperature of the dormouse rose or fell in the same time will give the specific heat of the dormouse. I performed many experiments and found that, calculated in this way, the specific heat of the dormouse varies between 0.812 and 1.18, so that if we set it down at 0.95 we shall probably be very near the mark. It is noteworthy that when the thermometer stand was first brought into a warm room the earlier observations always gave a slightly lower specific heat for the dormouse than the later ones, because at first the rise of the temperature of the animal was, in comparison with the rise of that of the water, slightly the faster. The consequent slight increase in the creature's specific heat was often uniform. Thus in one experiment the first observation gave a specific heat of 0.857; according to the second it was 0.917; according to the third it was 0.933; and according to the fourth it was 1.0. Another noteworthy fact is that if the specific heat was calculated from observations made when the thermometer stand had been removed from a warm room into a cold one, it was always slightly higher than if calculated from observations made when the stand was brought from a cold into a warm room, and also the earlier observations gave the lower specific heat.

The close agreement of these results with those obtained by calculation from the dead tissues shows, I think, that we may safely put down the specific heat of dormice as being slightly below that of water. There are in these experiments several unavoidable fallacies, so that the specific heat obtained

by this method is only approximately accurate; but the fallacies lead to only a slight error. On the whole, the specific heat calculated by this method is slightly too low. Probably the results are more accurate than those obtained by calculating the specific heat of each of the tissues of the dead body and from these results estimating the specific heat of the body as a whole; the figures obtained by the two methods are in close accord, but by the method here given the specific heat appears higher than when calculated from the dead tissues.

I should not have introduced these figures had I not wished to show that whether we calculate from dead or living tissues the specific heat of the body is high, and it is obviously an advantage for warm-blooded animals, as they have to maintain a constant temperature, to have a high specific heat, for bodies with a low specific heat cool much more readily when the temperature of the surrounding medium falls than do those with a high specific heat; and although it would be unfair definitely to apply results obtained from hibernating dormice to a man suffering from fever, I think we may say that at present there is no evidence that in man the rise of temperature is due to a diminution of the specific heat of the body, as whatever the temperature of these dormice their specific heat varied very little, for reference to the paper from which I have quoted will show that experiments conducted for the same time usually gave nearly the same specific heat. Further, there is no reason to believe that the human body disobeys the law that the specific heat increases with the temperature, and, therefore, until the contrary is proved we must suppose that, other things being equal, it takes a few more calories of heat, and therefore requires a greater production of heat to raise the temperature of the body from 104° to 105° than it did to raise it from 98° to 99° .

COMPARISON OF THE VARIATIONS OF THE LOSS ON HEAT WITH VARIATIONS OF INTERNAL TEMPERATURE.

The communication which Dr. Waller made to the Physiological Society on November 11th, 1893, suggests the possibility of easily finding out whether in man the production of heat is increased when the temperature is raised, and of studying in detail the means employed to keep the temperature constant in health. It is clear that when the surface temperature of the body rises the loss of heat by radiation and conduction will be increased, and that when the amount of sweat secreted is increased the loss of heat from evaporation is increased, so that if in any case it can be shown that the internal temperature has risen, and at the same time the surface temperature has risen and the amount of sweat secreted has increased, the production of heat must have increased, provided that the temperature of the surrounding air is constant, and in the wards of a hospital it is kept nearly so.

I will now describe the means I have employed to obtain these data.

The Internal Temperature.—This is taken with an accurate self-registering clinical thermometer. When only slight variations are expected, it should be taken in the rectum or the axilla, for the rapidity of respiration and the temperature of the respired air affect the mouth; but when, as in many cases of fever, the temperature is raised several degrees above normal, an observation taken in the mouth will suffice, and if the observer is single-handed it is a great convenience to take it in the mouth, for the internal temperature, that of the surface, and the amount of sweat secreted should all be observed simultaneously.

The Surface Temperature.—This should be taken with a flat-bulbed thermometer held gently on the skin until the index is stationary; if it be held too firmly the pressure exercised will send up the mercury. The skin of the abdomen near the umbilicus is the best place to take the temperature, for the skin here is less liable than that elsewhere to transient variations of temperature from exposure to currents of air. The temperature registered by the thermometer does not represent that of the skin with absolute accuracy, for part of the bulb is exposed to the air, and the skin covered by the bulb slowly gets warmed because the bulb protects it, but as my object has always been to obtain comparative, not absolute,

results these errors are of no importance. I exposed the extensor surface of my forearm for some hours to the sun; the vessels consequently dilated, and the part was red: the surface thermometer showed the temperature at a spot well inside the margin of the red area to be 34.25° C., one-third of an inch outside the margin—the vascular dilatation here being just visible—it was 33.25° ; and some distance from the red patch the temperature was 32.5° . I have tried this experiment several times and the differences are always well shown. It is of interest to observe that even extreme vascular dilatation quite as great as is observed in any fever, only increased the surface temperature of the skin 1.75° C.

The Estimation of the Sweat.—This was estimated by using calcium chloride as an absorbent. It was fused into the bottom of a shallow circular glass box. When a reading is taken the box is held inverted on the skin for a given time (in the observations I have made I have, except in a few preliminary cases, always held it for ten minutes) and the difference in its weight before and after gives the amount of sweat secreted. It is best held with one finger so that the heat of the hand shall not affect it. The most suitable place at which to take an observation is on the skin near the umbilicus. To obtain accurate results the boxes must not be weighed directly after they are removed from the skin, for the glass has been warmed by the skin, consequently it displaces more air and warm air currents proceed from it while it is on the pan of the balance, and I always found that as it cools its weight slowly increases.

As the calcium chloride would otherwise absorb moisture from the air, each box is provided with an air-tight glass lid, which is taken off immediately before the glass box is put on the skin, and is put on again immediately the box is taken off. The box is always weighed with its lid on.

LECTURE II.

TEMPERATURE RAISED BY HOT BATH.

DR. HALE WHITE commenced his second lecture by relating certain experiments made on himself:

HOT BATH.

Experiment 1.—At 7 P.M. the temperature in the mouth was 98.2° F.; of the abdominal surface 93.7° ; and in ten minutes 2.4 milligrammes of sweat were yielded by the abdomen. At 7.25 a bath at 112° F. was entered; it was left in 14 minutes, when its temperature had sunk to 108° F. The observer felt faint, and was breathing deeply; at 7.45 the temperature in the mouth was 103.2° F.; of the abdomen, 94.5° F., and the quantity of sweat in milligrammes in 10 minutes was 185.7. At 8.5 P.M. the temperature in the mouth was 99.5° F.; at 9 P.M. it was 98.0° F. The abdominal surface temperature was 92.8° F., and the sweat was 1.5 milligramme.

Experiment 2.—At 3.30 P.M. the temperature in the mouth was 98.2° F.; of the surface, 92.3° F., and the sweat 1.5 milligramme. At 3.40 a bath at 112° F. was entered; it was left in 14 minutes, when its temperature had sunk to 106° F. Faintness and rapid breathing were not so marked. At 4 P.M. the temperature in the mouth was 101.0° F.; of the surface, 93.7° F., and the sweat in 10 minutes was 92.4 milligrammes. At 5 P.M. the temperature in the mouth was 98.2° F.; of the surface 93.7° F., and the sweat was 3.9 milligrammes.

TURKISH BATH.

Before the bath the temperature was 98.5° F., and the surface temperature of the abdomen 90.5° F.; milligrammes of sweat from the abdomen in 10 minutes, 2.2. One hour and 20 minutes was spent in the hot rooms, mostly at 120° F., but for some time at 150° F. and 190° F. Shampooing and a shower bath followed, and the surface was dried. The temperature in the mouth was then 99.8° F., of the surface 92.3° F., and the sweat in 10 minutes 46.1 milligrammes.

Commenting on these experiments Dr. Hale White said that the temperature rises in the hot bath owing to the fact that the hot water stops the loss of bodily heat by radiation and conduction, and prevents evaporation of sweat. Probably the hot water does not warm the body much in so short a time, as the specific heat of the body is high. In both these experiments I could, by watching the skin, see that the secretion of sweat did not begin to be very profuse till a few minutes after I was out of the bath, but by ten minutes after it was extraordinarily abundant. It will be seen that in all respects the first bath produced more marked effects than the second; probably this was because the second cooled the faster. But both experiments teach that the evaporation of sweat is by far the most important agent in reducing the bodily temperature when it has been artificially raised by a bath, and it will be noticed that when such a large amount of sweat is secreted the surface temperature is lower than it was either before the experiment began, immediately after the bath, or some time after the experiment, when the secretion of sweat is nearly normal. This is

no doubt due to the rapid evaporation of the sweat, and this means of reducing the temperature is so energetic that it actually makes the loss by radiation and conduction less. Other striking facts are the rapidity with which a normal temperature is regained and the enormous quantity of sweat secreted. The Turkish bath experiment also showed what a very energetic agent sweating is in reducing an artificially-raised temperature, for after sweating profusely for eighty minutes in the bath the sweat was still being secreted twenty times as fast as before the bath, but the quantity was not enough to make the surface temperature lower than before the bath.

TEMPERATURE RAISED BY DAMAGE TO THE BRAIN.

In two papers² I have shown that damage to the corpus striatum in rabbits raises their internal temperature. So many experimenters have confirmed this observation that I need not now trouble you with the details.

Many instances in which the temperature has been raised in man after a hæmorrhage into the corpus striatum have been recorded by Bourneville, Ferrier, J. H. Bryant, myself, and others, and references to these authors will be found in an address I gave at a meeting of the British Medical Association at Bristol in 1894.³ These cases show that if after a hæmorrhage takes place into one corpus striatum the temperature is raised, it will be higher in the opposite or paralysed axilla than in the axilla on the non-paralysed side, and that the temperature on the paralysed side remains higher than that on the other even during the subnormal period which succeeds the rise. I have seen several cases which illustrate this, and I will show you charts of two of them (see Charts 1 and 2), and since I began to investigate

Chart 1, showing the temperature after hæmorrhage into the corpus striatum on one side. The dotted line shows that in the axilla on the side opposite to the hæmorrhage; the straight line that on the same side as the hæmorrhage.

Chart 2.—From another but similar case to the preceding.

surface temperatures two more such cases have come under my care. They gave the following results:

Temperature in Two Cases of Left Hemiplegia Due to Right Cerebral Hæmorrhage.				Tempe- rature in Axilla.	Differ- ence in Axilla.	Surface Tempe- rature (Fore- arm.)	Milli- grammes of Sweat (Fore- arm).
Case III {	Paralysed side	...	...	100.8	+0.8	92.8	41.1
	Non-paralysed side	...	...	100.0	—	92.8	21.2
Case IV {	Paralysed side	...	...	97.8	+0.8	93.6	5.4
	Non-paralysed side	...	...	97.0	—	92.8	6.1

It will be seen that in both these cases the temperature was always higher in the paralysed axilla—that is to say, the internal temperature was higher on the paralysed than on the sound side. In the first case the surface temperature of the two arms was equal, but the amount of sweat secreted on the paralysed arm was much greater than on the other, therefore the loss of heat must have been more rapid on the paralysed side; but as in spite of this the internal temperature was raised, the quantity of heat in the paralysed arm must have been greater than that in the unparalysed arm. In the second case, on one occasion the secretion of sweat was almost equal on the two arms, and on two other occasions it was greater on the paralysed side. The surface temperature was always considerably greater on the paralysed arm, so that here the losses from evaporation, radiation, and conduction are all greater from the arm with the higher internal temperature; therefore in this case, too, the quantity of heat must have been considerably greater in the paralysed arm than in the other. In neither case was there any visible dilatation of the vessels on the paralysed side, nor have I ever observed, even when the difference of temperature has been considerable, any dilatation of the vessels on the paralysed side in hemiplegia. This is evidence against the view that the increased temperature on the paralysed side is due simply to there being more blood in the paralysed arm, and when we remember that damage to the central nervous system, whether experimental or clinical (it is the commonest cause of hyperpyrexia in man), frequently causes such a high temperature that we must suppose the production of heat to be increased, the observations I quote do lend strong support to the view that when in hemiplegia the temperature is raised in the paralysed axilla the increased quantity of heat in the paralysed arm is due to a greater production of heat in the paralysed than in the corresponding unparalysed limb. If this is so we have an instance in which a lesion that paralyses the motor function of muscle excites the thermogenetic, and thus we see that the functions are distinct.

TEMPERATURE RAISED IN FEVER.

In any fever the general conditions and temperature of different patients vary very much, and, therefore, single observations would be of little value. I have consequently made several observations on different patients in the following fevers, and if all the cases of any particular fever point in the same direction we shall probably be justified in drawing conclusions. I have always compared the observations made on a patient with a febrile temperature with those made on the same patient before he got up, but when his temperature was normal, or very nearly normal, unless his death prevented this.

1. *Typhoid Fever*.—In all cases the amount of sweat secreted during the fever was small, and less than when the temperature had fallen, therefore during the fever the loss of heat from evaporation was diminished. Also in every case except one the surface temperature rose less than the internal; in other words, the loss from radiation and conduction was not increased proportionately to the rise of internal temperature, and was therefore, relatively to the internal temperature, less during the fever. Thus, there is distinct evidence that in typhoid fever the loss of heat is relatively to the rise of temperature diminished, and this diminished loss no doubt helps to raise the internal temperature. Still, as the

surface temperature always rose considerably, the actual loss by radiation and conduction must have increased during the fever, and probably the decreased secretion of sweat was not enough to counteract this. Therefore, we may conclude that although certainly in typhoid fever the pyrexia is partly due to a lessened loss of heat, yet probably, too, the production is increased. Typhoid fever is remarkable in that, although it is a long-lasting disease with a high temperature, it has a low mortality, and these results suggest that one reason for this may be that the temperature is raised with the least expenditure of energy, for it is obviously more economical of energy to raise the temperature by diminishing the loss of heat than by increasing the production. Then, too, it is interesting to note that experience has shown that the best way to reduce pyrexia in typhoid fever is by sponging, that is, by increasing the loss of heat. It is well known that the internal temperature of typhoid fever is irregular as convalescence approaches, and the same is true of the surface temperature. In one case as the patient got worse the surface temperature approached the internal—that is to say, he was dissipating energy the faster.

2. *Pneumonia*.—In all cases except one the excessive sweating associated with the crisis lasted some days after it. This sweating must aid in keeping down the temperature after the fever. In every case during the fever the secretion of sweat is less than when the temperature has fallen, but as then the secretion is still rather more than in healthy individuals and in typhoid fever patients whose temperature has fallen, probably the sweat secreted during fever is not much less than before the fever began. Therefore it appears that the diminution of the loss of heat owing to a diminution in the amount of sweat secreted plays but a small part in raising the temperature. In only two cases did the surface temperature rise less than the internal, and in two it rose more (this explains the hot skin of pneumonia); therefore often in pneumonia the loss by radiation and conduction during the fever must be enormous. As, therefore, in pneumonia the loss of heat is so increased that the rise of temperature is for the most part due to an increased production of heat.

3. *Erysipelas*.—In both these cases observed the sweat was increased during the fever, and the surface temperature rose more than the internal temperature, so the dissipation of heat by evaporation, conduction, and radiation must have been very much increased. Consequently, the rise of temperature must have been entirely due to an increased production of heat, and not at all to a diminution of the loss.

4. *Suppuration*.—Here the considerable amount of sweat secreted after the abscess was opened and the temperature had fallen, appears to show that sweating helped to keep the temperature down. In the first case at least the quantity secreted during the fever was so small that the raised internal temperature may have been partly due to a diminished loss of heat from lessened evaporation. In both cases, too, there was a diminished loss from radiation and conduction, for the rise of surface temperature was less than that of the internal temperature; in fact, on one occasion the surface temperature was actually lower during the fever than when the internal temperature had fallen, and that although there was no excessive evaporation of sweat to bring it down. We may conclude that in these two cases of suppuration the raised internal temperature was, in part at least, due to a diminution in the loss of heat.

5. *Tubercle*.—Although the temperature fluctuates considerably tuberculous patients are always suffering from their disease, and therefore it is very difficult to obtain a normal standard. Consequently I do not think the figures teach much about the process of this fever. The amount of sweat secreted was large in the cases observed even when the temperature was low, so that no doubt the increased secretion of sweat is often the cause of the fall of temperature. It is interesting to notice that, on the whole, the cases of general tuberculosis secreted less sweat than those of phthisis. In both forms of disease the surface temperature is, as a rule, low; this may be due to the large amount of sweat. As the evaporation of this sweat must lead to a considerable loss of heat it is probable that in tubercle the production of heat is, as a rule, increased.

It appears from these observations that in some fevers the

pyrexia is, in part at least, due to a diminished loss of heat; in others this is of but slight importance, and then the production of heat is very great; and this result goes to prove the truth of the suggestion I made earlier—namely, that the processes which lead to a rise in temperature are different in different fevers. We have also incidentally learnt much about the surface temperature, and the secretion of sweat in the fevers we have considered.

Before leaving this subject I should like to draw your attention to some interesting figures obtained from a case of leucocythæmia. There was one white corpuscle to every ten red. The hæmoglobin was reduced to 17 per cent. of the normal, and the red corpuscles to 24 per cent. An observation gave the following result. The temperature of the mouth was 102.2°. The surface temperature at the umbilicus was 102.2°. The amount of sweat secreted in ten minutes was 1.2 mg. There are two points of interest here. First we have the high surface temperature with very little sweat; probably it would be impossible to have so high a surface temperature if there were much sweat, for the evaporation of the sweat would reduce it. Secondly, as the surface temperature was so high the loss by radiation and conduction must have been excessive, so the production must have been very high, and therefore even the small amount of hæmoglobin and the plasma carried between them enough oxygen for the metabolism necessary for this increased production. My best thanks are due to my house physicians, Dr. A. Salter and Mr. C. J. Harnett, for much help in these investigations. I have many observations on other fevers which I hope to publish subsequently.

Maragliano has published some observations which show that in some fevers the temperature depends very much upon the state of the cutaneous circulation—that is to say, upon the loss from radiation and conduction. He applied the plethysmograph to the arm of fever patients after experiments on healthy men, and, in the fevers upon which he experimented, he found that as the temperature rose the volume of the arm diminished, and as the volume of the arm is chiefly dependent upon the flow of blood through the vessels of the skin, it follows that this was diminished; in other words, less heat was lost by radiation and conduction when the temperature rose.

In one case, a man suffering from malaria, the contraction of the vessels preceded the rise of temperature; this is very strongly suggestive that it was the cause of it. In another case, a young man suffering from ague, the vessels began to dilate before the temperature began to fall—a fact which also strongly suggests that the fall of temperature was due to the increased vascular cutaneous dilatation.

I should like now to refer to some observations which Dr. Archibald Garrod and I made upon each other with the object of finding out how healthy individuals varied their loss of heat under different circumstances. In five pairs of observations, all made after dinner in the evening, the temperature of the mouth of both of us was almost the same, but four times out of the five my surface temperature was higher than his—on two occasions as much as 3.6° F. A possible explanation for the fact that on the other occasion my surface temperature was lower than his, is that we made the observations in his house, which is about 200 yards from mine, and on the evening on which this observation was made I strolled to his house after dinner while there was a cold north-east wind blowing. On four out of the five occasions I sweated more than he did. The whole amount I perspired during the five observations was 15.3 milligrammes, but his total amount was only 11.1, so that I perspired 38 per cent. more than he did. As, therefore, my surface temperature was higher, and the amount of sweat that I secreted was greater, I lost heat from radiation, conduction, and evaporation faster than he, and therefore my production per kilogramme was greater than his. We consequently learn that even healthy individuals vary much in their production and loss of heat.

It is well known that muscular contraction is accompanied by a rise of temperature; therefore, were it not that the loss of heat is increased during exercise the temperature would always rise. In fact, often the increased loss does not compensate for the increased production, and so the temperature does rise. Jürgensen found that the work involved in sawing wood for six hours raises the temperature of a healthy

man 1.2° C. above normal. Davy showed that walking for two or three hours raised the temperature of the urine 0.8° C. Clifford Allbutt found that Alpine climbing raised the temperature of the mouth half a degree, and Liebermeister and Hoffman found their axillary temperature raised by the exertion both of coming up and going down a mountain. Hobday, too, has taken the rectal temperature of omnibus horses, and he finds that it is raised about 2° C. by hard work. Dr. Garrod and I thought, therefore, it would be of interest to make a series of observations on the effect of exercise on each other. A bicycle ride of seventeen miles in a high, dry, cold wind reduced the internal temperature of Dr. Garrod by 1.0, and my own by 0.8 of a degree. I do not think this can have been due to the inhalation of cold air cooling the mouth, for we did not take the temperature of the mouth until we had been in the house for twenty minutes after the ride, and it must have been due to the rapid loss of heat caused by our going quickly through the high, dry, cold wind. With the exception of this bicycle ride exercise always raised Dr. Garrod's internal temperature, while mine was either unaltered or raised, but, on the whole, much less affected than his. There are other interesting points of contrast between us. Dr. Garrod's surface temperature was always considerably raised by exercise, while mine was usually slightly diminished. On the other hand, he sweated but little in comparison with me, for his average increased amount of sweat was 3.5 milligrammes, while mine was 7.05. The difference is really greater than this, for I took two bicycle rides in a cold dry wind to his one, and in a very dry wind the sweat evaporates fast, and consequently not much is secreted when the observation is taken. Apparently, therefore, we are different in our methods of accommodating the loss of heat to the increased production due to the muscular exercise; he chiefly increased his radiating and conducting power; I chiefly increased my loss by evaporation. We have, therefore, a demonstrated proof that different people accommodate differently. Further, when at rest my surface temperature is always higher than Dr. Garrod's, and I perspire more than he does; in other words, I lose heat faster, and, therefore, must produce more calories per kilogramme. As in fever, so here the data are not sufficient for us to say how many calories of heat the muscular exercises causes to be produced; we can only say that it must at least have been more than a certain number; for instance, my weight without my clothes is 61 kilogrammes; supposing the effect of exercise was to raise my temperature 0.5° C., and the loss by radiation, conduction, and evaporation were all increased, then I, as soon as when taking exercise I had raised my temperature 0.5° C., I must have produced at least more than an additional 30.5 kilogramme Centigrade calories, as my specific heat is nearly that of water.

REFERENCES.

¹ These experiments were related and discussed in the *Journal of Physiology*, Supplementary Number, 1892. ² Published in the *Journal of Physiology*, vols. xi and xii. ³ BRITISH MEDICAL JOURNAL, November 19th, 1894. ⁴ Das Verhalten der Blutgefässe im Fieber und bei Antipyrese, *Zeit. f. klin. Med.*, Bd. xiv., 1888.

REPORTS OF SOCIETIES.

OPHTHALMOLOGICAL SOCIETY OF THE UNITED KINGDOM.

E. NETILESHIP, F.R.C.S., President, in the Chair.

Thursday, June 10th, 1897.

FILARIA LOA.

DR. ARGYLL ROBERTSON read notes of the further history of a case of filaria loa previously reported to the Society. The chief point of interest arose from the circumstance that on returning to Old Calabar the patient was again troubled with the appearance of a worm in different parts of the body, and on one occasion a worm was removed by a missionary who possessed some medical skill from the abdominal wall in the left lumbar region. The parasite thus removed was sent to Dr. Robertson for examination. Unfortunately the preparation was imperfect, consisting merely of the uterine tubes and appendages and alimentary canal of a filaria. The em-

bryos in the uterus presented all the characters of the embryos found in the interior of a female filaria loa. It would thus appear that the filaria loa does not confine its visits to the neighbourhood of the eye or eyelids, but may appear under the skin of any part of the body, attracting attention only when it appears at a part where the skin is thin and sensitive, or when it presents itself under the delicate thin conjunctiva. The blood drawn at different periods of day and night, the urine, the fæces, the saliva and nasal mucus were all carefully examined by Dr. Robert Thin, the patient's medical attendant, for the presence of embryo filariæ, but the result was negative. The preparation of the parasite from the abdominal wall, and the preparation of a female filaria loa removed from the same patient's eyelid were exhibited microscopically.

SYMPATHETIC OPHTHALMIA.

MR. F. R. CROSS related some cases. Ten years ago he had reported to the Society a series of cases in which Mules's operation had been done instead of enucleation; in 2 of them mild sympathetic ophthalmia had occurred seventeen and twenty-one days respectively after the operation. He now reported a serious case of sympathetic ophthalmia occurring after evisceration of the sclerotic. A boy, aged 15, had been struck with a stick three weeks before he was first seen; there was a perforating wound of the left eye in the sclero-corneal region, with prolapse of the iris and traumatic cataract. Vision was barely p. l. The right eye was well V = $\frac{1}{2}$ and J. 1, pupil active, media and fundus n. The left eye was eviscerated on the day of admission, and a small glass vitreous was introduced. Considerable chemosis followed, and on the fourth day a few conjunctival punctures were made, after which recovery was uninterrupted. The patient was discharged from the hospital apparently quite well fourteen days after the operation; twenty-six days after, on August 18th, he returned with iridocyclitis in the right eye; there was ciliary injection, discoloured iris, small sluggish pupil, posterior synechia, hazy media T.n.; V = $\frac{1}{2}$. The eye had become irritable a day or two after leaving the hospital. The stump of the eviscerated eye was at once removed; atropine, and later scopolamine, were used, and a dark bandage was applied. The eye gradually improved, but the pupil did not dilate well till nearly two months later. On stopping the mydriatic the iritis recurred at once, and it was found necessary to continue the treatment and dark bandage for three months longer. The patient remained in the hospital seven months after removal of the stump. Recovery seemed to have been complete; the pupil acted well; the fundus and media were normal—V = $\frac{1}{2}$ and J. 1. The author was of opinion that enucleation in the first instance instead of evisceration would probably have prevented the sympathetic inflammation in the other eye. Destructive irido-cyclitis, however, did occur in the other eye some time after enucleation for injury, and the author quoted cases illustrating this.

MR. ADAMS FROST thought that Mr. Cross's case was to be considered as one of those in which sympathetic inflammation in the other eye had occurred after enucleation, and that it did not bear on the question of the safety of evisceration. In his experience Mules's operation was very satisfactory; he had had 1 case of nine years' duration, 1 of eight, and perhaps 12 of more than four years' duration, and in none of his cases, which had healed completely without leaving a fistulous opening, was there any after-trouble; the stump was not irritable as frequently as after excision.

MR. BRICKERTON, referring to Mr. Cross's cases, said that in all three of them the recovery was perfect; he himself had had one case of sympathetic inflammation after Mules's operation, in which also there was a complete recovery, so that the sympathetic inflammation in these cases did not appear to be very serious. What was wanted was a case of sympathetic ophthalmia after Mules's operation in an eye which could not of itself have started the inflammation, as for instance a glaucomatous eye, before it could be said that the inflammation was the direct result of the evisceration.

MR. CRICHTON and DR. ARGYLL ROBERTSON thought that the percentage of cases of sympathetic inflammation after excision was extremely low.

THE PRESIDENT agreed with the two last speakers, and thought it would be hazardous to act upon the supposition