Fig. 4.—Cast of a guinea-pig, with intestines exposed and partially distended with air. Note the absence of evidence of any compression.

or to patch any portions which exhibit a tendency to crack. In order not to have to wait long for this new coating to set, potassium sulphate, about a drachm to the pint, may be dissolved in the water used to make the cream. This greatly hastens the setting of the plaster. As soon as the plaster has set quite firmly the moulds may be carefully removed with any precautions that the exigencies of the case may suggest.

MAKING THE CAST.

After seeing that the surfaces of the mould are free from particles of plaster, etc., the different sections are fitted together, and the line of junction smeared with plaster cream. No preparation whatever of the surface of the mould, such as oiling, etc., is necessary. Thus no detail is blurred or rendered indistinct. Into the hollow mould thus formed the plaster cream for the cast is poured in the ordinary way.

REMOVING THE CAST FROM THE MOULD.

Fig. 5.—Cast of a hand.

Of all the advantages which this process may be considered to present over the older methods, it is my belief that the ease with which the mould can be separated from the cast is one of the most important. The method is as follows: As soon as the plaster in the cast has well set, the whole mass is put into a bath of hot water, say at 110° F. In the course of half an hour or an hour, the heat has penetrated through the

outer plaster casing, and warmed the paraffin, so that it becomes soft and plastic. The mass is then removed from the water, and the outer plaster mould broken away in large masses by means of a chisel. Then it will be found that the whole of the paraffin mould can be removed with the greatest ease, simply by seizing it with the fingers and lifting it away from the surface, as one would lift a piece of wet gauze. Thus there is, with ordinary care, no danger whatever of injuring the surface of the cast, and the whole process of separating the mould from the cast, which by the old plaster method of casting is so laborious and fraught with so much danger to the cast, can be done with absolute safety, with perfect ease, and in a very few moments. Moreover, "undercutting" is of no importance whatever, because of the plasticity of the mould, and the possibility of melting it out if necessary without any injury to the cast.

It may be objected that the apparatus is cumbersome and costly, and that the process involves some practice and a good deal of trouble. In reply I would merely say—look at the results. Has it ever before been possible to cast a hand showing such minuteness of detail as depicted in Fig. 5? Has it hitherto been possible to cast such a subject as Fig. 4 at all? In these two subjects are seen the delicacy of the molten paraffin in adapting itself to minute variations of surface (Fig. 5), and (Fig. 4), the absolute absence of the compressing effects which are so noticeable when plaster or other heavy materials are used.

CASES OF DIPHTHERIA TREATED BY INJECTION OF ANTITOXIC SERUM IN UNIVERSITY COLLEGE HOSPITAL DURING 1896 AND 1897.

BY

SIDNEY MARTIN, AND
M.D., F.R.S., Assistant Physician to the Hospital.

G. BERTRAM HUNT,
M.D., Resident Medical Officer.

IN the BRITISH MEDICAL JOURNAL of 1896, vol. i, p. 197, is a report by Drs. Martin and Smith of the cases of diphtheria treated in University College Hospital during the year 1895, the first antitoxic year, and a comparison of the results thus obtained with those of the four previous years in which antitoxin was not used. Although the belief in the antitoxic treatment is more widely and firmly held now than at the time of the last report, it seems desirable to continue these comparisons so as to include the last two years; not merely to add another proof of the efficacy of the treatment by antitoxin, but also to illustrate some still unsettled points connected with the proper dose and the best methods of using the serum.

During the years 1896 and 1897 there were 185 cases admitted into the diphtheria wards, but of these cases 7 are not available; 5 (which recovered) because no antitoxin was given as the cases were not considered to be true diphtheria; and 2 (1 fatal) because the antitoxin was only given by the mouth. There are thus left 178 cases treated by the subcutaneous injection of antitoxin to be compared with the cases treated without it. The severity of the cases admitted into University College Hospital, as we believe, in the other general hospitals, is on the whole rather above the average, since owing to the limited number of beds all cases of diphtheria seen as out-patients cannot be admitted, and preference is given to severe cases, especially those with

laryngeal obstruction. Therefore these cases treated by antitoxic serum can only be fairly compared with the cases in the same hospital during previous years.

BACTERIOLOGICAL DIAGNOSIS.

Out of 178 cases the diphtheria bacillus was found in pure growth or mixed with cocci in 149; in 18 cases the bacillus was not found on examination, and in 11 cases there is no record of the cultivation. A few of the cases in which the bacillus was not found were undoubtedly diphtheria, in fact in two of them paralysis developed; others, though presenting the clinical picture of the disease, failed to give the bacillus in repeated examination and possibly were not diphtheritic. Still, for the purpose of comparison with preceding years, they must be included, since such cases were of course not excluded formerly.

It has been suggested that the lessened mortality during the past few years has been due not so much to the serum treatment as to the introduction of the bacteriological diagnosis of diphtheria into practical work, so that mild cases which formerly would not have been diagnosed as diphtheria are now admitted as such chiefly on the bacteriological diagnosis, that their inclusion in the total number of cases would account for the lowered mortality. This factor can have had no effect in the reduction of our mortality, for almost all the cases were admitted into the hospital on clinical grounds alone, the bacteriological examination being made afterwards.

TREATMENT EMPLOYED.

In all these cases a subcutaneous injection of antitoxin was given within twelve or twenty-four hours after admission, the dose varying somewhat from time to time. In 1896 the dose was between 2,000 and 20,000 normal units, the average being about 7,200; one-third of the cases receiving 2,000 to 4,000 normal units, and two-thirds receiving 5,000 to 20,000. In 1897 the number of units given was between 4,000 and 12,000, the average being 7,000 to 8,000; one-seventh of the cases receiving 4,000, and six-sevenths between 6,000 and 12,000 normal units. In the beginning of 1896 twenty cases were injected with the serum of the Institute of Preventive Medicine, and two with Behring's serum; since then in every case we have used the serum kindly supplied by Dr. Sims Woodhead from the laboratories of the Royal Colleges of Physicians and Surgeons, usually of a strength of 4,000 normal units in 5 c.cm., and have found this concentrated serum very convenient. Local treatment was applied to the throat in almost all the cases. In the majority swabs of chloride of zinc (gr. x to 3j) were employed; in others sprays of perchloride of mercury or douches of chlorine water were used. All cases had steam sprays of bicarbonate of soda at two-hourly intervals. When we consider the very frequent association of pus and septic micro-organisms with the diphtheria bacillus in the throat, it is clear that we must not rely on antitoxin alone, but must also use local antiseptics in order to obtain the best results. In a few of the cases in which these were not used, there were glandular abscesses in the neck, which were not seen when antiseptics were again employed. Nasal discharge was treated by douches of boracic lotion or chlorine water. For laryngeal obstruction tracheotomy was performed when necessary; in no case was intubation resorted to.

EFFECT ON MORTALITY.

This is shown in Table I, in which the mortality of the three antitoxin years is compared with that of the four preceding ones. It will be seen that the death-rate both of the laryngeal and other cases has been greatly reduced, and that the total mortality of all cases has fallen from between 33 and 43 per cent. to 28, 17.7, or 17 per cent. for the three antitoxin years.

The three chief points to be discussed in considering the effects of the antitoxic treatment on the mortality are (1) the average age of the patients, the disease being notoriously more fatal the younger the patient; (2) the proportion of laryngeal cases; and (3) the day of disease on which the patients were admitted, for the earlier the disease is treated the lower the mortality. A great preponderance of older patients, a low proportion of laryngeal cases, or a series of cases admitted early in the disease might be factors tending to reduce the mortality for any given year, especially when dealing with a small number of cases. On looking at Table I, the ages of the

patients and the number of laryngeal cases being given for this purpose, it will be seen that the first two of these factors do not come into operation. In the first place, the proportion

TABLE I.—Mortality of All Cases.

Year.	Age.	Laryngeal Cases.			Pharyngeal, Nasal, etc.			Total.		
		No. of Cases.	Deaths.	Percentage Mortality.	No. of Cases.	Deaths.	Percentage Mortality.	No. of Cases.	Deaths.	Percentage Mortality.
1891	5 years and under	25	18		14	4				
	Over 5 years	5	3	70.0	18	2	18.7	62	27	43.5
1892	5 years and under	21	14		12	1				
	Over 5 years	4	1	60.0	23	4	14.3	60	20	33.3
1893	5 years and under	32	24		26	6				
	Over 5 years	9	8	78.0	38	1	10.9	105	39	37.0
1894	5 years and under	26	3		22	5				
	Over 5 years	8	3	47.0	4	4	30.0	64	25	39.0
1895 (1st antitoxin year.)	5 years and under	23	7		25	10				
	Over 5 years	7	3	33.3	20	1	24.4	75	21	28.0
1896	5 years and under	13	7		30	6				
	Over 5 years	4	2	40.0	38	1	10.0	90	16	17.7
1897	5 years and under	24	6		29	7				
	Over 5 years	10	2	23.5	25	—	13.0	88	15	17.0

of patients under 5 years to the total number of cases is in the pre-antitoxin years 63, 53, 56, and 53 per cent. respectively, while in the three antitoxin years it is 64, 65, and 63 per cent.; so that as far as the age of the patient is concerned, since there is a greater number of patients under 5 in the antitoxin years, we should expect a greater mortality in the antitoxin cases. The percentage of laryngeal cases to the total number of patients from 1891 to 1894 was 48, 41, 39, and 52 per cent., while in 1895 to 1897, the antitoxin years, it was 40, 24, and 38 per cent. In only one year—1896—was the proportion of laryngeal cases much below the average. The effect on the mortality of this smaller proportion of laryngeal cases in the antitoxin years would be about equalised by the larger proportion of cases under 5 years. In Table III it will be shown that there is no great difference in the date of commencement of treatment in the two series. In the pre-antitoxin cases 35 per cent. were admitted after the fourth day of the disease; while in the antitoxin series 31.6 per cent. were admitted after the fourth day; but this slightly earlier time of admission is quite insufficient to account for the great fall in the death-rate of the last three years. The examination of the two series of cases, therefore, reveals no factor capable of lessening the proportion of deaths except the effect of treatment; and as the only change in the treatment has been the use of antitoxic serum, the lessened mortality may fairly be ascribed to it.

TRACHEOTOMY CASES.

Besides considering the effect of antitoxin on the general mortality, it is worth while looking at its results in the tracheotomy cases alone, for the reason that the severity of these cases varies less from year to year. A case of diphtheria

TABLE II.—Mortality of Tracheotomy and Intubation Cases.

Year.	Tracheotomy.		Intubation.		Total.		
	Cases.	Died.	Cases.	Died.	Cases.	Died.	Mortality %
1890	41	30	1	1	42	31	74
1891	17	15	17	9	34	24	70
1892	12	6	8	3	20	9	45
1893	39	30	—	—	39	30	77
1894	34	17	—	—	34	17	50
1895	25	5	—	—	25	5	20
1896	18	7	—	—	18	7	39
1897	29	7	—	—	29	7	24

requiring tracheotomy must always be a serious one; so that a marked lowering of the death-rate of these cases is more certainly than in the pharyngeal cases due to the result of treatment, and not to any variation in the intensity of disease for that year.

With the tracheotomy cases for the years 1890, 1891, and 1892, we must include the intubation cases, for in those years many cases in which the trachea would now be opened were intubated. In Table II it will be seen that the mortality of these cases has been reduced from 65.5 per cent. for the four pre-antitoxin years to 26.4 per cent. for the three antitoxin years; that is, the mortality has been reduced by 40 per cent. A noticeable and important point about these cases is that the operation was performed in almost every case either on admission or within twelve hours; and that no case which on admission was free from laryngeal symptoms developed them in hospital. This illustrates the power of the antitoxin to prevent spread of the membrane. The greater mortality of the tracheotomy cases both in 1896 and 1897 than in 1895, although the total mortality for the same years was much lower, is probably due to the large number of fatal cases which were admitted late in the disease during these two years; of the 14 fatal tracheotomies during 1896 and 1897 no fewer than 8 were admitted after the fourth day, and in 7 cases the membrane had extended into the bronchial tubes before treatment was commenced.

CAUSE OF DEATH.

It is necessary to state the causes of death, and especially to examine closely for any indication of spread of membrane in fatal cases after the antitoxin injection has been given. Of the 31 fatal results during the last two years, 11 died between the third and eleventh days, with suppression of urine usually accompanied by severe vomiting, the typical toxic anuria of diphtheria. In all these cases the membrane had disappeared before death. It was at first asserted that the antitoxic treatment increased the mortality from this complication, yet larger experience has shown, as in the cases collected by the Committee of the Clinical Society,¹ that the serum treatment has neither increased nor diminished the number of deaths from this complication; but as the deaths from other causes have been much reduced the deaths from toxic anuria probably bear a greater proportion to the total number of fatal cases, and so have come into greater prominence.

Eight cases, all after tracheotomy, died with membrane in the bronchial tubes, usually associated with broncho-pneumonia. In 2 cases death occurred on the first day, in 5 on the second day, and in 1 on the third day after the injection of serum. Two of the patients were moribund on admission, and in none did tracheotomy give complete relief, which, taken with the fact that they all died within three days tends to show that the membrane was present in the tubes before the commencement of treatment, though in these cases it is impossible to say whether or not it afterwards spread. Five patients died of pneumonia without affection of the tubes, 1 moribund on admission within eight hours, 1 on the second day, 1 on the fifteenth, 1 on the thirty-fifth, and 1 on the forty-fourth. In the last 3 cases the children were free from membrane, and the pneumonia was probably an accidental complication. Two patients died from cardiac failure, 1 within eight hours of admission and 1 on the fourth day. Two patients died of gastro-enteritis, 1 on the third and 1 on the fifth day; and 1 died of pneumothorax following tracheotomy on the first day.

Nearly all the fatal cases, except those dying within two days of the commencement of treatment, were free from membrane at the time of death; and in none was there any evidence that the fatal result was due to extension of the membrane after the injection.

MORTALITY ACCORDING TO DAY OF DISEASE IN WHICH TREATMENT WAS COMMENCED.

On dividing the cases into two classes, as in Table III, one containing those admitted from the first to the fourth day of the disease, and the other those admitted after the fourth day, we find that the antitoxic treatment has reduced the mortality of the first class from 37 to 14 per cent.; while in those admitted after the fourth day the reduction was much slighter, from 43 to 35 per cent.; in fact, in the first antitoxin

year, there was no reduction in this class, and it was only when larger doses of serum were used that any effect was produced in the mortality of cases admitted after the fourth day.

TABLE III.—*Mortality According to Day of Disease.*

	Admitted from First to Fourth Day of Disease.			Admitted after Fourth Day of Disease.		
	Cases.	Deaths.	Percentage Mortality.	Cases.	Deaths.	Percentage Mortality.
1891	39	19	48.7	21	8	38.1
1892	31	11	35.5	24	9	37.5
1893	50	14	28.0	34	17	50.0
1894	36	14	38.8	23	10	43.0
1895	46	8	17.3	28	13	46.4
1896	66	9	13.6	24	7	29.0
1897	60	7	11.7	28	8	28.6
			37			43
			14			35

This important fact that a marked lowering of the death-rate is only seen in cases treated during the first four days is one brought out by most collections of cases, and is well seen in the reports of the Metropolitan Asylums Board for 1895 and 1896. It has a distinct practical bearing, as it not only explains the importance of giving the antitoxin as soon as the case is seen, but it also indicates the necessity of giving a full dose at once, and not administering the serum in divided doses on successive days; for in this case some of the later doses in patients on whom the treatment is begun on the third and fourth days are not so effective in counteracting the disease as shown by the mortality table.

DOSAGE OF ANTITOXIN.

In Table I, besides the general reduction of mortality in all the antitoxin years, it is shown that the total mortality of 1895, the first antitoxin year, was 28 per cent., which was much greater than that of 1896 and 1897, the second and third antitoxin years, in which it was 17.7 and 17 per cent. respectively. During 1895 the antitoxin used was obtained from various sources, its strength in normal units was not known, but it was certainly very weak, so weak that an insufficient dose (often under 1,000 normal units) was generally given. The only change in the treatment during the next two years was the administration of a more concentrated serum of known unit strength; and of this a much larger dose, averaging 7,200 units in 1896 and 7,800 units in 1897, was employed. In the later year, six-sevenths of the cases received over 6,000 normal units, and it is now the practice of the hospital to give each patient on admission a dose of not less than 6,000 normal units. One of the most important and instructive results shewn by our cases is this great saving of life which has been effected by the increase of the dose of serum. (See also Table III.)

EFFECT OF ANTITOXIN ON TEMPERATURE.

The symptom on which the effects of antitoxin can be best demonstrated by means of figures is the temperature, and Table IV shows that in the cases which recover the temperature falls sooner in those treated by antitoxin.

TABLE IV.—*Effect of Antitoxin on Temperature.*

	1892-93-94— Pre-antitoxin Years.		1896-97— Antitoxin Years.	
	Cases.	Percentage.	Cases.	Percentage.
Temperature Becoming Normal.				
Before fourth day	68	48	129	83
Between fifth and eighth days ...	40	28	20	12
After eighth day	33	23	8	5

The table shows that, while only 48 per cent. of the pre-antitoxin cases attained a normal temperature before the fourth day, in as many as 83 per cent. of the antitoxin cases did the fever cease during that time, and that therefore there were a very much larger proportion of cases, in the pre-antitoxic series in which the cessation of fever was deferred

till after the fifth and eighth days. From this table all cases have, of course, been excluded which ran an afebrile course throughout, and also those dying during the treatment. The figures, therefore, rather understate the result, since in the antitoxic series the mortality is much less, and therefore the number of recoveries of severe cases with more prolonged temperature is greater.

EFFECT OF ANTITOXIN ON OTHER SYMPTOMS.

Paralysis was observed only in 23 of the 178 cases during the time they were in the hospital; but it occurred after leaving the hospital in several children who attended later as out-patients, and probably in others who were not seen again; so that it is impossible to say in what proportion of cases it was produced; but it is probable that in these cases, as in other recorded series, the frequency of paralysis has not been reduced by the antitoxin treatment.

As to the effect on the duration of membrane in the throat, albuminuria and other symptoms, it is impossible to speak definitely, as no exact particulars as to these facts can be obtained for the pre-antitoxin years, and so no comparisons can be instituted; but it is a matter of ordinary clinical observation that the duration of visible membrane is much shorter in the antitoxin cases; and also that it is extremely rare to observe any spread of the membrane either locally in the throat or to previously unaffected parts such as the nose and larynx, after twenty-four hours from the time of injecting the serum.

BAD EFFECTS OF ANTITOXIN.

In 49 of the 178 cases rashes were observed. The rash was urticarial in 19 cases, erythematous or scarlatiniform in 16, papular or morbilliform in 14. In 8 cases it appeared between the second and fifth day, in 25 between the sixth and tenth, and in 16 between the eleventh and fifteenth. In most cases the rash caused no disturbance or uneasiness except some itching; but in 13 cases there was a rise of temperature, and in four cases more or less severe joint pains with some synovial effusion, affecting only the larger joints, especially the hip, knee, and shoulder. In 3 of these cases with joint pains the temperature reached 104°. In 1 case an abscess formed at the seat of injection. This must have been due to some fault in the technique of the injection and not to the serum; as the serum then in use was at once examined bacteriologically with a negative result both at the laboratories of the Royal Colleges of Physicians and Surgeons and by us. No other bad symptom was seen, which could be ascribed to the antitoxin.

CONCLUSIONS.

1. The great reduction seen both in our total and tracheotomy mortality can be only due to the treatment with antitoxic serum, for the examination of the cases reveals no other factor capable of producing this result.

2. The full benefit of antitoxin was not obtained until fairly large doses (on an average between 7,000 and 8,000 normal units) were employed; the mortality then fell from 28 per cent. in the first antitoxin year to 17.7 and 17 per cent. in the next two years. Each case of diphtheria should have an initial dose of not less than 6,000 normal units.

3. Since antitoxin has but little effect on the mortality when given after the fourth day, it should be administered as soon as the patient is seen without waiting for the bacteriological diagnosis, and the full dose should be given immediately in one injection, not in divided doses on successive days.

4. Not only does antitoxin reduce the mortality, but it also distinctly lessens the severity of the disease in cases which recover. This is best shown by the effect on the duration of the fever, which in our series lasted only four days in a very much larger proportion of antitoxin than of pre-antitoxin cases.

5. The effect of antitoxin is chiefly seen in its specific action of the diphtheritic membrane, as shown by the cessation of its spread and more rapid clearance, and also by the greatly-lessened mortality of such cases in which the chief danger is from extension of membrane, the mortality of tracheotomy cases especially being reduced from 65.5 to 26.4 per cent., a greater reduction than seen in the other cases.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, 1898, vol. i, p. 1455.

FURTHER NOTE ON BILE AS AN ANTIDOTE TO VENOMS AND DISEASE-TOXINS.

By T. R. FRASER, M.D., F.R.S., F.R.C.P.E.,
Professor of Materia Medica and Clinical Medicine, University of Edinburgh.

I HAVE already shown that the bile of several animals possesses antidotal properties against serpents' venom and against the toxins of such diseases as diphtheria and tetanus, and that the bile of venomous, or more correctly of nocuous serpents is specially powerful as an antidote against the venom of serpents.¹ The experiments have been extended, with the result that further proof of these propositions has been obtained.

Taking, in the first place, the dry Indian cobra venom, whose minimum lethal dose for rabbits is slightly less than 0.0024 g. per kilo., in a series of *in vitro* experiments a definition has been obtained of the smallest quantity of the dry bile of several serpents, nocuous and innocuous, and of a number of higher animals, to render non-lethal 0.0025 g. per kilo. of this venom, when solutions of each were mixed together and left in contact for ten minutes. The quantity of each bile required per kilo. of animal was approximately:

Crotalus bile...	0.0005 gramme	Indian python bile	0.005 gramme
African cobra bile...	0.001 "	Grass snake bile	0.0065 "
Pseudechis bile	0.002 "	Rabbits' bile	0.0075 "
Puff adder bile	0.003 "	Human bile...	0.015 "
Daboia bile	0.003 "	Ox bile	0.02 "
Indian cobra bile	0.004 "	Pig's bile	0.02 "

Corresponding results have been obtained with the venom of several other species of serpents.

The most important results are that the bile of nocuous or venomous serpents is the most powerful antidote to venom, and is closely followed in efficiency by the bile of innocuous serpents, while the bile of animals having no venom-producing glands—as man and the ox, pig, and rabbit—while definitely antidotal, is less so than the bile of innocuous serpents, and much less so than the bile of nocuous or venomous serpents. It is remarkable to find that the bile of one species of venomous serpent may actually be a more powerful antidote against the venom of another species than is the bile produced by this species, and that there is no direct correspondence between the toxic activity of the venom produced by a serpent and the antidotal power of the bile of that serpent.

Extending these experiments to the toxins of disease, it was found, in the case of a specimen of diphtheria toxin whose minimum lethal dose for rabbits was 0.05 c.cm. per kilo., that a dose of 0.1 c.cm. per kilo. could be rendered non-lethal by:

Daboia bile	0.025 g. per kilo.	Ox bile	0.1 g. per kilo.
Rabbits' bile	0.02 "	Human bile...	0.15 "

When tetanus toxin, tested, however, in guinea-pigs, was similarly examined, in a dose of it exceeding the minimum lethal considerably more than in the experiments with serpents' bile and with diphtheria toxin, the results were that the smallest quantity of bile able to prevent death from occurring was with:

Daboia bile	0.005 g. per kilo.	Human bile...	0.15 g. per kilo.
Rabbits' bile	0.005 "		

The biles tested against toxins have been comparatively few in number, because the deterioration which rapidly occurs in the strength of toxins prevents a large number of consecutive experiments from being made with the same specimen. With the exception of human bile, all the biles were taken from the gall bladder. The bile, therefore, of the venomous serpents examined has also more antidotal power against the toxins of disease than the bile of the majority of non-venomous animals. It is noteworthy that among the non-venomous animals, the rabbit should produce a bile definitely superior to the others in antidotal quality against not only toxins but also venoms.

As disease toxins and venoms are excreted from the blood into the intestinal canal, the antidotal bile which they must there be brought into contact with, constitutes a potent means of defence of the body by rendering the toxins and venoms inert before they can be absorbed from the intestines into the blood. The therapeutic importance of remedies that are able to increase the hepatic secretion is thus indicated. Although