wrung out of the spirit corrosive fairly dry. Hartmann's corrosive wood-wool wadding forms the outer dressing. It is kept in well-fitting metal boxes. For the purpose of keeping dressings fixed, I employ fairly broad bands of adhesive strapping and elastic webbing bandages outside of the ordinary bandage, which must be firmly applied to keep the deep surfaces of the wound together and to limit exudation.

[The explanatory diagrams were made by Dr. W. G. Richardson.]

NOTES.

¹ There is now a good prospect of a new hospital. ² Since I learnt from Mr. Arbuthnot Lane of London some years ago the value of torsion, I have never used a ligature except in special localities like neck or axilla, or for some particular purpose. ³ Without any explanation it would appear as if I had been unfortunate in my operations for uterine fibroids. The reverse is the case. I have operated upon 66 cases by abdominal hysterectomy or myomectomy, and the only death I have had, with the exception of the two recorded was a single one ten years ago, in which the stump was treated extraperitoneally.

THE TREATMENT OF TYPHOID FEVER.*

By SIDNEY PHILLIPS, M.D. LOND., F.R.C.P.,

Senior Physician to the London Fever Hospital; Physician and Lecturer on Medicine to St. Mary's Hospital.

TYPHOID fever being an artificially produced disease, tends to vary with the conditions associated with its origin, and though such variations are slight individually and gradual in incidence, in their sum they suffice in time to produce a considerable variation in type from the original disease. Certainly there is considerable difference in the symptoms described fifty or even twenty-five years ago and those occurring to-day; the difference is marked in the lessened severity of the abdominal symptoms; the tongue is now often moist throughout the disease, instead of dry and baked; tympanites and diarrhoea are much less pronounced; probably also hæmorrhage and perforation are less common; tremors and dilatation of pupils are now uncommon; and instead of noisy, active delirium, the mind is often clear throughout even fatal cases. The typhoid state, with the patient sunk deep in bed, unable to move himself, and lying unconscious or semiconscious for days, is now quite exceptional. Nor is this a comparison merely between a grave and a slight case, for the mortality is not lessened. Nor is it the result of treatment, though this may produce some modifications. The difference is one of type, and is probably accounted for by the fact noted first by Vaillard¹ and since by myself,² Goodall,³ Moore, and others—that of late it is not very infrequent to find at *post-mortem* examination of typhoid cases that there is little or no ulceration of intestines. Such cases escape some of the dangers of enteric fever, and those therefore met with on the *post-mortem* table must represent many others with little or no ulceration of intestines that recover. The diminution in severity of abdominal symptoms is not therefore inexplicable, but corresponds with a tendency to lessened ulceration of intestines. Relapses, too, appear much more frequent of late years; in 1862 Murchison computed they occurred in 3 per cent. of cases; in 1897 they occurred in 13 per cent. of Metropolitan Asylums Board cases.

If typhoid of twenty-five years ago so differed from typhoid to-day, it probably differed still more fifty years ago; and when conditions existed such as made typhus rife, the distinctive characters of typhoid may well have been affected by such influences in a manner to account for typhus and typhoid fever being regarded at the time as one and the same disease.

In the treatment of the disease Jenner in 1879 insisted it was necessary to be acquainted with the "epidemic constitution of the period," and it is not less necessary now to be acquainted with the prevailing type of the disease.

PREVENTIVE TREATMENT.

The proposals which have been made to produce immunity by inoculations of bacillus cultures have not yet, at any rate, come into general use, and at present the prevention of typhoid fever depends upon those sanitary measures by which the proportion of persons dying annually in London from typhoid has fallen from 377 per million persons living in 1869 to 129 per million in 1897.

* Read before the Harveian Society of London, November 3rd, 1898.

But the fatality of the disease once contracted has little altered. Murchison found that the mortality of patients in the London Fever Hospital during the ten years ending 1862 was 18.5 per cent.; the same figure was shown in the cases occurring in various large hospitals in Great Britain and abroad. All figures available at the present day show about the same mortality. Thus, the Registrar-General's annual returns since the Infectious Diseases Notification Act in 1889 show that from 1890 to 1897 the mortality still remained over 18 per cent. The chances of recovery, therefore, of a person contracting typhoid to-day are little or no better than they were fifty years ago, notwithstanding the advances in medical knowledge and improved nursing and hygienic conditions.

In the treatment of a disease it is important to know the relative frequency of the various dangers to life: *post-mortem* records and returns of complications during life do not make these known, but an approximate estimate of the frequency of deaths from causes in each of the following groups can be made.

i. Death from general causes: toxæmia, hyperpyrexia, pyrexia, heart failure, "asthenia."

ii. Death from local lesions special to typhoid fever: perforation, hæmorrhage, etc.

iii. Death from intercurrent affections, pneumonia, etc., due to the streptococcus and pneumococcus mainly.

Of the causes in Group ii, perforation is most common; it has been variously estimated as causing 5.7 (Hoelscher⁴) to 20 per cent. of the deaths (Murchison, Hare⁵). (Other writers' estimates were given in the paper.) Certainly perforation will be over rather than under-estimated if credited with 20 of each 100 deaths.

Hæmorrhage may be taken as causing 10 of 100 deaths as an outside estimate. (The figures were given on which this estimate was based.)

Peritonitis, without perforation, accounts for 2.2 per cent. of deaths (Hoelscher). Other rare accidents such as rupture of mesenteric glands, of spleen, or gall bladder, can not together account for more than 2 per cent. of the deaths, and a liberal estimate cannot credit the causes in Group ii with more than 35 of each 100 deaths.

The intercurrent affections in Group iii are mainly pulmonary: pneumonia, bronchitis, etc. Nephritis, pericarditis, meningitis, are very rare causes, and this group is shown by statistics (which were given) not to account for more than 10 of each 100 deaths.

The causes grouped as ii and iii, therefore, do not together produce more than 45 of each 100 deaths, and if we allow for them, say, 50 per cent., there is ample margin for error. The remaining 50 of each 100 deaths must be due to causes falling in the first group, and these may be separately discussed here, though often associated in the disease itself.

First as to *toxæmia*. It is probably the more common cause of death, and the antiseptic treatment is based on the endeavour to lessen its dangers by destroying the toxins before absorption, and limiting the action of the micro-organisms in the alimentary canal.

The list of substances advocated for the antiseptic treatment is innumerable; for many the claim has been also made that they cut short the disease, but an "abortive" treatment of typhoid is not yet known. Many antiseptic substances have been experimentally proved to destroy the bacilli outside the body, for example, a 1 in 5,000 solution of corrosive sublimate, and the principle of the antiseptic treatment is no doubt correct. Jenner used to give charcoal long before the term "antiseptic treatment" was formulated, but it is bulky to take. Bichloride of mercury has been extensively used since it was advocated by Sir W. Broadbent⁶ in 1894, and it appears to me to be the most useful antiseptic, often lessening the temperature and the toxic symptoms; it rarely causes diarrhoea, but is probably better abstained from if there be hæmorrhage.

It is of interest that Murchison condemned the use of mercury in typhoid as "always useless and injurious." This difference in experience is another evidence of the altered type of typhoid fever.

Salol is another antiseptic substance which is useful. I find it a useful practice to give a daily enema of disinfecting solution, especially when there is constipation.

Hyperpyrexia is very rare in typhoid fever. Of 200 cases

secutive cases at St. Mary's Hospital it occurred only twice; the same percentage was found by Bryant in 608 cases collected from *Guy's Hospital Reports*; Ord,⁸ Osler, Fagge, and the statistics of the Metropolitan Asylums Board all show the rarity of its occurrence. But rare as it is in typhoid fever, it is infinitely rarer as a complication in itself; it is nearly always due to some local affection. In one of my cases pneumonia, in two perforation and peritonitis, in one suppurative meningitis, and in one rupture of the gall bladder were found *post mortem*. It should, then, never be assumed that hyperpyrexia is a complication in itself without a thorough search for some local affection to account for it. This may be of import in treatment, as the cold bath, which is the sole efficient means of treating hyperpyrexia, might do harm if it were due to perforation or peritonitis; in such cases treatment must be directed to the local condition and the temperature meanwhile lowered by sponging, quinine, etc.

Pyrexia is not now regarded as one of the chief dangers of enteric fever, and too much importance is not to be attached to the mere temperature. A pyrexia of 103° for two or three weeks in a previously healthy young adult lying in bed and taking food well will not produce serious consequences. Even high temperatures are more often an indication of some intercurrent affection than a danger in themselves, and severe cases may not have a high temperature. Dreschfeld⁹ and others record cases even fatal with no rise of temperature. The pyrexia usual in typhoid is for the most part effectively treated by tepid or cold sponging and quinine. When, however, the temperature is unusually high or unduly protracted, or with very irregular extensive rises and falls, or presents no remissions, other means must be adopted for its control, and of these there is none so effective as the cold bath. The claim made for it, that it reduces the mortality from 17 or 18 to 8 or 9 per cent. has not been disproved. It is recommended that whenever the temperature reaches 102.2° the cold bath shall be employed; practically, therefore, all patients have to be so treated, which means that the 82 who would recover even without the bath and the 9 who will die with or without the bath, that is, 91 of every 100 patients are subjected to repeated bathings unnecessarily, and that the 9 who die, have these repeated bathings added to their sufferings. Its infrequent use, therefore, in hospital, and still more in private, practice is due not so much to a doubt of its virtue as to its impracticability, and it is not likely to become of general utility unless some means can be found of selecting the cases which really require it. It is said that to obtain the advantages of the treatment it must be begun too early in the disease to predict if the case will prove a serious one. So far as hæmorrhage and perforation are concerned this is no doubt true, but these are accidents which are not rendered less frequent by the cold bath, and *quoad* the treatment, may be ignored.

The only causes of death which are reduced by the cold-bath treatment are those in Group 1, and it must be admitted that it is not always possible to tell early in the disease whether such dangers threaten, but if the temperature is not very high, is accompanied by remissions, is not very irregular in its rises and falls, if the pulse is not very rapid or feeble, and if the patient does not seem greatly affected by toxæmia, I think he might be spared the cold bath for a time at any rate. For it is not certain that the advantages of the bath are lost if deferred after the temperature touches 102.2°, Hare having obtained equally good results without following this rule.

Possibly, if a selection were thus made, the statistics of the cold-bath treatment would appear less favourable than when all the slight cases are submitted to the treatment, but I believe it would be more generally used, and more lives would be saved, than by adhering to a criterion of danger based on the discarded theory that pyrexia is the one danger of fevers.

Probably the temperature of the bath need not be so low as sometimes recommended; thus Hare's baths were usually at 75° or 80°, owing to that being the natural temperature of the water in Brisbane, and he obtained as good results as others have done, though it should be said in severe pyrexial cases he cooled the bath down by ice. In Dr. Barrs's tank treatment¹⁰ the temperature of the water is only 90 or 95°; this method, he finds, gives excellent results, but the special arrangements it requires are not to be obtained in a private dwelling. In patients not treated by baths it is generally agreed sponging

with tepid or cold water, affusions, and wet packs are better than the use of drugs. The best of the drugs is quinine, but it is slow in acting; acetanilide, antipyrin, and phenacetin are depressants, and act on the red corpuscles; this is denied in the case of antifebrin.¹¹ I never use these drugs in typhoid, but I have seen them do much harm in cases of fever at all approaching a malignant type.

Cardiac Failure.—Changes in the myocardium resulting from pyrexia and from toxins are recognised causes of cardiac failure in typhoid; they are not, however, of great frequency in the disease. Hoelscher only found advanced parenchymatous change in 19 per cent. in 2,000 necropsies of typhoid cases; it is indicated during life by the feebleness of cardiac impulse and sounds, and of the pulse; death may occur suddenly or gradually.

Sudden death sometimes occurs in typhoid in which no cardiac changes are found in the heart *post mortem*. Many explanations of these cases have been offered. Dewèvre,¹² though taking for granted such cases do result from cardiac failure, shows that none of the suggested explanations are satisfactory. These deaths appear of the same nature as deaths from diphtheritic paralysis, probably from peripheral nerve changes which are known to occur in typhoid.

Blood Failure.—Another class of deaths due to failure of circulation merit attention. These are cases without any distinctive signs during the acute stage of the fever, though there is often a tendency to diarrhoea and sweating, and sometimes melæna. But when the end of the third week arrives, and convalescence is expected, the temperature though it falls to 100°, or thereabouts, keeps daily above the normal—100° to 101° or so; the pulse may get rapid, rising 140 to 160 a minute, and though the tongue is clean, food well taken, and the special signs of typhoid fever have passed off, the patient fails to gain ground; he gets daily weaker and more and more listless, though the mental state is usually clear; profuse sweats may occur, and the vessels are wanting in tone; later on much tremor occurs from debility, and the patient usually dies, sometimes quite suddenly, sometimes merely flickers out. In other cases he may perhaps recover. Several such cases have been under my care. The symptoms differ from those usually attributed to toxæmia; and they cannot be attributed to heart failure,

1. Because myocardial change to such a degree as to lead to complete cardiac failure and death does not occur after a moderate fever of two or three weeks' duration, even in rheumatic fever, in which myocardial changes are specially marked.

2. Because the symptoms differ from those of cardiac failure; a failing heart tends to become slower, but in these cases the pulse often gets more and more rapid; the heart sounds and impulse too are not greatly weakened, though as the debility of the patient increases their debility increases with it; the extreme listlessness and debility of the patient are quite out of proportion to the heart weakness.

It appears most probable that in these cases there is an actual want of blood in the body, arising from failure in the blood-making function, and that death occurs from mere bloodlessness; I do not use the term *anæmia*, which would be strictly applicable, because it has become customary to use that term as applying only to the corpuscles.

The proof that symptoms in typhoid are sometimes attributable to want of blood are:

1. If the artery of a patient dead of typhoid fever after these symptoms be opened it will be found singularly empty of blood. In one case (where no melæna had occurred) it was quite empty and the inner wall of the artery quite dry. In some necropsies of typhoid patients, too, even where no melæna has occurred, the intestines are markedly blanched and exsanguined and the solid viscera pale and dry, though in most cases they are, as usual in febrile conditions, dark and softened.

2. Examination of the blood itself. Much less attention has been devoted to changes in the plasma and volume of the blood than to the changes in the corpuscles. Observers agree that there is an enormous decrease in the number of red corpuscles and in hæmoglobin throughout the course of typhoid fever which should be gradually made good in convalescence. There is also a great decrease in leucocytes, and Coe has found that the fibrin of the blood is

diminished in typhoid fever, the exact contrary of what occurs in other fevers. Hayem, too, finds that in typhoid fever there are sudden rises in the number of corpuscles which he can only attribute to losses of fluid by sweating and diarrhoea. All these results go to show that grave deterioration in all the constituents and in the quantity of the blood occurs in typhoid fever. Further, Dr. Thayer and others describe cases of post-typhoid anæmia, sometimes fatal, coming on as the typhoid passes off; it is admissible to think that if such fatal anæmia can immediately follow convalescence, it may occur at a time when convalescence commences, and prevent its accomplishment.

3. The symptoms of rapid pulse, increasing listlessness and debility, with fever and perfectly clear mental condition, are just those which follow want of blood, and indeed they often come on after severe hæmorrhage in typhoid, and after hæmorrhages from other causes.

And if it be conceded, as I think it must, that many of the symptoms and dangers of typhoid are due to changes in the constitution and amount of blood in the body, there is no lack of causes in typhoid fever to produce blood deterioration. To the causes of blood deficiency, pyrexia, toxins in the blood, and cloudy swelling of the internal organs which are common to the fevers generally, there are added in typhoid a special affection of the great blood-making organ, the spleen, and very often, also, a drain of material from diarrhoea, hæmorrhage, or profuse sweats. In addition, the widespread affection of the mesenteric and retroperitoneal glands must offer an obstruction to the absorption of nutriment from the alimentary canal, however well it is taken and digested. And I think the conclusion is justified that a certain number of typhoid patients die from mere bloodlessness and failure to make good the loss of blood material which results from the disease.

Why some patients should pass into this state and others not appears to depend more on the individual incapacity to make good the depletion by a new formation of blood than on any special character of the typhoid poison. Doubtless this defect in blood formation is in some cases associated with real cardiac weakness, but the cardiac weakness in many such cases is a result of it and not the cause. And the importance of measures to keep up the blood supply cannot be overestimated.

Another condition also seems not uncommon in typhoid fever—namely, a tendency for the venous side of the circulation to be full of blood while the arteries are unduly empty; in opening veins for injection of salines I have often been struck with the free bleeding from both proximal and distal ends of the vein, and care must be taken, if it is wished to avoid this, to put a ligature round the vein before it is opened, tightening it round the vein and cannula after the latter is inserted. Hayem has found, in examining the blood of typhoid patients, that changes which he attributes to loss of fluid by sweats and diarrhoea, occur in it suddenly, "such as seen in the algide stage of cholera," in which disease there is a tendency to accumulation of blood in the veins; and I think the overloading of the venous side in some typhoid (and other fever cases) is worthy of note, and among other things may predispose to thrombosis.

As regards treatment, it will be convenient to speak of heart failure, asthenia, and blood failure together. For all of them one essential is to prevent waste of material by diarrhoea, or hæmorrhage, or profuse sweats, and to supply as much nourishment as can be digested and absorbed.

As regards food, Dr. Barrs¹³ recommends the giving of solid food whenever the patient "likes it, wishes for it, and enjoys it." He argues that such food will not be likely to cause perforation, as it becomes softened before getting into the intestine, and that perforation only occurs in 2 or 3 per cent. of cases. I think cases do occur with little or no intestinal ulceration in which solid food might be given early without ill effect, but in the absence of certain means of distinguishing such cases, and remembering unfortunate results which have occurred in some patients who have surreptitiously followed Dr. Barrs's treatment, I think liquid food is the only justifiable mode of nourishment in the active stage of typhoid fever. And of all foods milk should be the staple diet, diluted with water, lime water, or barley water, according to the wish of the patient and the discretion of the medical attendant.

Where milk so diluted fails it will be sometimes better taken peptonised; in some cases it is borne well when diluted with tea, and there seems no reason to deny a typhoid patient either tea or coffee. We give caffeine for the heart, tannin to check diarrhoea, and allow sugar and water to be taken; why should a mixture of these be disallowed?

In some cases of vomiting or diarrhoea beef-tea or meat extracts have to be substituted for milk. The quantity of milk given should be as much as the patient can digest without flatulence or colicky pain or curds in the stools. As a rule three pints is enough in the twenty-four hours, but I can see no reason why more should not be given if the patient can take it well; persons often take five or six pints of milk a day under the Weir-Mitchell treatment, and a patient wasting and losing blood rapidly, as in typhoid, will benefit by as much as he can take without digestive disturbance.

As regards stimulants, I have nothing to add to the rules usually followed; they are not necessary as a routine part of treatment.

An important point is to secure sleep, and opium is quite admissible. Sir W. Broadbent deprecates the use of chloral. In the treatment specially directed to cardiac weakness digitalis, strychnine, and caffeine are to be resorted to, but cardiac tonics receive little response from a heart with organically-weakened muscle tissue; caffeine and strychnine act well together, but strychnine here, as in other conditions, may easily be pushed so far as to make the patient wakeful. Ether, ammonia, and sumbul, and diffusible stimulants are often of use. Alcohol, no doubt, stimulates the heart, but if pressed too much has the contrary effect.

In all cases, and specially in some, endeavours should be made to keep up the volume and composition of the blood; sweats should be checked by belladonna or oxide of zinc or agaric acid, and diarrhoea by enemas. There is no reason why cold water should not be given freely, due regard being had to preventing abdominal distension. In many cases ice sponging or the cold bath contracts the vessels and lessens their relaxation for some time after its use, and possibly in the bath or the tank fluid is actually absorbed cutaneously. Raw meat juice and every form of nutriment possible must be given in cases of gradual weakness from blood deterioration, and I have sometimes given iron and malt extract. Oxygen inhalations are also of use. Solid food must be given as soon as it can be taken with safety; after three days without fever at any time in the twenty-four hours it may be given; it is best to intermit one day at first to ascertain if a rise of temperature is produced. In cases where it is deemed probable that little ulceration occurred I give solid food earlier than in other cases. When cases drag on with little rises above the normal, solid food must be tentatively and gradually tried. In convalescence patients are hungry and require feeding up, and it is often in cases where solid food is long delayed that thrombosed veins occur.

When there are evidences of profound bloodlessness and weakness, whether following hæmorrhage or not, saline injections may be of use. Personally I have found the subcutaneous injection of salines gives so much pain that I prefer injecting the saline fluid into the basilic vein. This flows in better with a simple cannula, drainage tube and funnel than pumped in by the double action syringe made for the purpose. I have injected two pints, usually at a temperature of from 100° to 115°. After it there is usually sweating and rise of temperature temporarily. I believe the fluid should not be injected at so high a temperature as 105° in typhoid fever. In 2 cases I think the patients' recovery was due to it. In one of them, in 1896, the patient was in an apparently hopeless state of collapse, but recovered after two saline injections. Probably blood transfusion would be better, for in most cases saline injections only give temporary benefit.

Perforation may be accompanied by its well-known symptoms, but in some cases of typhoid where nerve tone is already lost, and the tympanitic belly is soft and doughy, perforation and after-peritonitis may occur most insidiously, with little pain, collapse signs, or alterations in temperature.

Cases of recorded recovery from perforation based merely on evidence of symptoms cannot be taken as conclusive, for it may be simulated by rupture of peritonæum over a mesenteric gland and other causes; indeed the abdomen has been opened for perforation in several cases where it had not oc-

curring. In Herringham's¹⁴ case nothing was found and the patient recovered; in Sheild's case¹⁵ rupture of gall bladder. In a case, however, recorded by Murchison a perforation had all but healed when erysipelas carried off the patient, and in a case at St. Mary's Hospital in 1894 a *post-mortem* examination showed a perforation all but healed when the patient succumbed to septicæmia. Three cases of 22 recorded operations for perforated typhoid ulcers have recovered, and as operation affords a better chance of recovery than that offered by Nature, it appears to me indicated in all cases in which perforation can be diagnosed—the difficulty in the diagnosis, however, will prevent the operation being very often performed. The only other treatment affording a chance of recovery is opium.

Hæmorrhage.—Some have regarded hæmorrhage as a favourable occurrence in typhoid fever, and there is no amount of hæmorrhage that may not be recovered from; in 1882 I saw, with Dr. Wakefield, a patient who bled two chambers full, and recovered. But the more general opinion is that it is a bad omen, and that death follows 30 or 40 per cent. of cases of free hæmorrhage, though sometimes long after the bleeding has occurred. It is generally arterial, but in a case of mine in 1888 the necropsy showed that the dark blood had oozed away during life from a vein in the ileum, opened up by ulceration. In its treatment opium pressed freely, and turpentine, are useful, but often no drug is so efficacious as tincture of hamamelis; in one case at St. Mary's Hospital in 1892 (J. M., Case 1398) it checked the hæmorrhage after all other remedies had failed, and I believe it is more to be relied upon than anything else; in 5-minim doses in a little water every half-hour while the hæmorrhage lasts, with or without opium. The application of the icebag to the abdomen seems beneficial, but it is depressing if kept on long after the hæmorrhage has ceased and it freezes the abdominal wall into a leathery consistence, and probably interferes with the vitality of the subjacent intestine. I have repeatedly seen hæmorrhage occur while the icebag had been on for days. The necessity for absolute non-movement of the patient when hæmorrhage occurs is obvious.

Of peritonitis without perforation opium is indicated.

Diarrhœa.—Murchison in the edition of his work in 1862 wrote that diarrhœa occurred in 96 of 100 cases; in 1884 he found it reduced to 80 of 100 cases; since that time it is much less frequent. Of 200 consecutive cases at St. Mary's diarrhœa occurred in only 115; constipation in 48; and the diarrhœa was seldom very severe. In many cases it had been set up by a purge given before a diagnosis was made. Diarrhœa is found in a very large proportion of cases in whom hæmorrhage occurs, and in my opinion adds to the danger of typhoid fever by preventing absorption of nutriment and by draining the blood of fluid. I think purgative drugs should never be given and diarrhœa should be checked in every way possible by adapting the nature and quantity of food to the patient's powers, by giving all food warmed and by enemata of starch with or without opium rather than by drugs given by the mouth. Of the latter, salicylate of bismuth in doses of 20 or 30 gr. doses three or four times a day is very useful.

Constipation must be treated if obstinate; enemata are better than purgative drugs for this purpose, and should have some disinfectant added to them. It is very important to overcome constipation before solid food is given; if not, it often sends up the temperature and an immediate recurrence of symptoms. Some attribute relapses to the giving of solid food, others to constipation. Certainly the two combined will produce a rise of temperature and disturbance for a considerable time, though not, I think, a true relapse with fresh spots.

Tympanites is commonest when there is much diarrhœa. That form in which the belly is soft and doughy as well as swelled is due to general loss of nerve tone and treatment affects it little. Diffusible stimulants, etc., may be tried. In the form of tympanites with tense abdomen hot fomentations are, I think, preferable to the icebag. Passage of a long tube into the rectum gives relief and often also sets up retention of urine.

In many cases swelling of abdomen is due not to true tympanites but to stomach distension, the result of liquid food and the recumbent position. Food must be given in small quantities only.

Retention of urine must be always remembered—it is often

overlooked in the general distension of abdomen, or because "overflow" occurs from the full bladder. Long retention may add uræmic dangers in a typhoid patient very readily.

The question of moving a patient in cases where the sanitary conditions are bad has sometimes to be considered, for he will be likely if he remain where he is to take in fresh doses of the poison. On the other hand, as Sir W. Jenner pointed out, the cases that do worst owe it sometimes to having travelled when well on in the disease. If it is very early, careful removal will be best. If the patient is well on in the disease he had better remain where he is as the risk of moving is too great. I show the chart of a patient I saw in June with Dr. Pettifer who had repeated rigors and rise of temperature pointing to septic conditions. We moved him; only one more rigor occurred and he recovered.

In the third group the intercurrent affections of typhoid such as pneumonia are to be treated on ordinary principles; pneumonia is usually recovered from, and it is apt to begin or to end in the course of typhoid with abrupt rises of temperature or a critical fall, after which the typhoid temperature reasserts itself. Acute bronchitis in typhoid is, in my experience, more serious than pneumonia.

REFERENCES.

- ¹ *Sem. Méd.*, March, 1890. ² *Trans. Clin. Soc.*, vol. xxiv. ³ *Ibid.*, vol. xxx. ⁴ *Munch. med. Woch.*, 1891, p. 43. ⁵ *The Cold Bath Treatment*, 1898. ⁶ *Lancet*, vol. ii, 1894. ⁷ *Guy's Hospital Reports*, 1893, vol. i. ⁸ *St. Thomas's Hospital Reports*, 1878. ⁹ *Practitioner*, 1897. ¹⁰ *Treatment of Typhoid Fever*, 1892. ¹¹ *Coe, The Blood*, 1898. ¹² *Johns Hopkins Hospital Reports*, vol. iv, No. 1. ¹³ *BRITISH MEDICAL JOURNAL*, vol. i, 1897. ¹⁴ *Trans. Med.-Chir. Soc.*, vol. lxxx. ¹⁵ *Lancet*, 1895.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, ETC.

NOTE UPON THE TREATMENT OF PNEUMONIA BY IRON PERCHLORIDE.

STEEL in the treatment of pneumonia has already many advocates, and I should like to add my testimony to its usefulness in certain forms of this rather common malady and under certain conditions. The following is an example of the kind of case in which it seems to be especially valuable: P. J. L., aged 34, came under treatment on October 18th, 1898. He was a fairly healthy looking man. He stated that he had been feeling very poorly for two or three weeks, and that on the day previous he had caught cold from getting wet while on a short railway journey. When seen on the afternoon of October 18th he complained of a tight feeling across the lower part of the chest, pain in the right side of his chest, back, and legs, and of a rather troublesome cough. There was found dullness above and below the right clavicle, bronchial breathing and bronchophony, and, in fact, the usual symptoms of apical pneumonia. Lower down there were crackling and subcrepitant *râles* to the sixth rib in front and moist sounds were also heard rather extensively posteriorly. The temperature was 102.2°, the pulse 118, and respirations 28. He was ordered carbonate of ammonia and citrate of potash in rather full doses every three hours, and hot applications occasionally to relieve pain over the right side. The following morning the temperature was 103.2°, pulse 124, and respirations 32. The tongue was still furred, the skin moist, pleuritic pains severe on moving or coughing. He had had very little sleep, and still felt very poorly. In the evening the temperature was 104.4°, pulse 126, and respirations 30; the moist sounds in the chest seemed to be extending; he felt, and evidently was, very seriously ill.

The question was whether to give a large dose of digitalis rather as a sedative than otherwise, as the first sound was still quite clear, or to give iron perchloride. I decided to give the latter, as I felt sure it would lessen secretion in the affected bronchioles, and act as a general tonic. The following mixture was prescribed: Tinct. ferri perchlor. mxx ; liq. am. acet., $\mathfrak{zj}$; sp. ether chlor. mxx ; aquam ad $\mathfrak{zj}$, to be taken every three hours.

On the following morning he said he felt much better; he had had two or three hours' sleep; his pulse was firmer, and