actively engaged, and in many diseases of the menstrual organ play a very important part.

As regards the adaptability of the mucosa tissue, it is chiefly of interest as evidence of the archaic nature of that tissue. The devastation witnessed within the uterus when the mucosa is denuded is astonishing, and yet a few days afterwards it is again possessed of glands, blood vessels, and superficial epithelium. This happens every month, and it is a remarkable feat of which no other organ in the mammalian body is capable. Moreover, it is noticeable that by far the larger number of capillary blood vessels, together with some of the epithelium, is formed from the indifferent elements of the mucosa—a remnant, it seems, of that power of reproducing lost parts which some of the lower animals possess.

A review of the various modern views of menstruation in the human female leads to the opinion that it will be found that while the histological phenomena accompanying menstruation in the latter may differ somewhat in detail from those in monkeys, they are practically identical processes; further, menstruation and "heat" are very much more similar than has hitherto been recognised.

With regard to the relation of ovulation to menstruation, if it is granted that menstruation in monkeys is the same process as menstruation in woman, then it can definitely be stated that ovulation does not occur during each menstrual period, and that it does not necessarily occur during any menstrual period. In *S. entellus* and *M. rhesus*² I have shown that menstruation frequently occurs without ovulation—probably this is the case during the whole of the non-breeding season; while Leopold and Miranoff have shown that ovulation may occur independently of menstruation in the breeding human female.

The want of power to ovulate is no doubt one of the causes of sterility, and the importance of all considerations dealing with the process of ovulation is obvious to those who have to combat sterility. It is a fuller knowledge of the cause of ovulation which is required, and it is hoped it will be forthcoming and be found of assistance to those who desire to induce ovulation.

NOTES.

¹ The various stages may be followed with the help of the drawings in Mr. Heape's paper on "The Menstruation of *Semnopithecus entellus*" (*Phil Trans.*, vol. clxxxv, 1894.) ² The Menstruation and Ovulation of *Macacus Rhesus*, *Phil Trans.*, vol. clxxxviii, 1897.

AXIS TRACTION WITH ORDINARY FORCEPS.

By JOHN F. LE PAGE, M.D.

THE method of delivery with traction in the axis recommended by Dr. T. Archibald Dukes in the *BRITISH MEDICAL JOURNAL* of November 5th as the "most powerful method" is open to criticism. I premise that axis traction has for one object the avoidance of "powerful" methods. *Arte, non vi* should be the watchword of every obstetrician. I take exception to his directions; "Keep both arms the whole time rigid and extended," "and pull with your back," "Fancy pulling an oar or anything else without using the back!" "One's whole strength can, when necessary, be exerted in urging the child's head in any desired direction," and the nurse is to "keep the patient on the bed." The last quotation has my unqualified approval, for it is an instruction which would commend itself as one of imperative necessity. It is, indeed, marvellous that with all this "powerful" traction, "no compression need be used unless the forceps are beginning to slip." As a matter of fact, this is certainly not so, but considerable compression would be necessary to prevent slipping whilst such Herculean efforts were in progress. It is erroneous to write, "let the handles assume their natural position close to the symphysis pubis," for, with the head above the brim, when the blades are in position the forceps are against the perineum, and not the os pubis.

Dr. Dukes expresses the opinion that "any intelligent man would use the instruments in the above way unless he were told to act otherwise." The intelligent man who did follow this method would, I am afraid, deserve censure.

I submit that, in view of the fact that our profession embraces thousands of young men who have not yet had much midwifery experience, and so have not had the material from which to evolve just and sound individual

opinions, but who still do as they have been taught because they were so taught, the foregoing advice is dangerous in the extreme, and, as contrary to the principles of the art, should be earnestly denounced.

Tactile dexterity in the conduct of instrumental labour, and not powerful traction, is an inevitable outcome of the advance in the science of midwifery during the past half century; an advance which leaves no excuse for conducting labour in an heroic manner, but which demands, as a precise art, the observance of scientific principles.

To denounce axis traction would be to repudiate a law in mechanics which is absolutely inviolable and unalterable; that is, that the direction of applied force, whether it be a *vis a tergo*, or a *vis a fronte* is always a straight line, whatever the shape of the instrument through which it is conveyed. The line of traction is not along the curve of the blades, but in an imaginary line from near their points to near the extremity of the handle. Hence the introduction of axis-traction forceps, which, by a second movable handle, allow of traction in a more backward direction. The arrangement advised by Dr. Dukes is not an axis-tractor, but a means by which the forceps are simply converted into a lever of the third class, the head being the weight, the end of the handle the fulcrum, and the lock the power. His directions are to "push with the right hand and pull with the left," that is to push the handle upwards and forwards, and pull the lock backwards. This is leverage, and is not admissible except for special purposes, such as lifting the head off the brim.

I entirely agree that axis-traction forceps do in many cases, especially in primiparæ, cause considerable pressure on the perineum. It is frequently quite impossible with them to use traction in the axis of the brim; and it was to obviate this pressure, and to allow of traction at any angle, and to supply a very ready and effective means of converting any ordinary long forceps in a moment, when *in situ*, into an axis-traction forceps, which overcome all the disadvantages of Tarnier's, Galabin's, and others, that my "axis tractor" was made some four years ago.¹

The figure shows approximately the relations of the long forceps, head, and pelvis. The line *A B* indicates the true axis, and *A D* the direction of power used with the forceps; *A E* is the line of traction with the axis tractor, and it is obvious that by using the handle of the forceps as a guide, the whole of the force expended may be directed in the true axis *A B*; whereas traction with forceps alone is along the line *A D*, and is necessarily expended to a very considerable degree on the anterior

brim, the effective force being far less than half the amount which is used. In the axis A B there is no loss of power, and risk of any injury to the structures over the pubic bones is reduced to a minimum. Although in its ordinary use this attachment is a tractor, it is possible with ease to lift a head over the pubic brim in pendulous abdomen (F S) and adjust it to the brim, or to the diameters of a deformed pelvis. In brief, by combined use of handle and tractor, the head can be moved into any position, and traction made in any direction with the greatest ease and precision. And whilst the lock is allowed considerable play, so that continuous pressure of the head is avoided, the force requisite for delivery is, in practice, found to be little, for it may be entirely used in the line of least resistance.

If instrumental delivery is conducted with precise knowledge of the mechanism of natural labour and of the direction of natural force—which is not downwards, but backwards into the hollow of the sacrum—and the instrument used is capable of accurately supplementing that force, and is, in its slightest movement, absolutely under the control of the operator, all difficulty disappears and with it the excuse for an undue and dangerous expenditure of force on parts which were never by Nature intended to suffer such abuse.

REFERENCE.

By Messrs. R. Sumner and Co., of Liverpool, by whom I am informed that many hundreds are now in use in this country.

DYSTOCIA DUE TO ANTE-PARTUM HOURGLASS CONTRACTION OF UTERUS: CÆSAREAN SECTION.

By COURTENAY C. WEEKS, M.R.C.S., L.R.C.P. LOND.,
Pinchbeck, Spalding.

HISTORY.

On July 29th, 1897, I was called to Mrs. B., aged 25 years, who was in labour with her second child. The pains occurred on and off for two days, and the "waters had broken" shortly before my arrival. On making an abdominal examination one was at once struck by the peculiar contour of the abdomen. It seemed to be occupied by two distinct tumours, one in its upper and the other in its lower part. The lower seemed to be about the size and shape of a child's head; the upper, larger, had its long axis transversely, and protruded very much forward. Between these two swellings a distinct groove could be made out; foetal limbs could not be palpated. The two swellings seemed (as the patient lay on her back) to occupy different planes, and suggested the possibility of twins.

Vaginal examination revealed the vertex presenting with occiput posterior. The os was well dilated, and the head just above the promontory of the sacrum, which was somewhat though not excessively prominent. An attempt was at once made to rectify the malposition, but quite failed. It was then noted that although the pains were strong and regular, and the uterus could be felt to contract, that absolutely no descent of the head took place. It seemed to be "strung up" over the promontory of the sacrum. Four hours after seeing the patient I gave chloroform, and made a further examination of the uterus preparatory to application of the forceps. The vagina and os seemed normal; the latter was well dilated. On passing the hand up by the side of the foetal head—(this was done with difficulty owing to the firm contraction of the lower segment of the uterus)—I found that the neck was firmly grasped by the uterus, which was powerfully contracted at this point. At first it was absolutely impossible to pass beyond this contraction. Bimanually the lower segment of the uterus was felt to be thinned, but not markedly so. It was also demonstrated that the sulcus, felt abdominally, corresponded with the contraction felt internally.

Forceps were applied, but absolutely no descent was produced. After a further interval I sent for Mr. S. H. Perry, and later for Dr. M. Perry, and from first to last we endeavoured either to rectify the malposition or deliver by forceps or version. All our efforts failed, even under deep chloroform anaesthesia and chloral hydrate. The pains throughout these manipulations kept very strong, and somewhat rapidly the

patient's condition became so serious that we decided that the uterus must be emptied.

OPERATION.

Craniotomy was resorted to but without any success. Cæsarean section was then decided upon sixteen hours after I first found the os fully dilated, and not until we fully agreed that it was the only chance for the patient. Chloroform was administered by Dr. Perry. The skin was rapidly prepared, and a very thorough mercurial vaginal douche given after one final attempt at forceps delivery. The abdomen was opened by a median incision, and the uterus exposed. Sponges wrung out of warm boracic lotion were then placed all round the uterus, and it was incised, whilst Mr. S. H. Perry placed his hands well down on each side of the uterus in the pelvis, and so most effectually and thoroughly controlled the hæmorrhage. The child was quickly delivered, and was rapidly followed by placenta and membrane; a large sponge was then placed for a few seconds in the uterine cavity, and firm pressure was kept up whilst the uterus contracted, which it did very well. No cause could be ascertained for the contraction; the uterus was a dark, slaty, blue colour. A double row of sutures—one taking up the whole of the thickness of the uterine wall as far as the submucosa, and the other taking up only the peritoneal cover so as to cause infolding of the edges, were used. The abdominal wound was closed by one set of sutures only.

AFTER-HISTORY.

Immediately after the operation strychnine and ergotine were given hypodermically, and soon after 15 c.cm. of anti-streptococcus serum, the operation being done in a not over-clean cottage. There was a good deal of collapse, but the patient rallied well, and her temperature at night was 100° F., and was normal the following morning.

For fifty-two hours the patient did very well in every way, the highest temperature being 101.4° at the end of forty-eight hours. There was practically no pain, no vomiting, no distension, flatus passed *per anum*, the tongue was clean and moist, respiration was abdominal, and the facial aspect good. She slept and felt and looked very comfortable. On the third day she had a little cough, and after one coughing attack (lungs appeared normal) she complained of severe abdominal pain, became very collapsed, and vomited a little bile-stained fluid. The pulse was very rapid, feet and hands were cold. There had been no chills or rigor, and now the patient complained simply of pain, and asked for a "good dose of salts." Stimulants caused a slight rally, but only for a while; a few hours after this initial attack of pain the abdomen was slightly distended. The abdominal wound was then examined; a few stitches had given way in two places, allowing knuckles of omentum to protrude. These were reduced and a couple more stitches removed, so that the uterus might be examined for any hæmorrhage. This was fifty-nine hours after the operation, and we found the wound beginning to heal and an entire absence of any sign of hæmorrhage or inflammation.

We attributed the collapse to the nipped omentum. Various stimulants, rectal and axillary injections of saline solution, were tried, but the patient gradually sank thirty-two hours after the attack of pain, and eighty-four after the operation. The respiration towards the end was of a peculiar sighing character, and resembled that seen in collapse due to loss of blood. The lochia were established, and never had any bad odour. Vaginal douches were used from the first. There was no vomiting until the "acute attack of pain," and only twice very slightly afterwards. There were no rigors, diarrhoea, nor tenderness of abdomen. Hearts, lungs, and great blood vessels normal. The clinical picture was one of good progress for fifty-two hours, and then sudden and eventually fatal collapse.

POST-MORTEM EXAMINATION.

There were no signs of any peritonitis; the uterus was well contracted, and of a light bluish colour. In the right iliac fossa was an ileo-cæcal intussusception, reduced fairly easily; only slight extravasation of lymph. The adjacent coils of intestine were slightly injected and finely granular on the surface.

REMARKS.

The interest of this case naturally centres in the cause of the dystocia. That we had a very definite "contraction of a