recent investigations on cases of pernicious anaemia, in addition to the defect they present as efficient grinders of food. I may say, however, that I have had more difficulty with regard to this part of the treatment than with any other.

Whether the cure becomes permanent depends very much upon the future life of the patient. If the conditions which produce this state of the stomach are resumed, the dilatation and proptosis may occur again. In one of my cases, that of a man whose employment necessitated his standing all day, a recurrence after two years brought him back to me, and he has been greatly benefited by undergoing the treatment again after this lapse of time. In most of the cases, however, so very decided an improvement has resulted that I have been encouraged to bring the details before the profession.

SOME PHYSIOLOGICAL ASPECTS OF GASTRO-ENTEROSTOMY.

By HECTOR CHARLES CAMERON, M.A., M.B., B.C.,
MEDICAL REGISTRAR, GUY'S HOSPITAL MEDICAL SCHOOL;
FORMERLY DEMONSTRATOR OF PHYSIOLOGY,
GUY'S HOSPITAL MEDICAL SCHOOL.

[From the Guy's Hospital Physiological Laboratory. The expenses of this work were defrayed by a grant to Dr. Pembrey from the Royal Society.]

SINCE the work of Pawlow¹ and his fellow-workers in St. Petersburg, and of Bayliss² and Starling in London, has become widely known, the question—Is gastro-enterostomy a harmless operation?—has been asked with increasing frequency. These workers have given to us a new conception—the *continuity* of the process of digestion. So long as the process was looked upon as a series of isolated actions and reactions, each independent and separate, so long as pancreatic digestion was regarded merely as following gastric digestion, and in no sense occasioned by it or modified by it, so long might surgeons have little hesitation in performing such short-circuiting operations as gastro-enterostomy, pylorotomy, or partial gastrectomy. But when we learn that the adequate secretion of pancreatic juice is dependent upon the changes which the food has already suffered in the process of digestion—that is to say, upon gastric activity—then we may very well wonder to what extent the mutilation entailed by the “short-circuiting” of the duodenum impairs the stability of the whole process. By the operation a link in the chain has been broken. The resulting modification in the patient's powers of digestion and absorption is at any rate of considerable *theoretical* interest.

The first recorded experiments which show the digestive capacity of patients after gastro-enterostomy are those of Joslin.³ From results obtained in Ewald's laboratory Joslin stated that in two cancerous patients upon whom gastro-enterostomy had been performed, the digestion both of proteids and fats was greatly impaired. Quoting these results, Paterson,⁴ in his Hunterian Lectures on gastro-enterostomy, remarks, “Joslin's experiments have been used as an argument against the performance of gastro-jejunostomy, except as a last resource, and have had a considerable effect in retarding a more general use of the operation in gastric ulcer and other non-malignant affections of the stomach.” That a knowledge of Joslin's results has delayed or vetoed the performance of the operation of gastro-enterostomy in a single instance I take leave to doubt. Surgeons are rightly in the habit of estimating the success or failure of their work by the measure of their own experience. The knowledge that an infant upon whom gastro-enterostomy has been performed for congenital pyloric stenosis may attain to a healthy and vigorous manhood, the evidence of the increasing weight of patients upon whom they themselves have operated, will far outweigh the results of such experiments as Joslin's, however indubitable and however consistent their results may be.

Therefore, I say that the result of such experiments is of only *theoretical* interest. It is foolish to tell a surgeon that gastro-enterostomy can no longer be considered a harmless operation. Tracheotomy is not a harmless operation, and yet it has its usefulness. The question cannot be decided in the abstract, but it must be answered for each individual case. A man with a gastro-

enterostomy may not be as good a man as his neighbour; and yet he may be in far better case than before the operation. There are, indeed, surgeons whom it may be salutary to remind that the operation does entail some alteration of function—surgeons who seem to regard the new relation of stomach and bowel as though it were an alternative scheme of Nature to be substituted as quickly as possible whenever the old arrangement shows signs of disorder—but the majority stand in need of no such advice.

The progress of surgery is, in a sense, retarded because of the priceless value of the material with which it deals. Advance must be made by the method of observation alone, for the method of experiment is to a great extent denied to it, and rival theories—the true and the false, the wheat and the tares—must grow together until the harvest. In other branches of science theory must precede experiment. In surgery, theory must come later—framed to accord with observation. It is true that experiments upon animals, in surgery as elsewhere, have increased our knowledge. In the study of gastro-enterostomy, the work of Cannon and Blake, and more recently of Legett and Maury, has been of great value and utility. Yet surgeons are rightly chary of transferring the results obtained from animals to the precious metal of humanity.

The theoretical considerations which follow have been brought forward not so much to suggest possible alterations in surgical procedure, as to ascertain to what extent theory and practice coincide, and to attempt, in the great variety of conditions relieved by gastro-enterostomy, to find the explanation of the benefits conferred in each. For instance, the experiments quoted below show that patients after gastro-enterostomy have complete power of digesting a mixed diet rich in fat, while their capacity for digesting a purely milk diet is considerably diminished. Apparently the experience of surgeons has led them to appreciate this difference, for it is their usual practice to abandon a purely milk diet as quickly as possible after the operation, usually after two or three days at the most.

Again, it will be maintained that the benefit of gastro-enterostomy in acute ulceration of the stomach is due to this—that the operation permits the regurgitation of alkaline duodenal juices into the stomach, and thus diminishes the gastric acidity; and it will be argued on theoretical grounds that most regurgitation will occur and the maximum benefit be conferred where the simple no-loop method is adopted, and entero-enterostomy, or the Y-shaped junction, avoided. In practice, guided by observation of the results obtained, since January, 1905, the Mayos have entirely given up the more complicated methods in these cases, and always perform the simple no-loop method.

The following points may be discussed in order:

1. The abnormal course followed by the food.
2. The removal of the valvular action of the pylorus in regulating the passage of the gastric contents into the duodenum.
3. The regurgitation of alkaline duodenal juices through the artificial opening.
4. The changes in digestion and absorption brought about by the operation.

1. The Abnormal Course followed by the Food.

In cases of obstruction to the pyloric outlet there can be no doubt that the food leaves the stomach by the artificial opening. The same is, of course, true where the gastro-enterostomy follows a pylorotomy or partial gastrectomy, unless an end-to-end anastomosis has been made. Where there is no obstruction to the outlet at the pylorus, it is in the highest degree improbable that the food deviates from the normal course. Cannon,⁵ in cats upon which gastro-jejunostomy had been performed, proved by the bismuth and x-ray method that in these animals the food took its normal course. In attempting to confirm these observations upon patients in Guy's Hospital I met with the difficulty that, although very large quantities of bismuth subcarbonate (3ij) were given, I was never able, in any position of the patient, to photograph the duodenum, nor even to identify it on the screen with enough precision and certainty. Clinical observation, however, strongly confirms Cannon's conclusions. Thus, a man suffered from a

duodenal fistula, which refused to close. A posterior gastro-jejunostomy was performed, but the fistula refused to heal, and the intestinal contents still escaped. A second laparotomy was performed; the pylorus was ligatured, and the fistula healed almost immediately. It may be concluded almost with certainty that no opening, however large or however placed, will divert any considerable amount of the gastric contents so long as the pylorus remains unobstructed. In cases where the pylorus was obstructed, Cannon and Blake occasionally observed that the food leaving the stomach passed for a short distance into the proximal loop, only to be swept back again into the stomach by a peristaltic wave. This movement, which at times was seen to be repeated again and again, could, however, at the most only result in a very transitory and insufficient contact of the acid chyme with duodenal mucous membrane, and must be regarded rather as an accident than as the normal condition.

More recently Legett and Maury⁶ have further investigated the course of food after gastro-enterostomy. To determine this point they have devised an ingenious plan. A dog was induced to swallow a bullet, which had been wrapped up in a piece of meat, and tied to a piece of string. In an instant the bullet, meat, and one end of the string were in the dog's stomach. The other end of the string was secured around the pterygo-maxillary ligament. The dog was killed some hours later, and the diagram (Fig. 1), copied from their paper, shows the

Fig. 1.

course adopted in one case: That is to say, the body A has passed out at the stoma, round the duodenum in the reverse direction, and through the pylorus, twice over. This experiment, though of great interest, certainly does not disprove Cannon's statement that with a patent pylorus the food does not pass out by the stoma. It has been known since Dr. Beaumont's observations on Alexis St. Martin that contact with a hard solid body causes spasm of the pylorus. The body A would, therefore, be forced, if it left the stomach at all, to leave by the stoma.

2. The Removal of the Pyloric Valves.

In cases with pyloric obstruction where the stomach is drained by the anastomotic opening. It is evident that the carefully adjusted and regular mechanism of the pyloric valvular apparatus is removed. Serdjukow has shown that while free acid in the stomach opens the valve, free acid in the duodenum closes it. So soon as free acid begins to accumulate in the stomach the pylorus relaxes and a "squirt" of acid chyme passes into the duodenum. The duodenum in consequence temporarily shows an acid reaction and the pylorus responds by becoming tightly closed. The acid in the duodenum then produces secretin in quantity just sufficient to summon to the duodenum just enough alkaline bile and pancreatic juice to neutralize the chyme which has escaped. Referring to this action of the sphincter of the pylorus Pawlow⁷ remarks:

This regulating action prevents disorder in the process of digestion and at the same time ensures regularity in the transition from the acid gastric to the alkaline intestinal digestion. If it were otherwise, and the acid contents of the stomach passed without control into the duodenum, the bile on mixing with it might arrest or greatly weaken the action of the pepsin, while the insufficient reduction of the acidity would interfere with the pancreatic ferment.

And yet it is just such an uncontrolled escape of the acid gastric contents which must take place in cases of gastro-enterostomy. It has, indeed, been stated more than once that in time something in the nature of a sphincter is developed around the artificial opening, but we can hardly believe that such a sphincter can be controlled by an elaborate local nervous reflex like that of the pylorus. Case 3 in the series described below is in this connexion interesting. Here no gastro-enterostomy, properly so-called, was performed, but the pylorus, and the pylorus only, was excised and an end-to-end anastomosis made. The patient, who suffered from a simple fibrous stricture of the pylorus, after the operation passed

26 to 28 per cent. of the fat of her milk diet undigested. The only change here introduced was the removal of the pyloric valvular apparatus, and the resulting defect may, I think, be explained in the words of Pawlow.

With the gastric contents escaping in this irregular way it appears, then, likely that the pancreatic secretion will be affected injuriously. With the pylorus working efficiently the stimulation of the pancreas is co-ordinated, long-continued, and at no time excessive; with the artificial opening draining the stomach and the pylorus obstructed it would seem that it could hardly help being inco-ordinated, explosive, and short lived, with the result that the activity of the pancreatic juice which is actually secreted in response to the abrupt stimulus is inhibited by the excess of acid present. It is probable, however, as a result of the regurgitation of alkaline duodenal juices through the anastomotic opening, to be discussed immediately, that the removal of this controlling apparatus becomes of only minor importance; for with this regurgitation the acidity of the gastric contents is considerably diminished, and the mechanism of pancreatic secretion in consequence profoundly changed.

3. The Regurgitation of Alkaline Duodenal Juices through the Artificial Opening.

To determine the presence or absence of this regurgitation through the artificial opening, Katzenstein⁸ has made an elaborate research upon dogs. In dogs, with gastro-jejunostomy performed, bile and pancreatic juice always find their way into the stomach at first continuously and later on after the ingestion of food. Their appearance in the stomach is hastened by the presence of fat in the food. In consequence of this admixture of alkaline intestinal juice the stomach acidity is reduced by a chemical reaction and also by reflex inhibition. That the introduction of alkalies into the stomach inhibits gastric juice reflexly—a fact first shown by Pawlow⁹ experimentally—is further proved by the observation that after gastro-enterostomy the total chlorides in the stomach are diminished.¹⁰ The hydrochloric acid-pepsin digestion is thus impaired by a gastro-enterostomy, and with a diminished or abolished acidity of the gastric contents there must be diminished manufacture of secretin in the duodenal mucous membrane. Clinical evidence also bears out the findings of Katzenstein. In a patient upon whom gastro-jejunostomy had been performed four months before, Katzenstein himself found bile and pancreatic juice in the stomach after an ordinary test meal. Kaiser¹¹ quotes numerous observers who found that after gastro-jejunostomy in human beings bile was almost invariably present in the stomach. Moreover, while it is true that peptic jejunal ulceration after a gastro-jejunostomy is from time to time recorded, the extreme rarity of these cases is in itself evidence that the gastric acidity is diminished. For if the acidity was normal, we should expect ulceration to be comparatively common as a result of the uncontrolled escape of the gastric contents through the artificial opening, and of the insufficient neutralization which would result. The practical importance is that in this regurgitation of alkaline duodenal contents, which has been proved experimentally by Katzenstein and others, lies the explanation of the utility of gastro-enterostomy in cases of peptic ulcer.

Fig. 2.—At A the bowel is divided and the ends closed.

Ulceration of the alimentary tract, as is well known, is confined to those regions where the reaction is acid, namely, the stomach, the first part of the duodenum, and with extreme rarity, as a sequel of gastro-jejunostomy, the jejunum just below the artificial opening. If the reaction of the stomach could be made entirely and permanently alkaline, the cure of gastric ulcer would be facilitated and relapse rendered in the highest degree unlikely. An operation upon the lines indicated by Fig. 2 would have the effect of depriving the stomach and the upper part of the duodenum of their individual acid reaction, and of

reducing them for good and all to the alkaline level of the rest of the alimentary canal.

After the performance of a gastro-jejunostomy in the usual way, the proximal loop has been cut across and the divided ends closed. In this way the bile and pancreatic juice are found to pass backwards through the pylorus in the reverse direction to reach the stomach, with the result that the stomach acidity is abolished. The operation is not a new one in surgery. Moynihan¹² has reported a case where he was forced to adopt this device because of the accidental rupture of the duodeno-jejunal flexure. The patient made a good recovery without vomiting or wasting. Tavel of Berne has reported a similar instance, quoted by Moynihan,¹³ where persistent vomiting after gastro-jejunostomy, unrelieved by several secondary operations and anastomoses, was finally cured by adopting this method. That the operation would be well borne by dogs is suggested by the work of Legett and Maury quoted above. It has, indeed, been performed many times and for many reasons, and the animals have lived and thrived after it. Stendel, Ohlumski, and Kelling, in investigations undertaken to explain the so-called "vicious circle," have all performed it. Kelling's dog was the only one to die directly from the result of the operation. He states that the food passed into the blind loop of the duodenum through the pylorus and extended it to such an extent that gastric contractions ceased, and the animal died of starvation. In Stendel's case the dog died on the eighteenth day of perforative peritonitis. Ohlumski's dog made a good recovery, and lived in apparently perfect health. For four weeks after the operation the animal was noticed to have excessive thirst, but this ultimately disappeared. All these operations were performed a considerable time ago—before the year 1899—and all are quoted in the *Annals of Surgery*, vol. xxxvi, p. 698.

Quite recently Maury has more than once performed the operation, and the dogs have lived. I would refer the reader to his interesting paper on death in high intestinal obstruction.¹⁴ When thorough and scientific medical treatment has failed to stop the process of gastric ulceration, and when, in addition, gastro-enterostomy has failed to relieve, an operation upon these lines is possibly indicated.

In the majority of cases, however, a simple gastro-enterostomy, devised so as to permit as much regurgitation of alkaline duodenal juices as possible, will bring relief or effect a cure. The testimony of surgeons upon this point is almost unanimous. It would, appear, however, that a distinction should be enforced between gastro-enterostomy done for drainage in pyloric obstruction and gastro-enterostomy done for acute ulceration. In the latter case the Y-shaped junction and secondary entero-enterostomies should be avoided, and the stoma should be placed as near the cardiac end as possible, when the pressure is low, in order to facilitate regurgitation as much as possible. When the operation is intended to provide drainage, on the other hand, as Cannon was the first to insist, the stoma should be near the pylorus. As mentioned above, it is interesting to note that since January 1st, 1905, the brothers Mayo have been led to abandon all the more complicated methods, and adopt only the simple no-loop operation, where this regurgitation is at a maximum.

That the vomiting which has occasionally followed gastro-jejunostomy is to be accounted for by the presence of bile and pancreatic juice in the stomach is rendered in the highest degree unlikely by the observations quoted above of Moynihan, Tavel, Kaiser, and Katzenstein. Equally striking is the statement of Cannon and Blake, that in their researches upon cats upon which gastro-jejunostomy had been performed:

Retention of food in the stomach, with subsequent repeated vomiting, such as attends the so-called "vicious circle," have been associated usually in our experiments with kinks and demonstrable obstacles to the easy passage of the food... this circulation of the food—that is, the passage from the stomach through the pylorus and duodenum back to the stomach, which they frequently observed—has never resulted in the clinical symptoms of a vicious circle.

4. The Changes in Digestion and Absorption brought about by the Operation.

Experiments on metabolism after gastro-enterostomy have been carried out by at least two observers. Joslin¹⁵ of Boston, working in Ewald's laboratory, published the

results of observations upon two patients with gastro-jejunostomy, performed for malignant disease, with pyloric occlusion.

The first case was that of a woman, aged 45, from whom two-thirds of the stomach had been removed for carcinoma. The ends of the duodenum and stomach were closed and gastro-jejunostomy performed. Her metabolism on a mixed diet was investigated. Analysis showed the following result:

	Absorbed.		Unabsorbed.	
	Per Cent.		Per Cent.	
Fat	81.3		18.7	
Nitrogen	87.4		12.6	

The subject of the second experiment was a man upon whom gastro-jejunostomy had been performed for carcinoma of the pylorus:

	Absorbed.		Unabsorbed.	
	Per Cent.		Per Cent.	
Fat	68.79		31.21	
Nitrogen	80.05		19.95	

In the second experiment a control observation was made before the operation, but its value is much diminished by the accident that the patient was vomiting frequently, and the estimate of the intake must necessarily have been extremely difficult.

In his Hunterian lectures on gastro-jejunostomy, delivered at the Royal College of Surgeons in 1906, Paterson¹⁶ criticizes these observations, and states that, in his opinion, their value is seriously diminished by the circumstance that the patients were suffering from cancer. He maintains that diminished power of absorption and digestion is only to be expected in cases of carcinoma, and that such observations cannot be accepted as evidence concerning the effect of the operation on the digestion and absorption of patients who are not suffering from malignant disease. Paterson himself repeated the experiments in four non-malignant cases. Few details as to the methods adopted are given. The patients were fed on a mixed diet. Tabulated, his cases show the following results:

	Period Since Operation.	Fat Absorbed.	Fat Unabsorbed.	Nitrogen Absorbed.	Nitrogen Unabsorbed.
		Per Cent.	Per Cent.	Per Cent.	Per Cent.
Case 1 ...	5 months	92.3	7.7	91.0	9.0
Case 2 ...	7 months	92.5	7.5	90.5	9.5
Case 3 ...	24 months	92.7	7.3	92.1	7.9
Case 4 ...	2 months	94.7	5.3	92.7	7.3

These figures show that in non-malignant cases the digestion and absorption of nitrogen and fat is impaired only to a very slight extent. Case 3 was one of almost complete stenosis of the pylorus. Cases 1, 2, and 4 were operated upon for gastric ulcer, for perforated gastric ulcer, and haematemesis respectively, and in none is any mention made of the existence of pyloric obstruction. Paterson himself answers the criticism that in Cases 1, 2, and 4 there is no evidence to show that the pylorus is not patent, and that the food does not take its old course, by the claim that in any case, whether the food leaves by the pylorus, by the anastomotic opening, or by both routes, his experiments show that digestion is not seriously impaired, while clinical evidence proves that the short-circuiting has a very beneficial effect upon the ulcer. Although the strength of this contention for practical purposes is obvious, it can hardly be maintained that it is a satisfactory answer to a question of such importance. It appeared certain, in view of the radical nature of the operation of gastro-enterostomy, that there must be some modification of the processes of digestion, and it was in search of evidence of such modification that the present series of experiments was undertaken. It was felt that merely to prove that the powers of absorption in these patients is unimpaired does not imply that there is no modification of the process of digestion. The results of digestion, in short, may be complete and yet the nature of the digestive process may be profoundly changed.

In all, seven cases have been under observation. Cases 1 and 4 had well-marked pyloric obstruction, not malignant, and in both cases there was a largely dilated stomach. Cases 2 and 5 had no obstruction at the pylorus, but haematemesis and indigestion. Case 3 had a simple fibrous stenosis of the pylorus; no gastro-jejunostomy was performed, but the pylorus was removed and the ends united by direct suture. Case 6 had a gastro-jejunostomy for simple fibrous pyloric stenosis eight years before. In Case 7 the pyloric half of the stomach was removed for malignant disease, the ends were closed, and a posterior gastro-jejunostomy was performed. In Cases 1, 2, and 3 the patients were fed entirely on milk; in Cases 4, 5, 6, and 7 a rich mixed diet was given—milk, cream, butter, eggs, bread, vegetables, fish, and lean meat; with tea or coffee—and the amount of fat in this diet was estimated in the following way: The quantity of fat in the milk, cream, and butter was calculated by extracting samples of known bulk or weight with ether in a Soxhlet apparatus. The amount of fat in the eggs was calculated from tables. Only fish with a very low percentage of fat was given (cod and haddock), and care was taken to remove all visible fat from the meat. With these precautions the amount of fat taken in food other than milk, cream, butter, and eggs was neglected. In Cases 1 and 2 both the nitrogenous and the fat metabolism was investigated; in the remaining cases only the fat. Attention was especially directed to the digestion and absorption of fat, because such a study provides the best means of estimating the activity of the pancreatic juice. It is true that on a milk diet, where the fat is presented in the form of an emulsion, the gastric lipase is able to split up almost the whole of it, but on a diet in which the fat consists of butter the lipase is ineffective, and the digestion must be the work of the steapsin of the pancreatic juice. By a study of the nitrogenous digestion and absorption, the activity of the pancreas cannot be estimated, because it is impossible to separate the effect of peptic digestion from that of the trypsin. In all cases a period of at least three days was allowed to pass after the special diet had been begun. The urine and faeces were collected from the fourth day onwards. The estimation was carried on for a period of at least five days. In Cases 1, 2, and 3 the nitrogen and fat in samples of the milk supplied were estimated twice—on the first day of the experiment, and again on the sixth day; the milk for the twenty-four hours was kept in a large flask, which was shaken before any feed was drawn off, to ensure that the sample was a true representative of the whole. The nitrogen in the urine and faeces in Cases 1 and 2 was estimated every day by Kjeldahl's method. In Case 3 the patient had some incontinence of urine, so that the estimation had to be abandoned. The total fat in the faeces for five days was estimated in all cases by extracting with ether in a Soxhlet apparatus weighed samples of the dried and powdered faeces, which had been collected for five days and mixed.

Table I (given for comparison) shows the results of similar experiments conducted by various observers upon normal persons; the last column represents the average of all cases. The diet in all was milk.

Table II, taken from Harley and Goodbody's *The Chemical Investigation of Gastric and Intestinal Diseases*, p. 133, gives

TABLE I.—The Normal Absorption of Fat on a Milk Diet.

Amount of Food.	Grams of Fat in Food.	Grams of Fat in Faeces.	Percentage of Fat in Faeces.	Percentage of Fat Absorbed.	Age and Sex.	Experimenter.
1,790 c.cm. milk ...	53.7	1.50	2.8	97.2	F. 12	Camerer
2,039 c.cm. milk ...	57.4	1.60	2.8	97.2	F. 10	Camerer
2,050 c.cm. milk ...	79.7	5.70	7.1	92.9	M. adult	Rubner
2,200 c.cm. milk ...	69.1	4.93	7.2	92.8	"	Müller
2,438 c.cm. milk ...	95.1	4.66	3.3	96.7	"	Rubner
3,075 c.cm. milk ...	119.9	6.70	5.6	94.4	"	Rubner
4,100 c.cm. milk ...	160.0	7.40	4.6	95.4	"	Rubner
3,000 c.cm. milk ...	94.2	6.57	6.9	93.1	"	Rubner
3,000 c.cm. milk ...	111.8	5.65	5.1	94.9	"	Prausnitz
2,632.4 c.cm. milk ...	93.4	4.97	5.04	94.96		

TABLE II.—The Normal Absorption of Nitrogen on a Milk Diet.

Number of Cases Examined.	Milk. Pints.	Average Number of Grams of Nitrogen in Faeces.	Percentage of Nitrogen Absorbed.
6	4	0.84	93.82
4	4½	0.76	95.03
2	5	1.03	93.76

similar information with regard to nitrogen absorption in normal individuals.

Cases 1, 2, and 3 in the present series gave the result shown in Table III, which may be compared with the previous tables.

Table III shows that the percentage of fat absorbed in cases examined shortly after the operation and kept upon a milk diet is distinctly less than normal (see note); and that this diminution is shown whether there is obstruction at the pylorus (Case 1) or whether there is no obstruction at the pylorus (Case 2). The percentage of unabsorbed nitrogen, on the other hand, is within normal limits. Case 3, which was examined as a control in which no gastro-enterostomy properly so-called was performed, and which has been referred to above, is interesting because the unabsorbed fat is the largest found—26.28 per cent. of the whole.

The digestion of milk, therefore, was faulty both in Case 1 where the pylorus was obstructed, and in Case 2 where the pylorus was not obstructed. In the latter case both experiments upon animals and clinical observation agree that there is no considerable escape of gastric contents through the artificial opening. In Case 1 it is certain that the stomach contents passed out by the artificial opening; in Case 2 we must conclude that they passed as

TABLE III.—Nitrogen and Fat Absorption in Cases 1, 2, and 3.

	Amount and Kind of Food per diem.	Grams of Fat in Food per diem.	Grams of Fat in Faeces per diem.	Percentage of Fat Absorbed.	Grams of Nitrogen in Food per diem.	Percentage of Nitrogen Absorbed.	Grams of Nitrogen in Urine per diem.	Weight of Dried Faeces per diem in Grams.	Age, Sex and Period since Operation to First Day of Experiment.	Clinical Notes.
1	1,700 c.cm. milk.	69.0	8.6	87.5	7.2	97.1	7.5	25.79	F. 40 years 20 days	Mr. Golding-Bird's case. Haematemesis at 23 years of age; since then constant pain in stomach; posterior gastro-jejunostomy; pylorus obstructed; stomach dilated.
2	2,835 c.cm. milk.	117.8	15.9	86.6	14.75	95.9	11.5	35.17	M. 53 years 11 days	Mr. Rowland's case. 2½ years' history of indigestion; haematemesis 16 months before posterior gastro-jejunostomy; pylorus not obstructed; stomach not dilated.
3	1,700 c.cm. milk.	67.0	17.7	73.72	—	—	—	7.74	F. 52 years 36 days	Mr. Lane's case. History of gastric pain for years; cicatricial contraction of pylorus found; pylorus excised and ends sutured together; no gastro-jejunostomy; microscopic report, "no malignant disease."

TABLE IV.—*Absorption of Fat in Cases 4 and 5 on a Mixed Diet Rich in Fat.*

The diet consisted of butter in large amount—about 3 oz. daily—milk, cream, eggs, cod or haddock, lean meat, bread, vegetables, and weak tea.

	Amount and Kind of Food.	Grams of Fat in Food per diem.	Grams of Fat in Faeces per diem.	Percentage of Fat Absorbed.	Weight in Grams of Dried Faeces.	Age and Sex. Period between Operation and First Day of Experiment	Clinical Notes.
4	Various	122.49	5.25	95.71	23.20	F. 43 years old. 18 days	Dr. French's case. Operation by Mr. Rowlands. Indigestion many years. Three ulcers, one on either side of pylorus. Posterior gastro-jejunostomy. Obstructed pylorus. Dilated stomach.
5	Various	201.3	7.00	98.57	41.31	M. 41 years old. 14 days	Mr. Lane's case. Old duodenal ulcer. Posterior gastro-jejunostomy. Pylorus not obstructed. Stomach not dilated.

TABLE V.—*Absorption of Fat in Cases 6 and 7 on a Similar Mixed Diet.*

	Amount and Kind of Food.	Grams of Fat in Food per diem.	Grams of Fat in Faeces per diem.	Percentage of Fat in Faeces.	Weight in Grams of Dried Faeces per diem.	Age, Sex, and Time between Operation and First Day of Experiment.	Clinical Notes.
6	Various	128.54	8.07	6.4	35.2	M. 68 years old. 8 years	Mr. Symonds's case. Had been operated upon for simple fibrous stricture of pylorus eight years before.
7	Various	158.40	5.00	3.13	12.6	F. 59 years old. 27 days	Mr. Swan's case at Cancer Hospital. Partial gastrectomy for cancer, and posterior gastro-jejunostomy.

before along the duodenum. It follows that the explanation of the deficient digestion of fat is not to be found in the "short-circuiting" of the duodenum and in the consequent want of contact between acid chyme and duodenal mucous membrane. The cause must be common to both, and is probably to be found in the regurgitation of alkaline intestinal juices through the artificial opening, with the consequent diminution of the acidity of the stomach both by chemical reaction and reflexly by inhibiting the secretion of acid gastric juice.

Cases 4 and 5 were fed upon a mixed diet containing a large amount of fat (*v. s. p.*). The analysis gave the results shown in Table IV, which may again be compared with Table I.

Case 4 had well marked pyloric obstruction. Case 5 had no obstruction at all, and again there is no marked difference in the result of analysis. Yet in one case the food passed over the duodenum, in the other it did not.

Comparing Cases 4 and 5, which were given a mixed diet rich in fat, with Cases 1 and 2, which were given milk only, it will be seen that while Cases 1 and 2 show a distinct deficiency in the power of dealing with fat, Cases 4 and 5 digested and absorbed larger quantities of fat with complete success. That the difference caused by the diet is not to be accounted for simply by the unappetizing character of a milk diet and the consequent diminution of the psychic appetite flow is, I think, proved by a reference to Table II, where the digestion of normal persons upon an equally unappetizing milk diet is shown. Moreover, it is to be noted that in Cases 1 and 2 the digestion of proteids is even more complete than usual.

The difference may probably be attributed to the diminution in the curdling ferment rennin, the secretion of which would be impaired in an equal degree to that of the pepsin. The effect of the presence of rennin in the stomach under normal conditions, by its coagulating powers, is to hold back the casein and fat of the milk while the whey is allowed to escape. With a diminution in the amount of rennin the milk would tend to remain in fluid form, with the result that it would escape with undue rapidity from the stomach. The digestion of butter and non-emulsified fats, on the other hand, would be unaffected by such a diminution in the amount of rennin. However this may be, surgeons have apparently recognized this distinction in practice, for they take a pride in increasing the amount and variety of the diet as rapidly as possible after the operation. At Guy's Hospital a pure milk diet is often abandoned on the second or third day after the operation, or even sooner.

Case 6, upon whom gastro-jejunostomy had been performed eight years previously at the Cancer Hospital for simple stricture, was examined with a very similar result; while Case 7, where the digestion of fat was also found to be normal, is interesting, because this result is directly opposed to the findings of Joslin in the two cancerous

cases mentioned above. Case 7 was the only case of malignant disease examined.

Except in the cases in which the diet consisted of milk alone these results agree with findings of Paterson, and are in opposition to the earlier work of Joslin. In view of the teaching of Bayliss and Starling that the stimulus to the pancreatic juice is dependent upon the contact of the acid chyme with the duodenal mucous membrane, these results are surprising. It is true that prosecretin is found in all but the lowest part of the ileum, in amounts gradually diminishing as the distance from the pylorus increases, but only the prosecretin in the first part of the duodenum can be of the first importance, for there only is acid found to bring about its conversion into secretin. The complete digestion and absorption of fat in Cases 1, 4, and 7, where the duodenum is "short-circuited," forces the conclusion that the mechanism of pancreatic secretion, demonstrated by Bayliss and Starling, and dependent on the contact of duodenal mucous membrane and HCl, can if necessary be replaced by other means equally efficacious. These cases tend to support the view that fats have themselves a specific effect upon the pancreas. Indeed, it would be strange if it were otherwise. Fat, as is well known, restrains the secretion of gastric juice, and, if fat had not also a directly stimulating effect upon the pancreatic juice, that statement would also imply that fat acted as a restrainer of pancreatic activity as well. Damaskin¹⁷ has described a very perfect experiment upon a dog which showed undeniably that not only does a definite secretion of pancreatic juice take place after the introduction of oil into the stomach, but that this secretion takes place when every trace of acid contents is absent from the stomach. The investigations of Walther¹⁸ go further and show that fat not only excites a free secretion of pancreatic juice, but also leads to an increase of the pancreatic ferment. Finally, Fleig,¹⁹ by rubbing up the mucous membrane with soap, has obtained a hormone, which he has named "saprocrinin." This reaction is independent of the existence of free acid, and except that the duodenum would take no part, would be uninfluenced by the changes introduced by the operation of gastro-enterostomy. Saprocrinin, more than sufficient, is furnished by the rest of the alimentary canal, in which, as a result of the fatty diet, there will be present a more than usually large quantity of the necessary soaps.

CONCLUSIONS.

1. On a purely milk diet, in non-malignant cases, after gastro-jejunostomy there is a slight but definite diminution in the power of digesting and absorbing fat.
2. On a mixed diet, rich in fat, for the most part in the form of butter, this diminution in power disappears.
3. On a milk diet and on a mixed diet, rich in fat, the results are the same, whether there is obstruction at the pylorus or not. It is therefore improbable that the diminution in power of digesting milk depends upon

whether the duodenum is or is not short-circuited out of the alimentary canal. The cause must be common to both obstructed and non-obstructed cases, and is probably to be found in the regurgitation of alkaline bile and pancreatic juice into the stomach and in the consequent reduction of gastric acidity, together with the inhibition of the rennin.

4. That in cases of recurrent intractable duodenal or gastric ulceration, an operation is indicated which will ensure the complete loss of the acid reaction of the stomach, while the benefit at present conferred in ulceration by gastro-enterostomy is due to the partial loss of that reaction. That such an operation would consist in closing the proximal end of the bowel and implanting the distal directly into the stomach. That where gastro-enterostomy is performed the opening should be near the cardiac end of the stomach, and that the anastomosis should be simple, not Y-shaped, nor with enter-enterostomy.

5. That the complete digestion of fats on a mixed diet is due to the specific stimulating effect on the pancreas of a hormone produced by the fats themselves.

6. A case of malignant pyloric obstruction, after partial gastrectomy and posterior gastro-jejunostomy, showed complete power of dealing with fats.

Finally, I take the opportunity of pleading the laborious nature of the experiments carried out as an excuse for their small number, and of thanking the physicians and surgeons under whose care the patients were at Guy's Hospital, and Mr. Swan at the Cancer Hospital, for their kindness in so readily granting facilities for work which can have no claim to be considered of the highest practical importance.

NOTE.

In an address delivered at the annual meeting of the American Medical Association, at Atlantic City, U.S.A., and published in the *Lancet* for September 21st, 1907, Mr. H. J. Paterson states that he has found little diminution in the power of digesting fat in 4 patients upon whom he had performed gastro-jejunostomy, and who were placed upon a milk diet. Details as to the method adopted and the amount of milk given are wanting. The fat absorption, taking the average of 4 cases, is stated to have been only 1.17 below the normal. The present analyses were completed before the publication of this paper.

REFERENCES.

- ¹ *Work of the Digestive Glands*, Pawlow, English edition, 1902, p. 164.
- ² Croonian Lecture on The Chemical Regulation of the Secretory Process, by W. M. Bayliss and E. H. Starling, *Proc. Roy. Soc.*, vol. lxxiii, 1904.
- ³ E. P. Joslin, *Berlin. klin. Woch.*, xxxiv, 1897.
- ⁴ H. J. Paterson, (Hunterian Lectures, Royal College of Surgeons, 1906)
- ⁵ Cannon and Blake, *Annals of Surgery*, xli, 1905, p. 686
- ⁶ *Annals of Surgery*, vol. xli, October, 1907, p. 519.
- ⁷ Pawlow, loc. cit., p. 163.
- ⁸ Katzenstein, *Deut. med. Woch.*, xxxiii, 1907
- ⁹ Pawlow, loc. cit., p. 145.
- ¹⁰ Denechau, *Les suites médicales éloignées de la gastro-entérostomie*, *Faculté de Médecine de Paris*, 1907.
- ¹¹ *Zeitschrift für Chir.*, lxi, p. 337.
- ¹² Mayo Robson and Moynihan, *Diseases of Stomach*, p. 238.
- ¹³ Loc. cit.
- ¹⁴ *Annals of Surgery*, vol. xli, p. 556.
- ¹⁵ Loc. cit.
- ¹⁶ Loc. cit.
- ¹⁷ Pawlow, loc. cit., p. 121.
- ¹⁸ Pawlow, loc. cit., p. 122.
- ¹⁹ Starling, *Recent Advances in the Physiology of Digestion*, p. 92.

ETHER ANAESTHESIA BY THE OPEN METHOD.

By H. BELLAMY GARDNER, M.R.C.S., L.R.C.P. LOND.,
LATE INSTRUCTOR IN ANAESTHETICS AT CHARING CROSS HOSPITAL.

SINCE the year 1876, when Mr. J. L. Clover invented his portable gas and ether apparatus, it has until recently been the custom in England to administer the vapour of ether as a general anaesthetic by means of a closed india rubber bag. This method produces a very rapid insensibility; the total amount of ether evaporated averages only 6 fl. oz. an hour, and little vapour escapes into the surrounding atmosphere. Its disadvantages are that the partial deprivation of air and rebreathing of expired products from the bag tend to produce:

1. Duskiness of the countenance and oozing of venous blood from the cut surfaces of the operation wound.
2. Considerable secretion of mucus from the membranes of the air passages.
3. Laboured breathing, with strong movement of the abdominal walls.
4. After-sickness in 75 per cent. of the cases.

These factors render the smooth conduct of ether anaesthesia with a bag inhaler more difficult than that of chloroform, with the result that many practitioners

do not attempt its use in private practice, and that chloroform is also frequently resorted to by specialists when complete tranquillity is desired.

The open method of administering ether has been used in America for several years, and the procedure which I now adopt, after a series of careful trials and experiments with various materials, consists in pouring ether upon a wire-framed mask covered with ten layers of white absorbent gauze such as is used for surgical dressings. The mask, which may be of the Schimmelbusch pattern, or constructed according to the author's special design shown in Fig. 1, is allowed to rest lightly upon a freshly-made gauze face-pad consisting of several layers of the same gauze made into a roll 14 in. long and 1 in. thick. The free ends of the roll are crossed and tied

Fig. 1.

together with a short piece of gauze or tape, as shown in Fig. 2, before the mask is applied. An 8-oz. medicine bottle, fitted with the double-tubed stopper illustrated in Fig. 3, is well adapted for the supply of ether, which can be poured from the end of either tube at a rate controlled by the finger held upon the other. To diminish the supply still further the bottle may be held somewhat upon its side, when a small stream of separate drops falls from the tube orifice. A larger stock of ether must also be at hand, as in some cases from 12 to 16 oz.

may be required for a long operation.

Fig. 2.

THE ADMINISTRATION.

The patient being in the supine position is first told to close the eyes, and the gauze face-pad is then loosely laid upon the face forming an oval, which covers the closed eyelids and encircles the mouth and nose. The dry ether mask is then superimposed, and the patient is told to breathe regularly and slowly in and out through the mouth. A few drops of ether are then allowed to fall upon the lower portion of the mask opposite to the mouth; and whilst verbally encouraging the patient to continue this form of breathing, the drops of ether are steadily continued so as to keep the lower half of the gauze moist to the touch but not wet enough to overflow. It is a good plan to keep talking steadily to the patient, instructing him how to breathe and describing the sensations he will experience until he drops off to sleep, so that the sound of the voice may guide and control him while he remains conscious.

The taste of ether when presented in this manner is not objected to by the majority of patients, because it is too weak to be pungent. There is hardly any excitement or struggling except in alcoholic subjects, for whom it should not be used. The only movement which may usually be anticipated is a tendency to slowly raise the head and sit up, which is often witnessed also during the induction with C_2E_2 mixture in adults. This need not be restrained, as it only lasts for a minute or so and if the ether be steadily continued the patient will slowly sink back upon the pillows in the third stage of anaesthesia. The time taken to induce this condition is, on the average, between six and seven minutes. As a rule, when slight stertor

Fig. 3.