A Clinical Lecture

ON

CHOREA OF AGGRAVATED TYPE WITH CERTAIN UNUSUAL PHENOMENA.

BY

R. W. PHILIP, M.A., M.D., F.R.C.P., F.R.S.E.,

PHYSICIAN AND LECTURER ON CLINICAL MEDICINE,
ROYAL INFIRMARY, EDINBURGH.

I HAD intended to address you to-day on another subject. The advent at the commencement of this week of a very aggravated case of chorea, presenting several unusual features, has determined me to direct your attention in that direction.

We are apt to think of chorea as a mild, almost trifling ailment—one of the lesser manifestations of the rheumatic diathesis, tending to occur especially in delicate, neurotic children. As matter of experience, some four-fifths of the cases do occur between the ages of 5 to 15, and the great majority of all cases are mild in character and yield readily to rest and treatment.

Contrariwise, the patient who forms the text of to-day's lecture is a well-developed lad of 18. The attack, which came on suddenly, has been of violent character. Most of the recognized symptoms are present in aggravated form, and there has appeared at least one symptom which I have not seen previously described.

HISTORY.

Let me first of all give you a short statement of the history of the case. W. I., aged 18, a joiner, was admitted to my wards in the Royal Infirmary on January 11th, 1908, suffering from intense restlessness and unceasing movement apparently of all his muscles—in a word from *chorea gravis*.

The patient seems to have been quite well till three weeks ago, when he got a severe chill, and returned home, complaining of cold and of being out of sorts. That night the movements commenced. His father, who slept with him, complained greatly of his restlessness, and the lad seems to have had a fit of emotional excitement. Next day he tried to go about as usual, but his father thought him dull and heavy-looking. He had a quiet time at New Year on account of the holiday. On January 4th he suffered from headache, and pain and stiffness in the knees and left elbow. This made him keep his bed. The doctor was called in, and said the patient was suffering from rheumatism. There was apparently no fever, and the pain disappeared in two or three days. On January 7th, while still in bed, he complained of sore throat. The same day his father noticed twitching in his hand and about the mouth, and thought there was some hesitancy in speech. Next day, January 8th, the throat was much worse, and he had great difficulty in swallowing. Since then the throat gradually improved, but the twitches and restless movements grew steadily worse, and spread all over the body. For two or three days they have been truly violent. He was brought to the Infirmary chiefly because of the difficulty of managing him at home.

Previous Illnesses.—The patient was in the Royal Infirmary two years ago with a sharp attack of chorea, which continued for a month. There is no history of rheumatic fever, nor, up to the date of present illness, had he suffered from sore throat or pains in the joints. Indeed, apart from so-called bilious attacks every now and again, which lasted a day or so, he seems to have been healthy. There is no history or evidence of gonorrhoea.

His general surroundings seem satisfactory. His home is comfortable. He is well fed. His work is pleasant, and his habits are good. The family history shows nothing out of the way. Both his parents and two brothers and one sister are alive and well. There is no history of rheumatism in the family.

PRESENT STATE.

On examination the patient presented the aspect of intense motor restlessness. The clinical picture might truly be described as one of muscular insanity (*folie musculaire*). The motor contortions were universal and

most disorderly. The face showed every variety of grimace and contortion. The forehead was repeatedly wrinkled and alternately knit. The eyebrows were similarly raised and depressed, independently of the movement of the forehead and often independently of each other. The eyelids were closed and opened irregularly, unequally, and often independently of each other. The eyeballs were constantly moved in every possible direction—upwards and downwards, from side to side, obliquely, sometimes in conjugate fashion, sometimes independently. In addition to this, there was present what I do not remember to have seen described anywhere in relation to chorea, namely, the condition of *hippus*, that is to say, the pupil was alternately dilated and contracted in extraordinarily rapid fashion.

The other muscles of facial expression were thrown into irregular contortions, sometimes bilateral, sometimes unilateral, giving rise to a variety of bizarre grimaces, hardly to be reproduced by the healthy subject. Extremely contradictory manifestations were effected, from the broad grin of good nature to the ferocious look of uncontrolled anger. Interest, pleasure, excitement, vexation, distress, resentment, hate, were all rapidly reproduced from moment to moment, more quickly than the lightning sketches of the music-hall artist.

Similarly the jaws were separated and approximated with considerable force. The tongue was projected spontaneously from time to time, and as rapidly withdrawn. When the patient was asked to put out his tongue, there was usually some delay, and then suddenly it was darted out, perhaps rolled from side to side, and drawn back with a jerk. The movement of the tongue against the sharp edge of the teeth had led to considerable scratching and scoring. The lower lip also had been much injured from contact with the teeth during closure of the jaw. The lower jaw was moved from side to side violently, and the head was similarly jerked in all directions, at one time suddenly raised from the bed, and the next moment thrown back again with dangerous force. Backwards, forwards, from side to side, the heedless, delirious movement never ceased.

The upper limbs were likewise active. The fingers, thumb and hand generally, made an endless variety of jerking movements. The fingers were alternately extended to their fullest, abducted, and flexed. The hand was alternately flexed and extended, pronated, and supinated. Wrist, elbow-joint, shoulder-joint participated in the gay dance, with every possible variation and combination.

The lower extremities were also disturbed, with alternate flexion and extension, abduction and adduction, inversion and eversion of foot. Movements of flexion and extension occurred at the knee and hip, with less or more rotation, abduction, and adduction. One foot was made to cross the other, so that considerable bruises resulted on the alternate dorsa of the feet. The two internal malleoli were brought together with considerable force, causing congestion and bruises. The legs were made to cross each other. The trunk muscles were nowise spared. Irregular and unequal contractions of different groups of muscles produced extraordinary movements of the shoulders, back and sides. It was only by constant watching that the patient was saved from being jerked out of bed.

The diaphragm and abdominal and costal muscles were greatly disturbed, and the breathing correspondingly embarrassed. The rhythm of the cardiac sounds and pulse was influenced, but to a less extent than might *a priori* have been anticipated from the extraordinary disturbance in other groups of muscles.

Speech was much deranged, more particularly in respect of articulation, which was jerky, highly irregular, and halting. At times it was quite unintelligible. The patient, when spoken to, seemed to apprehend what was said perfectly, but, while attempting to answer, failed to effect accurate articulation.

The movements were so pronounced and continuous that no increase under observation was certainly determinable. They continued without intermission through the night. For two nights there was complete insomnia. On the second night the patient's mind seemed to wander from time to time, but no maniacal symptoms were manifested. At the end of forty-eight hours from admission the patient showed evidence of extreme exhaustion. The face was flushed and covered with perspiration. The

extremities were similarly clammy. The temperature had risen to 101° F., and the pulse-rate was 120 per minute, with irregularity in time and force.

A fortunate feature up to the present has been the possibility—admittedly with difficulty—of feeding the patient. With continuous watchfulness on the part of nurses, this has been effected to some extent throughout. Sufficient nourishment in these cases is of the first moment. Its achievement in the present instance has doubtless served to ward off the fatal catastrophe which the extreme exhaustion would otherwise have precipitated.

Further examination of the nervous system was rendered difficult on account of the movements. It was impossible to consider sensory disturbance. The deep reflexes were generally present, and apparently natural, perhaps slightly increased. The organic reflexes were undisturbed. I have already referred to the striking phenomenon of hippus which continued for days. There was a certain amount of uncertainty and halting in connexion with micturition.

Other systems showed little out of the way. The circulatory system was examined with care. Apart from rapidity and irregularity of pulse little of importance was determined. The heart's action was rapid and excited, but the sounds seemed pure enough. There was no definite murmur either at apex or base. The urine was natural, apart from a large excess of urates. There was no albumen. Urea amounted to 16 grains per oz.

The condition is, then, of much interest from several points of view. In the first place the age of the patient is unusual. As we have seen, the large bulk of cases of chorea occur between 5 and 15. Of the relatively small group of cases which occur later, from 16 to 30, the great majority are in female subjects. Gowers points out that the disease is rare in lads over 16.

Family history in cases of chorea often reveals a hereditary tendency in one of two directions. There may be, on the one hand, a history of rheumatism, or, on the other hand, a neuropathic leaning. Neither of those is traceable in the present instance. Indeed, apart from slight ailments, the patient's family seem to show an unusually clean bill of health.

As to his personal history, there is no factor of special significance in relation to work, or manner of life or habits. It is to be noted that he had a previous attack of chorea when he was 16. The emotional state into which he seems to have got on being reproved by his father, points to a rather sensitive, neurotic temperament, but nothing more than is common enough. Of definite facts, we must keep in view the history of sudden chill, and the subsequent malaise, which made the patient return home from work. This was immediately associated with some degree of restlessness. After battling more or less successfully against the illness which threatened, he had, in the course of a fortnight, to keep his bed because of pain and stiffness of joints, which was quickly followed by sore throat. The sore throat which supervened three or four days before admission seems to have been of aggravated nature, as judged by the history. It was while the sore throat was conspicuous that the grave nervous manifestations supervened. Such anginous symptoms are not infrequent precedents of chorea. Whether they are always rheumatic in origin or sometimes due to other toxins is a moot point.

DIAGNOSIS.

In the present instance the diagnosis has offered little difficulty. The character of the spasmodic movements is quite distinctive. They have been much more pronounced than they commonly are in chorea, but in other respects essentially those to which the name "chorea" has been applied. Because of their severity, the condition may be properly described as *chorea gravis*, as opposed to *chorea minor*. The term *chorea major* has been retained for another condition, to which I shall refer presently.

Of conditions from which the case had to be separated, acute mania suggested itself when the movements were at their worst. As the patient threw himself about with what simulated suicidal determination, acute mania seemed possible. The absence of conspicuous delirium saved the diagnosis. Although some degree of wandering occurred towards night, the delirium was never of the constant, excited, jabbering type which one gets in acute

mania. Hysteria was likewise excluded. Hysteria of this type is unlikely in a young male subject. Further, there has been an absence of the excitement of so-called higher centres, characteristic of hysteria.

With other diseases associated with irregular movements there was little risk of confusion. Thus from hereditary chorea, or Huntington's chorea, the present case was distinguished by the abruptness of onset, the grave character of the movements, and the absence of hereditary history. From senile chorea it was separated by the patient's youth and the severity of the spasms. From multiple myoclonus, or paramyoclonus multiplex, it was readily separated by the nature of the movements, which differed widely from the lightning-like spasms of myoclonus recurring in the same muscle or group of muscles successively. In myoclonus the trunk and proximal muscles of the limbs are commonly affected in more or less symmetrical fashion. Contrariwise, in the case before us the movements are conspicuously irregular. The same remarks apply to what is known as electrical chorea, which occurs in North Italy. In that condition, the spasmodic movements affect commonly one of the upper limbs first. Thereafter spasms occur in the muscles of the leg of the same side, and later become more generalized. As the name suggests, the contractions are commonly shock-like. From disseminated sclerosis, which occurs frequently enough in young subjects, the present case is easily separated. In disseminated sclerosis there may be little disturbance save when the patient attempts a voluntary movement. Then a spasm supervenes, and the movement is executed in clumsy, jerking fashion—the so-called "intention spasm." It is hardly necessary to mention paralysis agitans, with which the present case could not be confused, first, because of the age of the patient, and secondly, because of the character of the movements. Paralysis agitans is a senile disease, associated with degenerative changes, and the movements, which develop gradually, are of fine, rhythmic character. In athetosis the movements are of quite a different character, being slower, more regular, undulating, and apparently purposive. Lastly, the movements were of too general and grave a type to lead to confusion with habit spasm and other varieties of spasmodic tic.

The case, is then, one of pronounced chorea, or *chorea gravis*. Before passing to discuss the prognosis and treatment, it may be convenient to introduce a word or two regarding the nature and relationships of chorea.

The name "chorea," or "dancing disease," was really stolen from another morbid condition, to which it is more literally applicable—a disease which seems to have been much commoner some centuries ago than now. "*Chorea Sancti Viti*," or St. Vitus's dance, in the early sense of the term, seems to have occurred in epidemic fashion in many parts of Europe. It was truly a choreomania, or dancing mania, especially prevalent in the fourteenth and fifteenth centuries. Patients were sent to various chapels and shrines with the view of calming their dancing ardour, and restraining the exuberance of their movements. The patients affected by the disease appeared to have lost all control over their senses and movements. They formed circles and danced in wild delirium for hours together, until at length they were overcome by exhaustion. In this state they emitted groans as if in the agony of death. By way of treatment they had the abdomen tightly bound with cloths, after which they commonly recovered, and remained sane until the next attack. During the dancing mania the patients appeared insensible to impressions through the senses but had visions. Spirits, whose names they shrieked out, were conjured up. After the attack patients maintained sometimes that they felt themselves immersed in a stream of blood, and that this had led them to leap.

The name "chorea" seems to have been applied first to the extraordinary mania in 1418, when it broke out in Strassburg, and the magistrate discovered that the best line of treatment was to dispatch the affected persons to the shrine of St. Vitus at Zabern, to intercede for the good offices of the saint. A relic of the therapeutic pilgrimage is preserved in the weird procession of so-called "jumping saints" which takes place yearly during Whitsuntide at Echternach in Luxemburg. Such dancing disease has been known variously at different times and places as "St. John's Dance," "St.

Anthony's Dance," and so on. In later times this pronounced form of hysterical or hysterio-epileptic seizure has largely disappeared. Traces of the affection have emerged now and again amongst such sects as the Jumpers and the Shakers. It is well recognized that neurotic excitement of this nature is apt to be contagious by imitation, and in this sense to become epidemic.

It was to such a condition that the name "chorea" was first applied. The theft or transference of the name to the condition which we have been studying to-day was made by Sydenham, who described the more limited spasmodic condition in 1686. It is in his honour that the condition is frequently labelled "Sydenham's chorea." It was also labelled "chorea Anglorum," more especially by Continental authorities, to separate it from the glorified chorea of the Continent to which I have just referred. The distinction between the two conditions is still recognized by many observers in the division into *Chorea major*—that is, the earlier chorea of hysterical or hysterio-epileptic character—and *Chorea minor*—that is, Sydenham's chorea.

ETIOLOGY.

As to the ultimate causation of chorea, there has been a good deal of speculation. It must be freely admitted that even yet the question affords room for theorizing. Its association with rheumatism, whether in the form of acute rheumatism or in some of the minor manifestations of rheumatism, has long been recognized. The frequent concomitance of chorea and heart lesions, more especially endocarditis, is one of the commonplaces of medicine. That there is a close association with these groups of diseases there can be no manner of doubt. It is more difficult to say what the exact bond of relationship is. The older view that chorea depended on the distribution of multiple emboli in the basal ganglia and corpora striata and the production of local areas of softening is not now generally admitted. The fact that chorea is not always associated with endocarditis or valvular murmur is against it. Further, there is an absence of pathological corroboration, even although it has been shown that certain irregular movements of choreiform character have been produced by introducing starch granules into the circulation. The extremely varying character of the movements, in some cases so slight, in other cases so very pronounced and widespread, is against the embolic view, as is the varying duration of symptoms in different cases. In some instances the symptoms disappear within a week or so, almost without treatment. Some cases are quickly affected by such drugs as arsenic. In other instances neither prolonged rest nor any of the usually prescribed drugs seem to have much effect—it may be for months.

What are we, then, to admit as the probable explanation? The varying character of the symptoms and their extreme violence in some cases are much in favour of the view that the disease is due to irritation. In particular, various toxins are admitted as irritants, especially that associated with rheumatism. It is to be kept in view that the disease supervenes, not only in association with rheumatism, but also after other infective conditions, especially scarlet fever, measles, small-pox, diphtheria, enteric fever, cholera, pneumonia, whooping-cough, pyaemia, and gonorrhoea. Similarly, chorea has been associated with several types of sore throat. More than one observer has emphasized the probable entrance of micro-organisms through the delicate mucous membrane of the naso-pharynx. In keeping with this view is the fact that the rheumatic diplococcus has been found in relation to the smaller arterioles in the brain cortex of animals inoculated with the organism and so rendered choreic, and also in a fatal case of chorea in the human subject.

It is more difficult to determine exactly the part played by certain usually admitted causes of chorea, such as fright, shock of various sorts, and the like. Such are common enough precedents of the attack, more especially in young children. It may be, however, that, after all, these are merely predisposing influences, which determine the direction in which the effects of intoxication may show themselves.

As to anatomical changes, it must be conceded that up to the present there is no certain, constant *post mortem* appearance which can be predicated as the anatomical

basis of the disease. That this is to be sought for in the brain rather than elsewhere in the cerebro-spinal axis is suggested by the following considerations—namely, the frequent association of widespread muscular movements with disturbance of more psychic nature; further, the fact that chorea minor is often one-sided (hemiplegic chorea), and the fact that choreiform movements occasionally follow on a cerebral hemiplegia (post-hemiplegic chorea).

PROGNOSIS.

The prognosis in the simpler forms of chorea is usually favourable. Sometimes the condition may last for several months and its treatment be tedious, but a favourable termination almost always results. The tendency to recurrence must be kept in view.

The prognosis is more anxious in the severer types. It is truly grave in the case before us to-day. Several dangers present themselves. The exhaustion caused by the constant, aggravated movement is immense. The change in the appearance of the lad in the course of 36 to 48 hours has been very striking. From a condition of comparative freshness he is reduced to one of extreme enfeeblement. His pulse-rate has been almost doubled, and the temperature has risen considerably. The exhaustion caused by the movements is further emphasized by the almost complete absence of sleep for a day or two.

Besides the exhaustion caused by the movements, much local injury of skin and soft parts has been induced by the frequent movement of one limb against the other, and the tongue and lips against the teeth, which, even with careful supervision and treatment, it has been impossible wholly to prevent.

The very widespread character of the neuro-muscular disturbance in the present instance is evidenced further by the presence of the phenomenon of hippus, which, as I have said, I do not recall having seen in a previous case, or read of in connexion with chorea.

One feature which sometimes renders prognosis more anxious has been fortunately absent so far in our patient—namely, difficulty in swallowing. The patient has been able, with the help of good nurses, to take food fairly well. This has been our sheet-anchor up to the present. He has been supplied abundantly with milk, along with eggs and a little stimulant at stated intervals. It remains to be seen how long this may continue. There is also an absence of important cardiac lesion. Another element which may aggravate chorea in a female patient of his age—namely, pregnancy—does not, happily, come up for consideration.

TREATMENT.

Slighter cases of chorea require comparatively little in the way of treatment beyond release from disturbing influences, whether these be found at home or in school. More or less complete rest in bed and satisfactory hygienic surroundings, along with a simple tonic, such as arsenic, prove readily efficacious. In the course of six to eight weeks the patient is commonly well. Even when the cure is longer delayed, the final result is obtained along simple lines.

It is different in graver cases, such as that we are now considering. Here energetic measures require to be adopted, in order to anticipate and remedy the state of exhaustion with which the patient is threatened, and to prevent his injuring himself in the way to which I have referred.

On this account it is desirable to have the patient on a soft mattress, preferably a water-bed; all hard structures in his neighbourhood must be carefully padded; the head and corners of the bed should be cushioned; the patient's elbows and wrists, knees and ankles should be swathed in cotton wadding. It is not a bad plan to attach a pillow to the inner aspect of one of his legs. Scrupulous care must be taken in relation to feeding. The food should be of a simple, nutritious kind. I am much in favour, in such cases, of the systematic exhibition of egg-flip, say, every two or three hours. Alcohol, perhaps best of all in the form of brandy in egg-flip, say, 3ij, is of much service. Hydrotherapeutic measures may prove serviceable. Where the movements are not too violent a warm bath is of service. In other cases the wet pack may be used with benefit. In all cases great care should be taken in the matter of cleanliness, more particularly in relation to

scratches or sores of mucous membrane and skin which may be produced through the violence of movement.

Of drugs for this graver type of case, the most serviceable seems to me a combination of bromides and chloral. In the present case, bromides were at first used every three hours with insufficient effect. It was only when chloral was added that a sufficient restraining action on the movements was effected. The bromides were given in doses of 30 grains every three hours or thereby. Chloral was then added, first to the extent of 20 grains, which had little effect, and then twelve hours later in 30 grain dose. Fortunately this succeeded in giving the patient four hours' sleep. On the following night a similar dose afforded seven hours' sleep. During the deep sleep the muscular phenomena disappeared almost completely. On awaking from sleep, the patient presented some of the old violence of movement, but he seemed extremely tired. After the second night's sleep, he remained profoundly exhausted, and this has continued. The pulse remains rapid and the temperature elevated (100° F.). This is of grave omen, all the graver that now he has begun to show disinclination for food. When I saw him, a few minutes before coming to lecture, it was hopeless to think of bringing him to the clinical theatre. I must content myself by inviting you to pay a quiet visit to his bedside. The outlook is most dubious. Fatal collapse seems imminently threatened.

POSTSCRIPT.

[For two days the patient continued to get progressively weaker. Feeding became extremely difficult. At the end of forty-eight hours the temperature fell abruptly to 97° F. The pulse got very rapid (about 144) and almost indeterminate. Six hours later the collapse ended fatally. Unfortunately the patient's father resolutely refused a post-mortem examination.

The day after the lecture was delivered there appeared in the *Lancet* (January 18th, 1908) a Note on Certain Pupillary Signs in Chorea by Dr. F. Langmead. After dwelling on the paucity of observations regarding the pupil in chorea, he reports that he has sometimes observed the occurrence of hippus in children more than slightly affected by the sthenic type of chorea. I have pleasure in citing this, because so far as a further glance through the literature reveals, it is, apart from the observations recorded in the lecture, the only reference to the phenomenon as occurring in chorea.]

A Clinical Lecture

ON

CHRONIC BRONCHITIS.

By A. G. AULD, M.D., M.R.C.P.,

SENIOR ASSISTANT PHYSICIAN TO THE PRINCE OF WALES'S GENERAL HOSPITAL, TOTTENHAM, AND LECTURER ON CLINICAL MEDICINE IN THE NORTH-EAST LONDON POST-GRADUATE COLLEGE; PHYSICIAN TO THE ST. MARYLEBONE GENERAL DISPENSARY.

THE treatment of cough, and more particularly of what is termed "winter cough," constitutes a large share of out-patient and of general practice at this season of the year. Now, a condition which is so common year by year, whose features are so familiar, is apt to lose in scientific interest, or to be set aside in the more fascinating pursuit of the newer methods of treating conditions supposed to be more interesting or more important. The words of Hooker, written in the earlier part of last century, apparently remain true:

The science of medicine in all its departments is in a very changeable state. The discoveries which are made from time to time in anatomy, physiology, and pathology, the theories which are put forth, and the new remedies and modes of treatment which are continually proposed, keep up a constant excitement in the profession.

A disease, therefore, which is not invested with the prevailing scientific halo is apt to be neglected. But no disorder, however commonplace, should be divorced from a continuous association with scientific thought. And

when old facts and old views are considered in this light, the commonest conditions acquire interest, and our hitherto defective view of them gathers clearness and definition.

THE DIAGNOSIS OF A BRONCHIAL COUGH.

We are perhaps too frequently apt to conclude that a cough betokens some chest trouble, and unless care be taken, great and sometimes ridiculous mistakes may be made. The diagnosis must always be carefully and systematically carried out. To the practised ear, it is true, a cough will often betray its seat or its causes from its characters, such as tone, intensity, etc. But in all cases the physical examination should be complete. First, the entire mucous tract from the nares to the bronchi should be inspected, as the cough may arise from any point in this path. The external auditory canals should also be examined in view of the possibility of ear cough. In children, teething requires only to be mentioned. Any area of hyperaesthesia, any congestion, hypertrophy, ulceration, granulation or growth occurring at any point in the upper respiratory tract may be the cause of cough. It is unnecessary to enumerate the various lesions which are specially apt to be overlooked. I have known, for instance, a cough which had baffled several physicians quickly disappear after cauterization of hypertrophied adenoid tissue at the root of the tongue. Cough having its seat in the upper respiratory passages is usually of the hard and dry character.

When in the course of this examination the larynx is under survey the trachea should always be inspected, as the information gained is often very valuable. In some patients the trachea, or at any rate its anterior surface, can be seen right to the bifurcation in the usual sitting positions, but in some cases Killian's positions are necessary. By practice one can usually make out the bifurcation or see into the right bronchus, and note is made of the degree of hyperaemia or anaemia, venous congestion, patchy coloration, mucous deposit, and any other irregularities. But these phenomena are so intricate in their relationships that their discussion would demand a separate lecture.

The examination of the chest is next carried out, but it must be remembered that conditions provocative of cough may be present simultaneously in both upper and lower respiratory tracts. In this examination, all possibility of cough arising from diseases of the lungs or pleurae, from pericarditis, aneurysm, tumour or enlarged bronchial glands must be considered. Should the physical signs in the chest be meagre or absent, we must not neglect to consider the possibilities of nervous or reflex cough. You remember the instance given by Graves of a patient whose cough had long baffled the faculty, but which was promptly cured by an old woman who administered something which caused the expulsion of a tapeworm. The stomach and the pregnant uterus are frequent sources of reflex cough of remoter origin. It is necessary, however, to note that the class of nervous coughs may be accompanied sooner or later by congestive conditions in the larynx and trachea due solely to the mechanical irritation. This is commonly found in children, in whom such coughs are frequent. Lastly, there is the purely hysterical cough, which is generally apparent from its loud ringing or barking character, though in some instances it is merely a "teasing hem." I have known iodide of potassium and ipecacuanha given for such a cough in the expectation that it might thereby be "loosened."

If, on the other hand, our examination proves the presence of a cough due to a bronchial catarrh or bronchitis of the chronic variety, we proceed to consider the scientific conceptions on which the treatment is to be based.

THE NATURE OF CHRONIC BRONCHITIS.

It is generally assumed that catarrhal conditions of the bronchial mucous membrane are caused by irritants of various kinds which affect the mucosa as a whole, and this view is doubtless to a certain extent correct. But morbid anatomy, clinical facts and the results of treatment lead us to recognize that a more definite idea of the nature of bronchitis must be entertained if confusion is to be avoided. The indications, however, become clear when we consider the particular elements or structures in the