it may also be exhibiting the same influence on the first division of the parent cell or cells of cancer. On the other hand, it may be said that the nervous influence (supposing it exists) which controls the spread of rodent ulcers may be called into operation only when cancer has started, and has by its presence affected such reflex nerve impulses as those of vaso-constriction or vaso-dilatation, for example. But, if such were the case, one would not expect to see the marked tendency to adhere to one of the areas where two or three overlap, and the rodent ulcer would not elect to spread in one of them in preference to another, and consequently both would be equally affected, and no definite nerve distribution would be visible.

I have another curious rodent ulcer distribution kindly sent to me by Dr. Sequeira. (See Fig. 17.) The disease began in the centre of the cheek. This might be said to be a pure spread in the lymphatic vessels of the part, but it may also be said, first, to be too transverse in direction for that to be the correct explanation; and, secondly, that its area of occupation may be one of a nerve distribution, because practically the same region is occupied by herpes zoster in Dr. Harvey Cushing's case, Fig. 19.

The case of Dr. Harvey Cushing depicted in Figs. 18 and 19 suggests that if the right Gasserian ganglion had been intact the distribution of herpes zoster on both sides would have been symmetrical. It is highly probable that the existence of a peripheral nerve is essential for the occurrence of herpes zoster as a conductor of irritant impulses from the ganglion, yet this is the only case I know which can be said to demonstrate this point. And suppose the spread of cancer be influenced by ganglionic disturbances they must be conveyed or not conveyed by peripheral nerves, and when a peripheral nerve is exposed and irritated by cancer and septic conditions, its power of conducting afferent and efferent impulses must be profoundly changed and the normal spread of cancer modified in consequence. In fact I have already shown grounds for believing that the growth of rodent ulcers are arrested, or at any rate do not spread so rapidly in denervated areas as elsewhere except when irritated by antiseptics, etc.² I have also pointed out inflammatory changes in posterior spinal root ganglion on whose cutaneous distribution cancer began and that I could not observe the same changes in the posterior spinal root ganglia into whose cutaneous distribution the cancer had spread.³ Whether the changes which existed in the ganglion were primary or secondary to the appearance of the growth I will not now discuss, but it is as difficult to imagine how their presence can fail to influence the state of cancer as it is to comprehend how the influence acts or what it precisely is. In the present state of our knowledge one can only speculate upon the possibility of its being an active or irritative influence rather than one of a passive or an inhibiting nature.

REFERENCES.

¹ Head and Sherren, Injury to Peripheral Nerves in Man, *Brain*, cx, p. 252. ² BRITISH MEDICAL JOURNAL, December 12th, 1903. ³ *Ibid.*, July 20th, 1907.

A FORM OF ORAL FILTER TO BE WORN DURING OPERATIONS BY ALL PERSONS INSIDE THE OPERATING ROOM.

BY STAFF SURGEON G. E. DUNCAN, R.N.,
ROYAL NAVAL HOSPITAL, PLYMOUTH.

OWING to Mikulicz of Breslau and Dr. Mendes de Leon, considerable attention has been drawn of late to the question of the surgeon's breath as a source of wound infection.

The fear of infection is not from micro-organisms given forth in the ordinary act of expiration, but in the act of talking.

It has been shown by numerous observers that micro-organisms are not given off in the breath in ordinary quiet expiration; in fact, the expired air is sterile.

The buccal secretions of healthy people contain micro-organisms, and an enumeration is given of the organisms found.

Examinations of healthy buccal saliva prove that the following organisms are commonly present: *Streptococcus brevis* (Lingelsheim), *Streptococcus medius*, *Streptococcus*

conglomeratus, *Streptococcus longus*. All the above cocci were found by Gordon to be present to the extent of from 10 to 100 million per c.cm. of saliva. Staphylococci in less numbers, 1 to 10 million per c.cm. No staphylococcus of the aureus variety was found. Bacilli were conspicuous by their absence. In addition to the above normally-found micro-organisms, every medical man is well aware of the accidental pathogenic cocci and bacilli that are to be found in the throats of apparently healthy individuals, such as the pneumococcus, bacillus of diphtheria, *Micrococcus tetragenus* (Koch and Gaffky). In the way of literature on the micro-organisms in the mouth there are: *The Mycology of the Mouth*, Goadby, London, 1903; *Die Mikroorganismen der Mundhöhle*, Miller, second edition, Leipzig, 1892; *Microbes de la Bouche*, David, Paris, 1890.

In ordinary speaking, loud talking, orating, singing, shouting, sneezing, coughing, etc., organisms are projected from the mouths of healthy and unhealthy people in saliva droplets to various distances and in various directions. Flugge, and also others of his school, made extensive experiments in proof of the above. Papers by Flugge, Laschtschenko, Beninde, Heyman, and Koeniger are to be found in *Zeitschrift für Hygiene*, Bande 30, 34, 38. Modelled on the above experiments, we have those of more recent date by Dr. Gordon. Details of the technique are given at great length in Appendix B, No. 2. Sections 1 to 7 of the Thirty-second Annual Report of the Local Government Board, 1902-3, also an interesting paper by Dr. Rupert Collins, An Experimental Inquiry into the Infection of Operative Wounds from the Skin, the Breath, and the Air. This paper will be found in the BRITISH MEDICAL JOURNAL, July 15th, 1905, p. 121. Gordon found that salivary micro-organisms were capable of being projected in the salivary droplets to a distance of 40 ft. in front and 12 ft. behind the speaker. In all his experiments control tests were most rigorously carried out.

The organisms such as are projected in the salivary droplets of healthy individuals are capable of pyogenesis on a suitable soil. In the medical officer's report, Local Government Board, 1903-4, Dr. Gordon, in the "Differentiation of Streptococci," finds that two of his streptococci isolated from patients suffering from ulcerative endocarditis gave reactions identical with the two found to occur with the greatest frequency in normal saliva. The virulent streptococcus which occurred with greatest frequency in septic processes was fourth in order of frequency amongst salivary streptococci.

If there be truth in the foregoing statements, I would submit that it is in the best interests of the patient who is undergoing operation that those in the operating room should wear some form of mask or mouth filter.

The form of filter is a simple one, consisting of a square of butter muslin measuring 3 ft. 1 in. along each border. This square is folded up like an ordinary handkerchief and is sterilized with the dressings. After the surgeon, his assistants, and others in the operating room have sterilized their hands and put on their operating coats they will apply the muslin filter in this manner. Unfold the muslin from its handkerchief form until one fold remains, that is, the muslin is simply

Fig. 1.

folded once, its area being halved. Grasp the folded edge with both hands and fling the muslin over the head bring the centre of the folded edge down to the centre of the forehead just above the eye-brows, pass the folded edges behind both the ears exposing the free ends down over the shoulders on to the front of the chest. (See Fig. 1.) Now strip (on both sides) the anterior free edge from the posterior free edge up to the level of the lobe of the ear. (This stripping is shown in Fig. 1 on the left shoulder, whilst the fold on the right has not yet been touched.) Now bring the posterior free edge across the face at the level of the bridge of the nose. (See Fig. 2.) Likewise the remaining posterior free edge across the nose tucking the end in, about midway between the outer canthus of the eye and the external auditory canal. Now pull up

the collar of the sterilized coat round the muslin. (See Fig. 3.) With very little practice and some small amount of dexterity it is possible to bring over the nose and

Fig. 2.

mouth a doubled edge making in all four folds of muslin through which the expired air may be filtered.

During the recent visit of the Fleet to Japan I saw this filter in use in the operating theatre of the University of Osaka. The operation was an abdominal section lasting about an hour and a quarter, at no time during the operation did it seem necessary to have to readjust the

Fig. 3.

muslin. This oral filter is now in daily use at the Royal Naval Hospital, Plymouth.

A point not yet mentioned is the improbability that hairs and epithelium with their accompanying micro-organisms should fall into the wound, since the muslin covers the entire scalp, moustache, and beard.

BIBLIOGRAPHY.

John Tyndall, *Floating Matter of the Air*, p. 47, second edition. John Tyndall, *Fragments of Science*, vol. ii, p. 464. Cornet, *Zeit. für Hygiene*, Band 5, Leipzig. Straus, *Ann. de l'Inst. Pasteur*, vol. ii, p. 181. Gunning, *Zentralbl. f. d. med. Wissenschaften*, 1882. Anderson, *Phil. Trans.*, 1887.

APPENDICITIS AT SEA.

WITH REMARKS ON THE SURGICAL EQUIPMENT OF THE
MERCANTILE MARINE.

By A. E. JOHNSON, M.B., F.R.C.S.,

LATE SURGICAL REGISTRAR, MANCHESTER ROYAL INFIRMARY.

In the course of two voyages in vessels of the Royal Mail Steam Packet Company I had the somewhat unique experience of having to perform two operations for appendicitis.

The first case occurred on board the R.M.S. *Sabor*. On July 16th, 1907, the day after leaving Havana for Vera Cruz—a three days' run—I was called to see the captain's servant, aged 18, who complained of acute pain in the right iliac region and other symptoms pointing to acute appendicitis. On the two previous days he had been bathing in Havana harbour. He got up at 6 a.m. on July 16th, feeling ill and with abdominal pain, and worked until 8 a.m., when he was forced to turn in.

Though firmly believing in early operation for appendicitis, I considered that the difficulties of operating at sea warranted

postponing operation as long as possible. During the afternoon of July 17th, however, the pain became more severe, the temperature rose to 101° F., and the pulse-rate to 110, with indications of perforation and impending diffuse peritonitis. Immediate operation offering the only chance of recovery, as we did not expect to arrive at Vera Cruz until the next day, I proceeded to operate with the assistance of Captain Cheret and Mr. Gibbard, the Chief Officer. After making all preparations, I anaesthetized the patient with chloroform, and Mr. Gibbard continued the anaesthetic under my direction. On opening the peritoneum a little clear fluid escaped, the appendix was found to be perforated, quite free from adhesions, and was removed. A small faecal concretion, about the size of a melon seed, was found lying loose in the right iliac fossa. A drain was inserted at one angle, and the rest of the wound closed. Fortunately all went well, and I take this opportunity of expressing my thanks to Captain Cheret for his able assistance at his first case, and to Mr. Gibbard for the admirable coolness he displayed in the administration of the anaesthetic under trying conditions. The patient had much to contend with, owing to heat, mosquitos, and flies; but his sufferings were very much alleviated by the kind way in which Captain Cheret provided him with every possible comfort. For ten days after the operation his temperature was always about 100° F. in the morning and about 101° F. in the evening, although free drainage was provided and there were no local nor general signs to account for the pyrexia. For want of any other explanation I was forced to account for it by the presence of an open wound in the tropics, and the temperature certainly fell to normal when we got out of the intense heat. Convalescence was uninterrupted, the wound being completely healed and the patient walking at the end of a month.

A point to which I would draw attention is the fact that the illness commenced after the comparatively violent exercise of swimming following a period of three weeks of the forced inactivity of life on board ship. Since the importance of inquiring for a history of trauma in cases of appendicitis was pointed out by Kelly and Hurdon¹ and attention drawn to the subject in this country by Mr. F. A. Southam,² a careful investigation of the histories in all cases may lead to the inclusion of many under a traumatic heading, which would otherwise have been omitted. As Kelly points out patients will often attribute tumours, especially of the breast or abdomen, to an injury, but in acute abdominal affections their first thought is towards an indiscretion in diet or exposure to cold, and this may lead to a history of trauma being overlooked unless specially sought for. Small,³ in 1898, collected from literature and individual reports, 15 cases of appendicitis caused by trauma, and stated his belief "that the true cause of the greater percentage of appendicitis in young men is found in the more frequent exposure to accidental injuries and strains, and to the strong contractions of the abdominal muscles necessary in their work." Six years later Kelly found 50 cases "with but little effort." In 40 out of these 50 cases the appendix deviated in some way from the normal, and in 30 concretions were found. The character of injury is of three kinds—blows, falls, and muscular exertion. Twenty of the cases collected by Kelly were due to exertion, including skipping, dancing (2), violent shaking of the body with the head downwards, long hunting, over-exertion, and fatigue, lifting heavy weights (2), long bicycle ride (2), jumping (2), running, playing football, and severe strain while operating. Two of Mr. Southam's four cases might also be put down to strain. One case was that of a boy who strained himself jumping, and the other is that of a boy who was about to dive into a swimming bath and was suddenly pushed in, so that he fell flat upon his abdomen, which forcibly struck the water. In this case the immediate onset of symptoms might be put down to the blow, but to my mind a more likely explanation is the muscular strain in trying to recover himself, or the sudden translation of the viscera tearing some adhesions. I would describe my case as one of traumatic appendicitis, the onset of symptoms so soon after exertion bringing it into line with other recorded cases. There was here a faecal concretion, as in so many of the recorded cases, and one must recognize, in considering trauma as an exciting cause of acute appendicitis, that it is extremely unlikely to lead to symptoms in the case of a normal appendix. There had been no previous attack in this case, but it is common to meet with patients who have recovered from an attack, and who in the quiescent period suffer from slight pain in the right iliac region on over-exertion, and such warnings, indicating some abnormal condition, should lead to operation without delay.