It is a point that should be carefully looked for and noted in future archaeological work, for if it may not be a sign that is of actual medico-legal importance, it may help to throw some light upon the life conditions of an ancient people.

REFERENCE.

¹ Report of the Unrolling of the Mummies of the Kings Siptah, Seti II, Ramesses IV, Ramesses V, and Ramesses VI, *Bulletin de l'Institut Egyptien*, 5e. série, tome 1, 1907.

THE EXAMINATION OF THE BODIES OF 100 MEN EXECUTED IN NUBIA IN ROMAN TIMES.*

By FREDERIC WOOD JONES, M.B.LOND.

It is not often that the examination of a large number of bodies of people killed by various wounds falls to the share of the anatomist. Such cases as reach our museums are, for the most part, isolated instances of death from violence inflicted by accident, and the medico-legal interest attached to the damage of soft parts so greatly outweighs the importance of the bony injuries that the *post-mortem* evidences to be found in the skeleton may easily be overlooked or neglected. Among the vast number of bodies that Dr. Elliot Smith and I have handled during the first three months' work of the Egyptian Government Archaeological Survey of Nubia, we have been fortunate enough to find much material that possesses the advantages of being dated with historical accuracy and of being preserved in the wonderful manner that is common among bodies interred in this part of the world.

No burial site could be better suited to the study of the effects of violence inflicted upon the human frame than the region around the Pillars of Konosso that mark the frontier line of Ancient Egypt. Vast cemeteries of most historic and prehistoric periods lie in this neighbourhood, and the bodies that have been buried in these cemeteries are those of the peoples who have come from the north and from the south, as traders or as soldiers, to this boundary line of commerce and of politics. So far as any records go back, this region of the first cataract of the Nile has been an arena for all those contending peoples

who have lived and fought, who have died and been slain, in the myriad frontier battles waged around Shellal by the Nubians from the south and the more highly-cultured dwellers, or conquerors, from the north.

From isolated instances a great quantity of material has been examined that has been of the utmost interest in the study of *ante-mortem* wounding, but the cases to be described are those of two interments only. The bodies from these burials date from late Roman and Byzantine times, and the trenches in which they were thrown were portions of an extensive Roman fort, built upon the eastern bank of the Nile, and facing the historic island of Philae. In one trench 62 bodies were thrown in one confused mass, and in the other were 40 more (Fig. 1).^{*} Both burial trenches are of the same date, and in both cases all the bodies in the trench were thrown in at one act of

burial. All the skeletons were those of males, and, with the exception of a youth of about 18, all were grown adults in the prime of life. Each corpse was wrapped around with coarse linen, and the cords that bound their legs together, and trussed their arms to their sides, were not even removed before burial. Even the rope with which the great majority had been hanged was found *in situ* around the neck of one of the victims, and in a great many other cases thinner cords encircled the cervical vertebrae in firmly knotted loops. The injury most commonly found, and the one of the greatest interest, was that inflicted upon the skull by the process of hanging. The lesion is a peculiar and very constant one, and although it is not the injury most naturally expected, still its origin is easily seen, and all its features are readily explained. In all cases the lesion was confined to the base of the skull; no *post-*

Fig. 5.—Exit wound of a spear thrust through the loin from the front.

mortem evidences of hanging were to be found in the spinal column, there being no single case of damage inflicted to a cervical vertebra in the whole series of over 100 bodies. The injury to the skull consists of a dragging open of the sutures running antero-posteriorly, and a tearing apart of the whole of the cranial base, so that the lateral halves of the skull tend to be disunited (Fig. 2).

The final effect is an oblique opening of the sutures, so that the one portion of the skull, represented by the occipital and one temporal bone, becomes pulled aside from the other portion, represented by the facial part of the skull and the other temporal bone. The basilar suture in most cases becomes disunited, and so the median axes of the front and back portions of the base of the skull are not continuous.

* Figs. 1, 6, and 7 will be found in the special plate.

* An account of the circumstances under which these bodies were found by Dr. Reissner, to whom I am indebted for the determination of their age, will be found in the first *Bulletin of the Archaeological Survey of Nubia*. This report is published with the permission of the Egyptian Ministry of Finance and Captain H. G. Lyons, F.R.S., Director-General of the Survey Department, under whose direction the survey of the country that will be submerged in the Assouan reservoir is being carried out.

TO ILLUSTRATE DR. WOOD JONES'S PAPER ON THE *POST-MORTEM* STAINING OF BONE.

Fig. 1.—The skull of an Egyptian (Ptolemaic Period) exhibiting a fracture surrounded by blood stains.

Fig. 2.—The fractured mandible of a prehistoric Egyptian showing blood stains.

TO ILLUSTRATE DR. WOOD JONES'S PAPER ON THE EXAMINATION OF THE BODIES OF 100 MEN EXECUTED IN NUBIA.

Fig. 7.—Skull showing the passage of a sharp instrument that severed the head from the body.

Fig. 1.—View of the bodies undisturbed in the trench.

Fig. 6.—Sword cut of the parietal bones.

The lesion commences behind, at the meeting of the lambdoid suture with the parieto-mastoid and occipito-mastoid sutures. The occipito-mastoid suture is torn widely open, and the whole basal portion of the occipital bone is separated unilaterally from its connexions, the jugular and basilar processes being torn across their synchondroses. The line of disruption now passes forwards to the other side of the skull, the great wing of the sphenoid being separated from the temporal bone, and the squamous suture being opened in a varying extent of its course.

In the great majority of the cases the line of opening of the sutures passed from behind forwards from right to left; but in some instances the process was exactly reversed, and in these the left occipito-mastoid suture and the right spheno-temporal suture were opened (Fig. 3).

The factor that caused the change from the right to the left-sided opening of the hinder sutures was doubtless the varying position of the knot of the hangman's rope; and whilst in the great majority of cases it was placed upon the right side, in some few instances it was reversed, and placed upon the left.

The mechanical causation of this lesion is easy of explanation. It was doubtless the sudden jerk, accompanied by an upward pull, imparted to one portion of the head, and a downward pull to the other, that caused the sutures to be strained apart. The upward jerk of the head acts upon one side only—the side upon which the knot is placed; and the pull acting from the temporal bone—for the knot doubtless caught on the mastoid process and ramus of the jaw—caused that side of the head to be violently strained upwards and downwards. The counter-acting pull of the body was transmitted via the occipito-atlantal joint, and so the occipital bone, and the opposite side of the skull, were pulled violently downwards.

The temporal bone of the knot side naturally carried the facial part of the skull with it, for the stress was transmitted forwards through the jaw; whilst the body weight acted on the basi-occipital, and so carried the non-knot temporal in the opposite direction. The bases of all these skulls that give evidence of hanging also show a widespread reddish staining of the bone; the stain is doubtless due to the blood exuded in the neighbourhood at the moment of death. This *post-mortem* stain, so strangely permanent, which is imparted to the bones by the *ante-mortem* shedding of blood, we have come to regard as a sign of special importance, and it has been more fully discussed in a special note.

The exact methods employed in these ancient executions by hanging are, of course, not readily determined, and it is not impossible that in some details they may depart from those used in judicial hangings practised to-day. But if there may be some doubt as to the manner of hanging, there can be none as to its effects upon the skeleton; and, unless these cases are for some reason exceptional, then the best *post-mortem* evidences of death by hanging are those to be found in this peculiar lesion of the base of the skull. It is possible that in the *post-mortem* examination of persons killed by judicial hanging this injury to the cranial base may have been overlooked, for if a sufficient cause of death could be found in other parts of the body this lesion might easily escape detection.

Besides these people who had met their death by hanging, others were found in these executioner's trenches, and they had died from a variety of injuries. Some bodies showed extensive recent fractures of ribs. One (Fig. 4) had a typical *ante-mortem* fracture of the right parietal bone, with the fragment still inside the cranial cavity, and, as in the other cases, there was abundant evidence

of the blood staining of bone. Some had met their death from blows of sharp instruments; one body showed in the right loin the skin wound of the exit aperture of a spear thrust from the front (Fig. 5); one had the manubrium sterni divided obliquely by a sharp blade; and one had a large penetrating wound of the skull, in which the sword had actually raised a portion of the right parietal bone as a partial flap (Fig. 6).

One skull only gave evidences of decapitation, and it showed in a remarkable manner the clean-cut track of the blade that had divided the left mastoid and occipital condyle in the act of severing the head from the trunk (Fig. 7).

REMARKS UPON EXCISION OF THE BODY OF THE SCAPULA.

WITH AN ILLUSTRATIVE CASE.

By R. P. ROWLANDS, M.S.LOND., F.R.C.S.ENG.,
ASSISTANT SURGEON AND SURGEON IN CHARGE OF THE ORTHOPAEDIC
DEPARTMENT, GUY'S HOSPITAL, LONDON.

[WITH SPECIAL PLATE]

SOME important points have to be considered before deciding upon the most suitable operation, if any, for every individual case of growth of the scapula. The choice has to be made between three chief operations, namely: (1) Partial excision, (2) excision of the whole scapula, and (3) interscapulo-thoracic amputation.

It is necessary to estimate the relative advantages and disadvantages of these procedures for each case as it presents itself. Only the first two operations had to be compared in the case reported, for the growth was neither extensive nor malignant enough to suggest such a mutilation as amputation of the forequarter.

The most important points to discuss are: (a) The immediate risks, (b) the chances of recurrence, (c) the loss of function, and (d) the amount of deformity associated with each operation.

Statistics,* although they have their value, must not be taken too seriously, for any considerable collection

of cases of such uncommon operations includes a very large proportion of those which were performed before the days of aseptic surgery, so that the chief value of the records of the past is that they clearly show absolute surgical cleanliness to be of vital importance for these severe and extensive dissections. In most other respects they are misleading, because they deal with too few cases, many of which are so dissimilar as to be hardly comparable. Percentages based on such a collection of published cases are apt to give erroneous impressions.

A. The Comparative Risks of the Two Operations.

In the past the operative mortality of complete excision has been a little less than that of partial excision, and considerably less than that of nearly complete excision of the scapula. The main causes of death have been sepsis and haemorrhage, both of which have been more common after incomplete removal of the bone. There is no reason why they should be so in the future, for haemorrhage can be better controlled, especially by systematic ligation of the scapular vessels at the beginning of the operation. It has been asserted that the vessels retract under the portions of bone that are left in partial resections, but surely this must be due to incomplete exposure, and neglect of the obvious precaution of tying the large vessels at the earliest possible opportunity.

Oozing from the cut sections of bone in partial excision

DESCRIPTION OF FIGURES IN SPECIAL PLATE.

Fig. 1.—Back view after nearly complete excision of the scapula.

Fig. 2.—Back view after complete excision of the scapula. The deformity is much greater after complete than after nearly complete excision.

Fig. 5.—Excision of the body of the scapula; the glenoid cavity and the processes have been preserved. The coracoid process is held in its normal relation to the clavicle by the conoid and trapezoid ligaments, therefore there is hardly any flattening of the shoulder. There is a considerable interval between the humerus and the ribs.

Fig. 6.—Excision of the whole scapula. The head of the humerus lies close to the ribs and the outer end of the clavicle projects outwards.

Fig. 7.—Excision of the body of the scapula; the glenoid cavity and the processes have been preserved. The glenoid and the coracoid, in one piece, are lying internal to the head of the humerus, suspended by the conoid and trapezoid ligaments from the outer third of the clavicle. The arm has been passively abducted.

Fig. 8.—Excision of the whole scapula. The head of the humerus lies close to the ribs, and the clavicle is also displaced. The arm has been passively abducted.

* There are less than 100 partial resections recorded, and only about 75 complete excisions (Dr. Buchanan, *Phil. Med. Jour.*, 1900, vol. vi, p. 73).