followed by subsequent surgeons. Where the prolapse only affects the mucous membrane, then its excision, as practised by Patron, Malherbe and Pousson, probably offers the best method of dealing with it. It also has been suggested that the prolapse should be excised, even when it involves all the coats of the bladder, save when the ureters are involved in the displacement.

1. BUMKE (*Jahrb. f. Kinderheilk.*, Bd. 29, 1889).—Female, aged 6 months. Partial prolapse of bladder. Reduced and kept up by pad and bandage. Cured.

2. CROSSE (*Trans. Provincial Med. and Surg. Assoc.*, 1846).—Female, aged 3 years. Partial prolapse showing the openings of the ureters. Cured by pad and bandage, but slight incontinence of urine persisted.

3. FOUBERT (*Schmidt's Jahrbuch*, Bd. 100, S. 225).—Male of advanced age. First stage of inversion seen at autopsy.

4. DE HAEN (Reported in *extenso* by Hoin, *Essai sur différentes hernies*, p. 342. Paris. 1768).—Female. Inversion and prolapse of the entire bladder. Vagina and rectum also prolapsed. Death from gangrene of bladder.

5. LEEDHAM-GREEN.—Female, aged 18 months. Partial prolapse of bladder showing openings of ureters. Cured by injection of hard paraffin.

6. LEVRET (*Journ. de Méd., Chir., Pharm.*, etc., 1773, xl, 269).—Female. First stage of inversion seen at autopsy.

7. LOWE (*Lancet*, 1862, vol. i, p. 250).—Female, aged 2½ years. Congenital prolapse of bladder. Cured by cauterization of the urethra, but slight incontinence of urine persisted.

8. MALAGODI (*Racoglitare Medico di F.no*, 1855).—Female. Complete inversion of bladder. Rectum also prolapsed. Death from gangrene of the bladder.

9. MALHERBE (*Ann. des Mal. Organes Génito-Urinaires*, 1884, ii, 734).—Female. Partial prolapse of mucous membrane of bladder. Cured by ligature and excision.

10. MECKEL (*Traité d'Anatomie pathologique du Lohstein*. Paris. 1829).—Female. Complete inversion of bladder. Only case unaccompanied by displacement of the other pelvic organs.

11. MURPHY (*London Medical Gazette*, vol. xi, 1833).—Female, aged 4 years. Partial prolapse, showing orifices of the ureters. No recurrence after replacement.

12. NIEHAUS (*Correspondenzbl. f. Schweiz. Aerzte*, No. 5).

13. NOEL (*Mém. de l'Acad. R. y. de Chir.* 1769. T. II, p. i).—Female. Congenital prolapse of a translucent cyst, the size of a hen's egg, filled with urine. (Probably a prolapse of a cystic ureter.—C. L. G.)

14. OLIVER (*Edin. Med. Journ.*, Bd. 22, p. 769).—Female, aged 16 months. Some congenital defect of uterus and vagina. Prolapse reduced and kept up by pad and bandage. Cured.

15. PATRON (*Archives Gén. de Méd.*, 1857).—Female. Partial prolapse of mucous membrane of bladder. (Cystic lower end of ureter (?).—C. L. G.) Ligatured and excised. Cured.

16. PEROY (in Boyer, *Traité des Maladies Chirurgicales*, 1824. T. xi, p. 86).—Female, aged 52 years. Inversion and prolapse of portion of bladder. Cured by the tying in of an elastic bougie.

17. PILGRAM (*Inaug. Dissertat.* Bonn. 1892).—Female, aged 10 months. Cared by use of vaginal tampon.

18. POUSSON (*Ann. des Mal. des Org. Génit.-Urin.* 1894. Vol. xii).—Female, aged 30 years. Partial prolapse of mucous membrane of bladder. Cured by excision.

19. RUTTI (*Schmidt's Jahrbuch*, Bd. 100, p. 225).—Male. Autopsy showed first stage of inversion.

20. SOLINGEN (*Observ. de mulier. et infant. morb. chir.* P. 741).—Female. Prolapse of vesical mucous membrane. Cared by use of bougie, as in case of Percy.

21. THOMSON (*Lancet*, 1875. P. 46).—Female, aged 40 years. Did not recur after digital replacement.

22. TIZONI (*Gazzetta medica di Lombardia*, 1855).

" SOLITARY KIDNEY," WITH THE NOTES OF A RARE CASE.

By BERNARD J. WARD, F.R.C.S. Eng.,

LATE SENIOR HOUSE-SURGEON TO ST. PETER'S HOSPITAL FOR STONE,
AND PATHOLOGIST TO QUEEN'S HOSPITAL, BIRMINGHAM.

My object in reporting this example of "solitary kidney" is to draw renewed attention to a class of case which now and again crops up, when least expected, and often ends disastrously.

Now that kidney operations are so often undertaken, these cases have been found to occur fairly frequently in the practice of any large hospital; a surgeon performs some operation (it may be only a slight one) on one kidney, which necessarily interferes with its function for the time being, and which under ordinary circumstances (the other kidney being healthy) would result favourably; but if the other kidney happens to be absent, or so diseased as to be incapable of dealing with the waste products of the body, uraemia ensues, and another of those appalling accidents which are the bugbear of every urinary surgeon has happened.

The term "solitary kidney" is widely and somewhat loosely used to denote several distinct and separate classes of case, but it has enjoyed such an extended vogue that it is useful to retain it, whilst recognizing that it includes the three following classes:

1. *Congenital absence of development of one kidney*, its vessels and the corresponding ureter. Occasionally this is associated with non-development of the genital organ on the same side of the body. In a few cases the ureter is only partially absent, the upper end being represented by a fibrous cord, whilst the lower end may even be pervious for a few centimetres and have an opening into the bladder. The kidney on the opposite side is hypertrophied, and may occupy its normal position in the loin or be displaced downwards into the pelvis or in front of the sacrum. Its ureter most frequently opens into the bladder in the normal position, but may open anywhere on the posterior wall or base. This is the rarest form of "solitary kidney," and the case which I am about to describe belongs to this class.

2. *Fusion of the two kidneys into one mass*, the simplest and most frequent form of which is the well-known "horseshoe kidney," in which two of the poles (usually the lower) are united by a bridge of kidney substance across the middle line; they vary from this to an irregular-shaped mass of kidney tissue. In this form there are never less than two ureters, which usually enter the bladder in the ordinary position; three or four ureters have been frequently met with in this type. They may unite before opening into the bladder or open separately, in which instance they usually open close together in the normal positions. I have throughout the above descriptions laid particular stress on the number and position of entrance of the ureters into the bladder, because any abnormality in either the number or position of these openings noticed during a cystoscopic examination should suggest some malformation of the kidneys, which might not otherwise have been suspected.

3. *Those cases in which one kidney is either rudimentary, degenerated, or so diseased as to be incapable of itself of carrying on the excretory needs of the body*. This class is probably that most frequently met with by surgeons, particularly in tuberculosis and nephrolithiasis, which diseases so frequently affect both kidneys at the same time, whilst the symptoms very often point to disease of one organ only, and it is not until suppression of urine follows some operation performed to relieve these one-sided symptoms that disease of the other organ is suspected.

The case which I am about to describe belongs to the first of these classes.

During the routine examination of the body of a man aged 30 years, who died from cardiac failure following chronic mitral disease, I was surprised, on feeling for the right kidney, to find that the usual site of that organ in the loin was quite empty; and, further, that on close inspection no trace of any kidney on this side was to be found, either in the pelvis or elsewhere. On dissection it was also apparent that neither the corresponding renal vessels or ureter had been developed; there was not even a pad of fat or other tissue to represent the missing organ.

THE Pan-American Medical Congress will hold its fifth meeting at Guatemala from August 6th to 10th, 1908. The work of the Congress will be distributed among twenty-one sections. English, Spanish, French, and Portuguese are the official languages of the Congress. The general secretary is Dr. José Azurdia, of the Guatemala School of Medicine.

The adrenal gland was present and in its usual situation, but was supplied with blood by branches of the diaphragmatic arteries only, the usual aortic branches being absent.

The left kidney was twice the normal size, and weighed 12 oz.; it was of the usual bean shape with a smooth surface, and it lay in the normal position. A section showed chronic congestion, the result of a gradually failing heart; its vessels were at least twice the normal size, but the pelvis and ureter were normal both in size and shape.

The bladder was healthy, but the trigone gave a very curious appearance, the result of the absence of the right ureteric opening, the plica ureterica, and the interureteric bar. The left ureter opened in the usual situation, and its orifice was healthy; but the band of muscle which passes down from the ureter, and which should interlace with its fellow of the opposite side before passing downwards (forming the prominences of mucous membrane known as the plica ureterica and the interureteric bar), in this case ran straight down to the crista gallinaginis of the prostatic urethra, which it appeared to join. The result of this was to give the base of the bladder a curious lop-sided appearance which must be quite characteristic and could not be mistaken if seen by the cystoscope.

Fig. 1. shows the trigone and prostatic urethra of a normal bladder, the ureteric bands of muscle interlace to form the interureteric bar, and then unite to pass down to the crista gallinaginis of the prostatic urethra. A, Interureteric bar. B, Ureteric opening. C, Plica ureterica. D, Crista gallinaginis.

Fig. 2 shows the trigone and prostatic urethra as it appeared in the case of "solitary kidney" just reported. Only the left ureteric opening and band of muscle are present, and the latter passes straight down to join the crista gallinaginis, a condition which could be at once recognized by the cystoscope. E, Ureteric opening. F, Plica ureterica.

Having described the various forms of "solitary kidney" and given an example of one of them, the question arises, given one of these cases, how are we to diagnose it?

Beyond the usual clinical ones there are several methods at our disposal, one or other of which we should be ready to make use of when the occasion arises; and it is well to bear in mind that, although we may suspect complications in a given case, yet the accident usually occurs when the surgeon feels most sure of himself, therefore some attempt to estimate the working value of the *other* kidney should always be made before any kidney operation is undertaken.

The methods at our command are as follows:

1. *Cystoscopy*.—Examination of the bladder with the cystoscope may be done practically without risk by any one accustomed to its use. By means of it we can see the number, shape, and general appearance of the ureteric orifices and the state of the urine as it is expelled from each, and by the use of the ureteric catheter we can collect samples of urine directly from each kidney and apply the usual tests to them, which will often tell us all that we require to know; but, if necessary, we can also make use of special methods for estimating the comparative function of the two kidneys, such as their activity in excreting methylene blue, or in forming sugar after the injection of phloridzin. Within the last few years we have learnt that a diseased kidney will very frequently produce changes in the corresponding ureteric orifice, which in certain cases are quite characteristic, and in others allow us to draw valuable deductions as to the state of the kidney. These changes can only be observed by a skilled use of the cystoscope.

2. *Segregation*, or separation of the bladder into two halves by a special instrument, which enables us to collect the urine from the two kidneys separately. This process is valuable in those instances (happily rare when the observer is a skilled cystoscopist) in which, for some reason or other, it is not possible to use the cystoscope. It can also be combined with the methylene blue or phloridzin-glycosuria methods of comparing kidney function, but by its use alone the valuable positive evidence of sight gained by the cystoscope is lost, besides which it is not nearly so reliable as ureteral catheterization.

3. *Inspection of the two Kidneys*.—This requires a second operation to be undertaken and increases the severity twofold, besides which it is often quite impossible, without dissecting an organ, to say whether or not it is seriously diseased; therefore, when either of the above non-operative methods is available, the extra manipulations, with their added risk, are not nowadays often made.

If this imperfect summary of an ever-increasing branch of urinary surgery should lead to a more frequent use of the cystoscope and its adjuncts before any operation is undertaken on the kidneys, I feel sure that surgeons will be convinced of its utility and feel amply repaid for the little extra trouble it entails.

REMARKS ON TRAUMATIC RUPTURE OF THE KIDNEY.*

By CORNELIUS A. GRIFFITHS, F.R.C.S.,
ASSISTANT SURGEON, CARDIFF INFIRMARY.

THE remarks I have to make are based upon two cases of subparietal rupture of the kidney which came under my care during 1907.

CASE I.

Arthur T., aged 22, a coal trimmer, was admitted to the Cardiff Infirmary on January 17th, 1907, at 4.45 p.m., with the following history: He was running along the iron deck of a ship, slipped and fell sideways to the left, and in falling struck his left side just below the ribs against a large piece of coal on the deck, with a sharp edge upwards. He got to his feet at once unassisted, and, though he felt very faint, managed to climb a ladder to the upper deck, down another ladder to the deck side, and walked to a shed close by and sat down; he felt no particular pain in his side, but was "generally bad." About five minutes after the accident he tried to pass water, and was alarmed to find he was passing what appeared to be pure blood; after this he felt much worse, and nearly fainted, but he did not lose consciousness, nor did he vomit. He was assisted on to an engine, and afterwards on to an ambulance, and brought to the Infirmary.

Condition on Admission.

The patient was suffering from shock; pulse 84, temperature 98.6°; not greatly blanched. He complained of pain all over the abdomen, but more especially over the left lumbar region. There was marked tenderness under the left hypochondrium and all down that flank, preventing deep palpation; the rest of the abdomen was only slightly tender. Percussion did not elicit any unusual dullness. There were a few slight abrasions upon the left flank but no fractured ribs. He passed 1 oz. of bright red bloody urine, and complained of a burning sensation on passing it. No clots were passed, and there was no pain in the bladder. He was put to bed, kept warm, and a hypodermic injection of ergotin citrate ($\frac{1}{10}$ grain) given. A little later the temperature was 98.5° F., the pulse 110; he was weaker, and had passed 1 oz. of nearly pure bright red blood. At 6.30 I saw the patient and decided to explore the kidney for rupture.

Operation.

The usual oblique lumbar incision was made over the left kidney. The perirenal fat was infiltrated with a quantity of recently-effused blood, which was cleared away; the kidney, when brought into view, was seen to be torn on its posterior aspect, about the middle, the tear extending from the convex border to the hilum, deeply into the kidney substance, and communicating by a small opening with the pelvis. The torn surface of the kidney was still bleeding freely. Having ascertained that the kidney contained no stone, the tear was sewn up with iodized catgut, and bleeding stopped. The wound was closed in layers with catgut, provision having been made for drainage by gauze wick reaching the kidney.

Progress.

The next day, January 18th, drain removed, urine blood-stained. January 20th, only a trace of blood in the urine. January 25th, urine clear, a trace of albumen. January 27th, urine clear, wound healed "primarily." January 29th, allowed up. February 4th, discharged well.

* A paper read at a meeting of the Cardiff Medical Society, February 11th, 1908.