The way in which the infection was carried in this case is not clear.

The case is also worthy of record for the unusually large tumour below the left lower eyelid. From its appearance as well as from the feel of it it was diagnosed to be a sebaceous cyst. This was proved to be correct on its removal by operation.

I am greatly indebted to my House-Surgeon, Mr. E. F. Finch, M.B., for his careful observation of the case, and for the manner in which he carried out the treatment.

THE VALUE OF BILATERAL URETEROSTOMY (SHORT-CIRCUITING OF THE URINE) IN ADVANCED CANCER OF THE BLADDER OR PROSTATE

AND AS AN ADJUNCT TO THE REMOVAL OF THE BLADDER IN
UNCONTROLLABLE LOCALIZED DISEASE.

By E. HURRY FENWICK, F.R.C.S.,

SURGEON TO AND A LECTURER ON CLINICAL SURGERY AT THE
LONDON HOSPITAL.

I HAVE always declined to operate upon certain forms of infiltrating bladder cancer, except of course to establish drainage, being convinced that the sufferings of the patients

Fig. 1.—Vertical half of the bladder removed from Dr. Clitherow's patient. (Royal College of Surgeons Museum.)

were vastly increased by meddling with growths that could not thoroughly be removed. I considered, moreover, as one or other of the ureteric orifices were nearly always involved in carcinoma attacking the bladder, that it was useless to transplant them into the bowel and remove the bladder. I cannot say just how many I have rejected upon these counts in the course of my work, but I have operated 270 times in selected cases of growths of the bladder with a fair measure of success and a low mortality. When I mention that the largest mass I have removed filled a quart measure, it will be realized that I have turned aside from nothing which I deemed honestly surgical. I have always, however, been keenly alive to the final misery and the agony of those patients I rejected as being beyond the limits of true surgery.

A suggestion by F. S. Watson of Boston¹ has now been before the profession two years. When it first appeared it

arrested my attention profoundly. I have tested the principle in my own way, and I now desire very cordially to bear testimony to its value, and to urge its serious consideration upon my *confrères*.

F. S. Watson, after criticizing the high mortality and frequent recurrence in the operative treatment of vesical tumours, advocates a preliminary nephrostomy on both sides, in order to divert the urine, and in certain favourable cases he suggests as a further measure a total removal of the bladder, with its contained tumours. Whether he has done cystectomy on these lines or not I cannot say, but those who are compelled to watch the torture involved in advancing cancer of the bladder or prostate, and who often find themselves powerless to relieve the sufferings of the incessant attempts at agonizing urination, will realize the relief which merely short-circuiting the supply of urine can afford.

But bilateral nephrostomy is, in my opinion, far too severe an operation as a mere measure of relief in advanced cancer. Instead, therefore, of incising both kidneys, I usually make a small transverse incision over both iliac crests, and bring out the ureters on to the loin. This simple expedient—for it is simple in expert hands—is admirable in affording instant ease from suffering, and that without shock. The rationale is, of course, the same as that of colotomy in inoperative cancer of the rectum.

Fig. 2.—Separation of bladder from the peritoneum.

CYSTECTOMY.

But there is a further step: short-circuiting the ureters is the secret of a successful cystectomy. It does away with one of the greatest elements of failure in the entire removal of the bladder—and in many instances early removal of the bladder, when soft carcinoma attacks that organ, is the only real chance a patient has of cure. It will prove of infinite service in recurrent cases of soft luxuriant villous vesical papilloma with its power of bleeding and its capacity of obstructing urination without killing the patient.

With a preliminary ureterostomy I can now remove a bladder without the certainty of a fatal issue, if ordinary *vis* is present.

Three examples will suffice.

A man, aged 48, who had been twice operated upon by me for recurrent villous-covered growth of the bladder, was sent back to me last year by his medical man, Dr. Clitherow of Peckham Rye, with a note to say that his sufferings were unbearable and

unrelievable. They were described as follows: "Every act of micturition, and it occurs every ten minutes, resembles the labours of childbirth." The bladder could be felt projecting above the pubes, for it was so full of soft growth that it, with its containing urine, felt like a child's football. I brought both ureters on to the loin at one sitting.* There was no shock and his suffering was greatly reduced. A week later I removed his bladder (Fig. 1) by a method I adopt in non-malignant cases, which I may roughly designate as "shelling." He has been free from pain since, and it is now six months since the operation.

The technique of excision of the bladder is simple. Its secret consists in the operator using a strong electric head lamp, appropriate retractors, and in shelling out the *distended* bladder without opening it except at the vesical orifice by keeping strictly to the outer muscular layer of the organ. A vertical skin and a transverse muscle incision is made suprapubically. The bladder being brought into view, its posterior aspect is first separated by scissor-clipping from the peritoneal layer as far down as the posterior border of the prostate (Fig. 2). The separation proceeds *between the prostate and the bladder base* until the trigone is reached. If there is any suspicion of cancer, the vesicles and prostate must be removed, and in this case the shielding hand protects them from the rectum while the scissoring continues. The anterior surface is now separated from the

Fig. 3.—Separation of bladder from the pubes.

pubes as low as the prostate (Fig. 3). With a little tactile dexterity the scissors (a broad-bladed pair) can be made to cleanly cut through the vesical orifice—if the prostate may be left—and the trigone. The freed bladder is now lifted on to one side to allow of the ureters being traced; these are detached a little way up and divided. If the prostate has to be ablated, the scissor aims lower and cuts through the membranous urethra to join the posterior incision.

CYSTECTOMY, PROSTATECTOMY, AND SEMINAL VESICULECTOMY.

* Another patient, a gentleman aged 56, sent me by Dr. Reynolds of Stamford Hill in April, with a soft carcinomatous growth of the right lateral wall of the bladder, a growth which just filled a 6-oz. measure glass, was in great distress with frequent and painful urination.

*It is a detail which does not invalidate the principle, but this patient had three ureters and I overlooked one of them.

Both ureters were first short-circuited, and a week later the bladder, prostate and vesicles, and lower part of the ureters were skinned out entire and without opening the peritoneum. He is up and healing well.

UNILATERAL URETEROSTOMY AND HEMICYSTECTOMY.

We can extend the principle in women, but not, I think, in men, to unilateral ureterostomy and subsequent hemicystectomy, when the vesical cancer has distinctly invaded only one ureter and the other is clear.

I was asked by Dr. Lewis Smith to examine a female patient sent to him by Dr. Collie of Balham. She was 40 years of age, and had suffered from continuous haematuria for weeks. The right kidney was enlarged and tender.

The diagnosis was carcinoma of the bladder overlapping and blocking the right ureteric orifice, causing back pressure on the right ureter and right kidney pelvis, complicated by ascending pyelitis. I brought the right ureter on to the loin for drainage. It was swollen, dilated, and puriform urine oozed from it. At the same sitting I split the bladder in two, vertically, and removed the right half of the organ with 3 in. of the lower right ureter. I left the vesical orifice of the bladder and repaired in the same way as that adopted in removal of half the tongue, attaching the upper cut convexity to cut concavity of the trigone. She is bidding fair to do well. The growth covered the right ureteric orifice; it was hard and carcinomatous and equalled half a Tangerine orange in size.

I gratefully acknowledge that the principle of bilateral kidney drainage is due to F. S. Watson of Boston. I think as a principle, even as an operation, it will rank high in palliative urinary surgery. I do not think, however, it is realized how simple, expeditious, and free from shock ureterostomy is, and as an operation of relief it will be preferred to nephrostomy.

I should like to advance the following propositions:

1. What can be done for uncontrollable and luxuriant benign villous growth of the bladder, in which the patient suffers from exhausting haemorrhage or uncontrollable pain, or is commencing to suffer from ascending septic changes along the ureters? In such a case I agree that bilateral nephrostomy, or, better still, ureterostomy, should be performed, and a fortnight later the bladder should be removed—if the patient's physical condition and future seem to permit of this mutilation.

2. If the bladder is affected by interstitial cancer, which has so invaded the surrounding areas that no chance remains of removing the organ, I submit that bilateral ureterostomy is wiser, for this will relieve the patient of the agony of urination.

3. If malignant disease is detected early, as it can be by the cystoscope, it is wiser to perform double ureterostomy; and if the physical and renal condition of the patient permit, let the bladder, with or without the prostate and seminal vesicles, be entirely ablated.

REFERENCE.

¹ *Annals of Surgery*, December, 1905.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

SURGICAL EMPHYSEMA.

My partner, Mr. Blaber, and I were recently called to a case of surgical emphysema that was curious, first in the way it originated and secondly in the absence of symptoms. The patient, at about 6 p.m., stepping off a motor omnibus before it had stopped, fell upon his face, giving his neck a severe jerk and causing his nose to bleed. He walked home, and an hour and a half later, the epistaxis having ceased, partook of dinner, consisting of cutlets, vegetables, etc., and a glass of soda water. His wife then noticed that his face and neck were beginning to swell, and he also felt that his collar was tight. An examination showed extensive surgical emphysema, surrounding his neck and reaching from his eyes to the fourth rib on each side, to be present. There were two or three superficial abrasions upon the nose. No facial bones or ribs were broken, and he complained only of a slight stiffness of the neck. At the back of his palate was a clot of blood, and the removal of this disclosed a black object sticking into his nasopharynx just above the pillars of the fauces of the right side. This was detached with some difficulty, and proved to be the vulcanite mouthpiece of his pipe, measuring 2½ in. in length. Until then we had heard