This physiological senescence is quite independent of any external factor; its date of onset varies with the different subjects. It may be premature—that is, it may manifest itself at a period of life which is not that of normal senescence of the individual. I do not think this process is limited to the large group of family diseases of the nervous system, nor even to that of family and hereditary diseases considered as a whole. I am convinced that this is a far-reaching process, which deserves to be placed on a par with intoxication and infection, the two factors which nowadays govern medical pathology.

Nowhere could I have discussed these questions of nervous and general pathology before an audience better prepared to hear them. You belong to a nation which has achieved conspicuous distinction in general medicine, and especially in neurology. Do you not reckon among your great men of the past Sydenham, Fothergill, Charles Bell, Brodie, Abercrombie, Parkinson, Todd, Graves, Gull, Waller, Little, Lockhart Clarke, and Sir William Broadbent, who died recently, and who was the first President of the English Section of the Entente Cordiale Médicale? And at the present time can you not be justly proud of your brilliant phalanx of neurologists, who would not forgive me were I to sing their praises? In the name of French medical science I greet them as the worthy successors to a glorious past.

An Address

ON THE

PRESENT POSITION OF ABDOMINAL HYSTERECTOMY FOR FIBROIDS IN LONDON.

DELIVERED TO THE GLOUCESTER BRANCH OF THE
BRITISH MEDICAL ASSOCIATION,

BY JOHN BLAND-SUTTON, F.R.C.S.ENG.,

SURGEON TO THE MIDDLESEX HOSPITAL; SENIOR SURGEON TO THE
CHELSEA HOSPITAL FOR WOMEN.

It is surely a triumph of surgical art to improve an operation which in 1896 was attended with such a high mortality that few surgeons performed it, and then only in highly favourable, uncomplicated, and carefully selected cases; yet by 1906 the mortality of this operation in some hospitals in London had fallen to less than 2 per cent. This is the history of abdominal hysterectomy for fibroids in London.

In the year 1896 the results of abdominal hysterectomy for fibroids in eight hospitals in London were as follows:

St. Bartholomew's	7 with 3 deaths.
St. George's	1 " 0 "
St. Thomas's... ..	5 " 2 "
Middlesex	6 " 1 "
University College	3 " 0 "
Hospital for Women (Soho) ...	1 " 0 "
Samaritan	17 " 4 "
Chelsea Hospital for Women ...	9 " 1 "
	49 11

Ten years later—that is, in 1906—the latest returns from the same sources, and the New Hospital for Women, are:

St. Bartholomew's	26 with 4 deaths.
St. George's	8 " 0 "
St. Thomas's... ..	40 " 2 "
Middlesex	50 " 0 "
University College	21 " 1 "
Hospital for Women (Soho) ...	60 " 1 "
Samaritan (for Women)	37 " 2 "
Chelsea (for Women)	80 " 1 "
New Hospital for Women	26 " 0 "
	348 11

In regard to my own results: During the years 1906 and 1907, at the Middlesex Hospital and the Chelsea Hospital for Women, I performed abdominal hysterectomy for fibroids on 101 patients; they all recovered.* I performed abdominal hysterectomy in these two hospitals for other morbid conditions and the patients recovered.

* The short notes of the 101 hysterectomies for fibroids, and the subsequent history of the patients, are published in the *Journal of Obstetrics and Gynaecology of the British Empire*, May, 1908.

The notes concerning the operations in this series have been compared with the hospital books by the Registrars, Mr. Somerville Hastings, F.R.C.S. (Middlesex Hospital), and Mr. Stanley Dodd, M.B., M.A. (Chelsea Hospital for Women), and these gentlemen allow me to state that they have verified them.

It will be gathered from a perusal of these statistics that the removal of the uterus by the abdominal route, when it is occupied by troublesome fibroids, has become a fairly safe proceeding, nevertheless, hysterectomy, like all other major surgical operations, is attended with risks, even when it is carried out with all the careful surroundings which form such a prominent feature of modern surgical operations.

Setting aside the risks and sequelae common to operations in general, such as dangers from the anaesthetic, pulmonary complications, and the like, there are a few special dangers connected with it which require careful consideration at the hands of those who are daily occupied with this class of surgery. In this communication I shall deal with some of them in detail. I shall also consider the question which is much debated at the present time—namely, the advantages and disadvantages of the two methods in vogue for removing the uterus, subtotal and total hysterectomy, especially in connexion with an important matter, the occurrence of cancer in the cervical stump left after subtotal hysterectomy. The fate and value of belated ovaries is also well worth consideration, for some recent writers in America seem to attach little importance to the retention of ovarian tissue.

In addition to careful statements on these matters, which are of high importance for the subsequent comfort of patients, I propose to discuss certain matters concerning the pathology of uterine fibroids in relation to carcinoma and sarcoma, and to some secondary changes which affect them, especially in relation to pregnancy. These observations, more or less of a pathological kind, have become of great value in assisting us in the clinical side of our work; and they have helped to convince even the most experienced that "fibroids," which were commonly supposed to be the most easily recognized species of pelvic tumour, are not only simulated by other tumours, but in the secondary changes to which they are prone, mimic other well-known morbid conditions of the internal pelvic reproductive organs and lead to startling errors in diagnosis.

It is no part of my intention this evening to discuss details of technique, because the operation of hysterectomy, as a radical method of treating uterine fibroids, is now practised extensively throughout the civilized world; but I am anxious to put in a plea for the employment of sterilized rubber gloves by those who perform this operation or take any part in it, whether they be surgeons, assistants, or nurses. It not only reduces the mortality of the operation to a minimum, but the unpleasant sequelae are almost abolished also, especially trouble with the ligatures and sutures.

There is a risk which attends the total removal of the uterus, whether by the abdominal or the vaginal route, which should be referred to, namely, the liability of injuring one or both ureters. This subject I have discussed recently so that I need only refer to it here, but I mention it because it is an accident which demands the anxious consideration of all surgeons who perform hysterectomy.¹

Unpleasant sequelae, and especially that form of sudden death during convalescence which is usually attributed to pulmonary embolism, are more frequent in the practice of those who habitually operate slowly. It is a natural inference that a patient whose operation is completed in thirty minutes by a dexterous and careful surgeon, has a better chance of recovery and less liability to post-operative complications than the patient whose operation is performed by one who requires two hours for a simple hysterectomy. Everyday experience teaches that celerity in operating counts in favour of the patient.

TOTAL AND SUBTOTAL HYSTERECTOMY.

The great success which followed the use of the short ligature in ovariectomy induced several surgeons to apply the same principle to the cervical pedicle when removing the uterus for fibroids. The result was dismal failure. Matters improved somewhat after Koeberlé introduced the serre-noeud, and this continued the safest method until

1892. In the meantime antiseptics had begun to take effect in pelvic surgery, and attempts were made by Bardenheuer (1881), Polk, and other surgeons to avoid the dangerous difficulties connected with the treatment of the stump, by removing the cervix as well as the uterus (total hysterectomy), and some attained an encouraging measure of success. Nevertheless other surgeons (Goffe, Milton, Heywood Smith and Stimson) felt that the enucleation of the cervix was not always necessary, and sought to find a way of avoiding it. The credit of solving the difficulty fell to Baer, of Philadelphia (1892), for he showed that it is dangerous to constrict the neck of the uterus with ligatures, it is only necessary to secure the arteries.

Baer's method of supravaginal hysterectomy, or, as it is now commonly termed, the subtotal operation, soon supplanted the total method of Bardenheuer. The publication of Baer's paper had great consequences, it came at a time when the attention of surgeons was centred on improvements in hysterectomy.

The method was promptly tested and adopted in London. The effects of this improvement in technique in a few years revolutionized the surgical treatment of uterine fibroids, as the statistical results which I have set before you amply prove.

The great advantage of Baer's method is its simplicity and safety; but there is a disposition on the part of a few surgeons to prefer the total operation, mainly on the ground that the cervical stump left after subtotal hysterectomy is liable to become attacked by cancer.

As far as I can ascertain, Dr. M. Mann, of Buffalo, was the first to draw attention to the occurrence of cancer in the neck of the uterus after the body of the organ had been removed. He stated in 1893 that he "removed an ovarian tumour and the body of the uterus, by accident, along with it; the cervix was left." The patient recovered. "Six months afterwards cancer developed in the cervix, from which she died."

When cases of cancer, supposed to arise in the stump left after subtotal hysterectomy, come to be critically analysed, they fall into four groups:

1. The disease was present in the neck of the uterus at the time of the primary operation, but was overlooked.
2. Cancer attacked the cervical stump subsequent to subtotal hysterectomy.
3. The fibroid which necessitated the hysterectomy was really a sarcomatous tumour of the uterus.
4. The suspected growth on the cervix is not malignant but a granuloma.

Each of these postulates requires separate consideration. Many observations have been published which show beyond dispute that surgeons have performed subtotal hysterectomy in ignorance that the cervix was already cancerous, and the haemorrhages of which the patients complained before the operation were due as much to the cancer in the neck of the uterus as to the fibroids. This should serve as a warning that in cases where the surgeon contemplates performing a subtotal hysterectomy he should carefully examine the cervix beforehand; at the time of the operation he should also critically examine the cut surface of the cervix, and if it be in the least suspicious he should remove the whole neck of the uterus. It is necessary to remember that cancer attacks any part of the cervical endometrium, therefore an early cancerous ulcer in the middle of the cervix will run a great chance of being missed by a surgeon who is content with a subtotal hysterectomy.

It is certain that cancer does occasionally attack a cervical stump left after subtotal hysterectomy at such an interval after the operation as to make it certain that the cancer did not exist at the time of the operation. Such a case occurred in my practice. I performed subtotal hysterectomy in 1901 on a woman 42 years of age, mother of one child; eighteen months later there was a cancerous ulcer on the cervix; the whole of the cervical stump was promptly removed and the nature of the disease established microscopically. At this date (1908) the patient is in excellent health.

In another case under my care I performed total hysterectomy for fibroids in ignorance that the patient had cancer of the cervix. Some months after the operation, cancer recurred in the vaginal vault and scar of the hysterectomy; the neck of the uterus had been preserved by the medical attendant, and on examination the cancer was found. In

this instance, although total hysterectomy was performed it had no effect in staying the course of the disease.

It is necessary to utter a caution in regard to the occurrence of cancer of the cervix after subtotal hysterectomy. I removed a uterus containing a large globular submucous fibroid from a barren married woman 45 years of age. Six years later she came under my observation with a large granulating and bleeding growth on the cervix uteri. I had no doubt from the naked-eye characters that this was a primary carcinoma, although it surprised me to find it there, especially as the woman had never been pregnant. On my urgent representations she allowed me to remove the cervix. On microscopic examination the suspected cancer turned out to be a granuloma. Two years later the patient was in good health. Polk has recorded a similar experience. These facts show that caution is necessary in accepting reports of cancer of the uterine stump after subtotal hysterectomy.

Cancer of the Body of the Uterus and Fibroids.

In deciding between total and subtotal hysterectomy for fibroids, the probable presence of cancer requires consideration in another aspect. Many writers in recent years have expressed their suspicions that the presence of a submucous fibroid favours the development of cancer in the corporeal endometrium. Piquand, in 1905, drew attention to this matter, and emphasized what other observers had pointed out—namely, that a submucous fibroid is often associated with changes in the mucous membrane of the uterus, which not only causes excessive bleeding, but sets up inflammatory conditions giving rise to leucorrhoea, salpingitis, pyosalpinx and morbid changes in the endometrium, rendering it susceptible to cancer. His statistics support his conclusions, for they represent that in 1,000 women with fibroids, 15 will probably have cancer of the body of the uterus. My own observations support this opinion. This complication is found most frequently between the fiftieth and the sixtieth years of life. If we narrow the ages of the patient and exhibit the liability in its most emphatic form it would run thus—that in patients submitted to hysterectomy for fibroids over the age of 50 years, about 10 per cent. of them will have cancer of the corporeal endometrium.

In 1906, I looked through the case notes of 500 patients who had been submitted to operation for uterine fibroids under my care. Of these 63 patients had attained the age of 50 years and upwards. Among these 63 women there were 8 cases of cancer of the corporeal endometrium, the nature of the disease in each case being verified by careful microscopic examination.² Consequently, in performing subtotal hysterectomy for fibroids in women of 50 years and upwards, the surgeon should have the uterus opened immediately after its removal, and assure himself that the endometrium is free from cancer. If there be any suspicion in this direction he should remove the cervix.

Sarcoma.

The most insidious danger which besets the surgeon in dealing with fibroids of the uterus is the occurrence of an encapsulated sarcoma in the guise of an innocent fibroid. I have for some years dropped the name of myoma for these common uterine tumours, preferring to apply the term fibroid in a generic sense to all encapsulated tumours of the uterus. Every histological condition is found in them—from the hard calcified body looking like a block of coral to a soft diffuent collection of oedematous connective tissue, and tumours composed of tissue indistinguishable from spindle-celled sarcomata.

I have elsewhere recorded briefly a case in which I removed the uterus from a woman 40 years of age which contained a tumour as big as an ostrich's egg. On section it appeared to be a moderately firm fibroid, with its tissue whorled, as is usual in hard fibroids, and enclosed in a complete capsule. Some months later the patient complained of pain, and on examination a hard mass occupied the floor of the pelvis. A portion of this was excised and submitted to three competent histologists, who reported the growth to be an innocent fibroid. The patient died fourteen months after the primary operation with her pelvis filled with recurrent growth. The tumour was a spindle-celled sarcoma.

Much has been written regarding the sarcomatous degeneration of fibroids. In this matter I have maintained

an attitude of active scepticism. My experience amounts to this: The case which I have briefly described is the only example in a thousand cases of hysterectomy in which an encapsulated sarcoma in the guise of an innocent fibroid has come under my observation, therefore I come to the conclusion that it is an uncommon event, and on turning to the literature of the subject you will find that unequivocal examples are few.

Sarcoma of the uterus occurs, certainly, but not often as an encapsulated tumour. A curious example is represented in Fig. 1. The uterus is beset with multiple tumours, some of which project from the wall of the organ like subserous fibroids. These tumours were soft and compressible when living, and the interstitial lumps visible on the cut surface of the uterus look like common fibroids, but when examined carefully they are seen to lack capsules, and on microscopic examination exhibit the characters of spindle-celled sarcomata. In such a case the surgeon could not stumble in regard to the nature of the tumour. Unfortunately, here even total hysterectomy avails little, but it relieves pain at least for a time. From a careful study of the question, I have formed the opinion that if a woman with fibroids and concomitant cancer of the neck of the uterus seeks advice on account of haemorrhage, and the cancer has attacked the vaginal portion of the cervix, the nature of the case will be appreciated. The cases likely to be overlooked are those where the cancer is situated somewhat higher in the cervical canal than usual, so that it is not easily detected by the examining finger, and so low in the cervix that the disease

Fig. 1.—A uterus beset with sarcomatous knobs resembling subserous fibroids.

is not exposed when the body of the uterus is amputated in the course of a subtotal hysterectomy. A knowledge of this, as well as the fact that cancer of the cervix is almost exclusively a disease of women who had been pregnant, should make the surgeon particularly careful in performing subtotal hysterectomy for fibroids in women who have had children, and assure himself that it is not complicated with cancer.

In addition to the liability of the stump left after subtotal hysterectomy to become cancerous, it is stated by some surgeons that the patient is more liable to intestinal obstruction than after the total operation. This objection is easily met, because a perusal of their writings shows clearly that they do not perform the operation properly.

In subtotal hysterectomy performed according to Baer's instructions there should be no stump projecting from the pelvic floor, but merely a thin seam underlying the base of the bladder.

I have dealt in detail with these two methods of hysterectomy, because when it can be performed subtotal hysterectomy is, as a rule, a simpler operation than total hysterectomy. There are conditions in which it is imperative to remove the whole of the cervix, especially when the canal is very patulous and perhaps septic; when it is large and hard or large and spongy; and especially if there is the least suspicion of malignancy in the cervix or in the body of the uterus.

It must, however, be borne in mind that cancer has

attacked the scar left in the vagina after a total hysterectomy (Quénu). At the present time the subtotal method enjoys the greatest favour in London; but it must be remembered that where the total operation is most indicated it is often difficult of execution.

Although I have a decided preference for the subtotal operation, especially in spinsters and barren wives, I have performed total hysterectomy in more than 200 patients, so that I am in no way blind to its merits.

THE FATE AND VALUE OF BELATED OVARIES.

The only improvement of any importance made in Baer's operation of subtotal hysterectomy concerns the ovaries. These Baer removed with the Fallopian tubes; but in 1897 I advocated at the Obstetrical Society, London, that they were of great value to the patient, and pointed out that their conservation, when healthy, spared the patient the annoyance of that curious vasomotor phenomenon known to women as "flushings," which is the only obtrusive sign of the menopause.

It is now admitted by those surgeons in London who have had much experience in hysterectomy for fibroids that the immediate results of preserving at least one healthy ovary in this operation are admirable, especially in women under forty years of age, for the retention of an ovary is of striking value "in warding off the severity of an artificial menopause" (Crewdson Thomas).

Fig. 2.—The same uterus in section, showing that many of the nodules lack capsules.

Although I have left one or both ovaries in the performance of abdominal hysterectomy for fibroids in more than 300 patients, in only two instances have I found anything detrimental in the practice. In these two patients it was necessary to remove one of the ovaries subsequently. Since 1906 I have modified the method by leaving only one ovary, even when both were healthy, and find that the immediate good consequences of the operation are in no way impaired. There is reason to believe that whatever good effects follow the practice of leaving a belated ovary (that is, an ovary divorced from the uterus and left in the pelvis) they are temporary, for in the course of a few years the ovarian tissue disappears and the patients experience the usual symptoms of the menopause. It is possible that the rate of atrophy of the secreting tissue of a belated ovary depends on the age at which a patient is submitted to hysterectomy.

In 1898 I performed subtotal hysterectomy on a woman, 31 years of age, for fibroids, conserving the right ovary. Nine years later (1907) I operated again for intestinal obstruction and found this ovary healthy and functional, for a ripe corpus luteum was visible on its surface. Even a portion of an ovary, if it contain follicles, will maintain menstruation.

In performing abdominal hysterectomy for fibroids there are three points which require consideration in the interests of the patients, and they depend mainly on the conservation of a healthy ovary. These three points relate to

(a) the patient's comfort in securing freedom from flushings; (b) if she be married, her marital relations; and (c), if single, her nubility.

In regard to marital relations in women with a belated ovary nothing trustworthy is forthcoming, but I believe the retention of an ovary is an additional factor in promoting domestic bliss. The question of nubility is interesting. I am able to state that spinsters who have had subtotal hysterectomy performed with conservation of one ovary have married and lived happily with their husbands; and I am of opinion that the preservation of the vaginal segment of the neck of the uterus is an important factor, as it leaves the vagina intact, and though such women are sterile they are certainly nubile. Without overstating the case it may be said that a belated ovary is a very precious possession to a woman under 40 years of age, whether she be married or single. Some experienced observers maintain that an ovary is valuable to any woman who menstruates, even at the age of 50 years, the persistence of menstruation being obtrusive evidence that the gland is functional.

In regard to the fate of such ovaries, in the present condition of our knowledge it may be stated that—

In a woman under the fortieth year of life, a belated ovary remains active and discharges ova.

An ovary belated after the fortieth year of life atrophies and menopause symptoms will often ensue in the course of a few months after the operation. The retention of an ovary minimizes the menopause disturbances, and they are never so acute and prominent under these conditions as they are when the menopause is induced by the sudden and complete removal of all ovarian tissue.

Experimental evidence obtained from rabbits proves that the removal of the whole uterus has no deterrent effect on ovulation, and it does not prevent the occurrence of oestrus and ovulation at periodical recurring intervals.

There is no necessity to appeal to experiments on animals in this matter, as clinical observations on women are most eloquent in proclaiming the great value of a conserved ovary when the uterus is removed on account of troublesome and dangerous fibroids.

UTERINE FIBROIDS IN RELATION TO PREGNANCY.

Our knowledge of the natural history of fibroids has been increased and rendered more accurate since hysterectomy has been freely practised. The belief that fibroids disappear at the menopause is a bubble that has been pricked by the scalpel. The frequent prolongation of menstruation by fibroids many years beyond the date of the normal menopause is a fact. The suspicion that submucous fibroids predispose to cancer of the corporeal endometrium is likely to become a grim reality; and whilst case collectors are arguing whether fibroids cause sterility, or sterility cause fibroids, we know that women with fibroids often conceive; and that when fibroids and pregnancy coexist it is often a malicious combination.

Obstetricians urge that the dangers of fibroids complicating pregnancy have been exaggerated, but if any one takes the trouble to inquire into the matter he will find them real enough. No one suggests that a few fibroids of the size of acorns or of a golf ball in the wall of the uterus are inimical to successful pregnancy; but when a tumour the size of a fist occupies the neck of the uterus, or grows from the lower segment of the body of the uterus, it is a different matter.

Among the new things which surgical treatment of fibroids has brought to light is a knowledge of that change to which these tumours are liable known as "red degeneration." This increase in our knowledge of the pathology of fibroids has been extremely useful in diagnosis. Red degeneration is especially liable to occur in fibroids lodged in a pregnant uterus, and it has the extraordinary effect of rendering them painful. One of the most striking features of a fibroid known is its painlessness, and equally remarkable is the painfulness and tenderness of a fibroid when in the condition known as red degeneration, but these signs are only markedly exhibited by such fibroids when associated with pregnancy, and they are a source of many errors in diagnosis.

If those who are interested in this matter take the trouble to read the clinical reports of cases in which abdominal hysterectomy or myomectomy has been performed during pregnancy two facts will quickly assert themselves.

1. A considerable number of the hysterectomies, particularly those carried out in the early stages of pregnancy, were performed in ignorance that the women were pregnant. The patients, many of them spinsters, complained of the increasing size of the tumour.

2. Many abdominal myomectomies have been performed on an erroneous diagnosis. A fair proportion of these errors is due to ignorance of the fact that red degeneration, when well established, produces pain.

In many instances the pain is sudden and acute, resembling that experienced by patients suffering from an ovarian cyst with an acutely twisted pedicle, or the shock produced by the bursting or abortion of a gravid Fallopian tube.

Although the directions in surgical writings are clear concerning the course to be pursued when pregnancy or labour is complicated by fibroids, the difficulty which confronts the surgeon, when he is face to face with an actual case, is the great uncertainty which often exists regarding the nature of the tumour. Although he may begin an operation under the impression that he has to deal with an ovarian tumour, it may prove to be a fibroid, a tumour of the pelvic wall, a misplaced spleen, a kidney in the hollow of the sacrum, a tubal pregnancy, a sequestered extrauterine fetus (lithopaedion), or a calcified hydatid cyst. In an unusual case, an experienced surgeon (Olshausen) removed a gravid uterus under the impression that it contained a cystic fibroid which would obstruct delivery. When it was examined after removal, the suspected fibroid proved to be a large sacral teratoma growing from the fetus.

It sometimes happens that a child is born, and then the uterus extrudes a large submucous fibroid. This so resembles the head of a fetus that practitioners in several instances have attempted to pull it out in mistake for a twin, and even applied forceps. I have recorded an example in 1898 and have found other records of the same kind.

Many curious errors are made in determining the nature of pelvic tumours even after the abdomen is opened.

Some years ago a woman was submitted to Caesarean section in a London hospital, because delivery was obstructed by a solid tumour. In the course of the operation, a tumour, being firmly incarcerated in the pelvis by the gravid uterus, was regarded as a sarcoma growing from the pelvic bones, and therefore irremovable. Four years later the woman came under my care at the Middlesex Hospital. The tumour had grown and reached above the navel. I performed coeliotomy, and found it to be a solid ovarian (fibroid) tumour and removed it. The woman recovered.

It is important that the operator should satisfy himself as to the nature of the obstructing tumour. Blacker performed Porro's operation in a case of supposed calcified fibroid obstructing delivery. "A week after the operation the wound just above the stump gave way, and a number of hydatid vesicles were discharged, the tumour in reality being a calcified hydatid cyst."

The method of treatment which commends itself to obstetricians in London at the present time, when pregnancy is complicated with fibroids, consists in keeping the patient under observation until the child is viable. In many instances the pregnancy progresses to near term, and the child is born without difficulty. In other women troubles arise which necessitate operative interference. Then the child can be delivered by Caesarean section, and the uterus with its fibroids removed, either by the subtotal or total method, according to the tendencies of the surgeon in whose hands she falls.

It would greatly help to place the treatment of this complication on a sure foundation if those who practise midwifery would record the cases in which patients with this complication come to grief whilst waiting for the fetus to become viable.

It is only necessary to study the *Transactions of the Obstetrical Society of London* to be assured that pregnant women with fibroids do frequently require aid from surgery, and that such efforts are rewarded with success. There is no condition of the uterus which simplifies hysterectomy so much as pregnancy.

In some lectures *On the Surgery of Pregnancy and Labour Complicated with Tumours*, published in 1901, I drew attention to the marked tolerance exhibited by the pregnant uterus to abdominal myomectomy, and that it

was possible to enucleate large and troublesome sessile fibroids from the wall of the uterus without disturbing the pregnancy. On that occasion I published a table of cases showing the good results which follow from the operation.³

Since 1900 more than 30 cases of hysterectomy and myomectomy for fibroids complicating pregnancy and labour have been published in the *Transactions of the Obstetrical Society of London*. These records include cases which show that when the uterus contains fibroids, either of the submucous or the subserous pedunculated kind, and the pregnancy goes to term and safe delivery ensues, the patient's risks are by no means at an end, for the necrotic changes sometimes place her life in such grave peril that she is only rescued at the risk of myomectomy or hysterectomy.

The view which I expressed in 1901 has been amply confirmed by a greater experience of this matter. It amounts to this:

In a very large number of instances fibroids and pregnancy coexist and no harm ensues, for though the tumour occupies the pelvis in the early stages, it is so much "part and parcel" of the uterus that as the enlarging organ rises out of the pelvis, it carries the fibroid with it.

I am of opinion that ovarian tumours give more trouble to parturient women than fibroids; but fibroids have been far more lethal, as they frequently destroy pregnant women from sepsis.

If any practitioner has any doubts as to the reality of these risks, I would advise him to read the clinical history of a case recorded by Dr. Tate in the *Transactions of the Obstetrical Society*, 1907, in which a patient with a uterine fibroid miscarried of a dead fetus at the sixth month, and endured a grave septic illness which involved the removal of the uterus seven months later.

For one case of this kind which is published many remain unrecorded, and it is useless to urge that antisepsis in midwifery has abolished the dire consequences which often arise when fibroids complicate pregnancy.

THROMBOSIS AND EMBOLISM AFTER HYSTERECTOMY.

Thrombosis.

After operations on the pelvic organs, thrombosis occasionally occurs in the iliac, femoral, and saphena veins, accompanied by fever, pain, especially along the course of the saphena vein, and oedema of the limb. It is noticed most frequently about the twelfth day after operation. In some patients the thrombosis is confined to the superficial veins of the calf and thigh, but when the femoral and internal iliacs are involved the oedema is of a solid kind. Apart from the danger which ensues from the detachment of a fragment of clot and its arrest in the pulmonary artery, this complication is often very serious for the patient, for it entails a long confinement to bed, a tedious convalescence, and the oedema of the limb will sometimes persist for many weeks or months, in spite of topical applications, careful bandaging, or judicious massage.

Post-operative thrombosis was formerly fairly common after hysterectomy for fibroids, and in the later stages of malignant disease of the uterus. Its frequency after operations for fibroids was attributed to the profound anaemia in patients who had severe and exhausting metrorrhagia. I am convinced that it is due to sepsis, and since we have made it an almost general practice to operate with the hands carefully protected by sterilized rubber gloves this serious complication of pelvic operations has almost disappeared.

In several instances I have caused the clot found in thrombosed veins to be examined bacteriologically, and pathogenic microscopic organisms have been isolated. A careful comparison of post-operative results in a general hospital showed that thrombosis was much more frequent in the wards of the surgeons who worked with bare hands than among the patients of the surgeons who systematically wore sterilized rubber gloves.

I am also satisfied that in some cases of thrombosis of the veins of the thigh, especially those limited to the saphena vein, the clotting spreads from the superficial veins of the hypogastrium, which are infected from the abdominal incision.

Embolicism of the Pulmonary Artery.

In perusing the clinical histories of a series of cases of ovariectomy, hysterectomy, myomectomy, and, indeed,

after almost any surgical operation, here and there a record may be read to this effect: "The patient appeared to be doing well after the operation, when she sat up, laughed and chatted with the nurse, then suddenly fell back and died in a few minutes."

Anything more tragic than this it is difficult to conceive, and, as a rule, after such a sad occurrence the relatives are so distressed that they rarely permit an examination of the body. Death in such circumstances is usually attributed to embolism of the pulmonary artery. In some instances this is an assumption, but there are many in which an embolus has been demonstrated, and a few in which the source has been detected.

Post-operative embolism of the pulmonary artery is an important matter for surgeons interested in the operative treatment of uterine fibroids, for it follows such operations more frequently than any other.

In order to afford some notion of the relative liability of patients to this accident after subtotal and total hysterectomy for fibroids, I have gathered the following statistics, which are interesting as showing an extraordinary variation in the practice of different operators:

Baldy ascertained that among 366 operations for fibroids in the Gynecean Hospital, Philadelphia, there were 13 sudden deaths attributed to pulmonary embolism.

W. Duncan, in 18 abdominal hysterectomies, lost one patient from this cause. Spencer, in 85 total hysterectomies, had two deaths from pulmonary embolism. R. Lyle, in 8 cases of subtotal hysterectomy, had one sudden death.

Mallet collected the records of 1,800 coeliotomies: there were six deaths attributed to embolism, and of these, three followed operations for uterine fibroids. Charles P. Noble, in 42 vaginal myomectomies, lost two patients, one from septic endocarditis, the other from embolism; in the latter case the fibroid was gangrenous.

Olshausen, from the year 1894 to the end of 1905, performed 366 hysterectomies for fibroids; twenty-seven of these patients died. Five of the fatal cases were due to embolism.

Since 1894 I have performed more than a thousand operations of various kinds for fibroids, and have lost one patient from pulmonary embolism. This happened in 1900. The woman was 45 years of age, and profoundly anaemic from profuse and long-continued menorrhagia. Twelve days after subtotal hysterectomy she asked to be pillowed up in bed; this was done, when she suddenly slipped down the bed in agony and died in fifteen minutes. At the *post-mortem* examination the right pulmonary artery was found plugged with a thick clot. No thrombosed vessels were found in the pelvis.

The symptoms of pulmonary embolism may occur at any period from the hour of the operation up to the thirtieth day. In the majority of patients embolism happens about the twelfth day. The symptoms supervene with great suddenness, and seem to be preceded by movement, such as sitting up, getting out of bed, and especially straining during defaecation. Withrow tells of a patient who was attacked whilst "putting on her clothes to leave the hospital." She died in twelve hours. Reclus, at a meeting of the Société de Paris, 1897, mentioned that a patient quitting the hospital, apparently convalescent from hysterectomy, fell dead in the courtyard from pulmonary embolism.

It is important to note that these fatal cases of pulmonary embolism occur when they are least expected, and it is an unusual sequence in patients with obvious thrombosis of the femoral and saphena veins.

The most constant symptoms are urgent dyspnoea accompanied by great distress; in some instances the patient becomes pallid, and in others cyanotic. Death may follow in a few minutes; in some instances it is delayed several hours; the patient remains conscious, but suffers severe mental agony.

A pulmonary embolism is not necessarily fatal, for a woman after a pelvic operation may complain of sudden pain in the chest, urgent dyspnoea, exhibit great mental distress, and in a short time spit up sputum mixed with blood. In a few hours the urgent symptoms subside, and in two or three days pass away and the patient recovers. I have seen five examples of this mild form of pulmonary embolism after hysterectomy. One of the patients appeared to suffer from a succession of small pulmonary emboli.

Somerville Hastings, in an admirable article,⁴ refers to a woman 36 years of age, anaemic from profuse, long-continued menorrhagia due to a uterine fibroid, who, whilst waiting in the hospital for hysterectomy, was seized with

pulmonary embolism and died three hours later. An embolus occupied the pulmonary artery, resembling a blood clot found in the left common and internal iliac veins. Hastings also states that in a patient who died from pulmonary embolism after an operation a thrombus occupied the right cardiac ventricle, and he thought it possible that this intraventricular clot furnished the embolus.

We must bear in mind that individuals apparently in good health die suddenly in the street, in the arm-chair, in a bath, or even during sleep; it is a fair assumption that some of the instances of sudden death occurring during convalescence from surgical operations may be due to failure of the heart absolutely unconnected with the operation. It is, however, undeniable that thrombosis of the pelvic veins after ovariectomy or hysterectomy is a source of fatal emboli.

At present there is very little evidence available as to the cause of the thrombosis, but it can scarcely be doubted that sepsis—it may be only of a mild type—is responsible for some of the cases.

A careful consideration of the matter reveals beyond any doubt that pulmonary embolism occurs much more frequently after hysterectomy for fibroids than after any other operation, and it is especially liable to happen in women who are profoundly anaemic from profuse and prolonged menorrhagia. This indicates that long-continued and irregular losses of blood induce some change in the composition of this important fluid which favours its coagulation.

It is probable that the thrombosis is due to a combination of causes, such, for example as sepsis, cardiac disease, chill and exposure, especially in prolonged and difficult operations, damage to the walls of blood vessels, and circulatory disturbances due to anaesthetics.

It has been suggested that the practice of keeping patients strictly confined to bed for two or three weeks after hysterectomy, and allied operations, is responsible for the thrombosis which is the source of these fatal emboli. Some American surgeons act on this suggestion, and insist on their patients getting out of bed a few days after such operations. This method does not commend itself to British surgeons. In my own practice I make it a rule, even in the most favourable conditions, to keep the patients confined to bed for two weeks. No patient is allowed up until her temperature has been normal for at least three days.

In cases of embolism of the pulmonary artery death does not always occur immediately, but may be postponed for an hour or more after the lodgement of the embolus. On one occasion I saw a patient within ten minutes of the lodgement of a pulmonary embolus. The patient, a woman 35 years of age, was livid but did not breathe, yet her heart beat regularly and forcibly for five minutes after my arrival, then stopped suddenly.

Trendelenburg is of opinion that in some of these cases it might be possible to remove this clot by surgical intervention. After careful consideration of the matter he carried out this operation on a woman aged 63 years. He raised an osteoplastic flap on the left side of the thorax, exposed the conus arteriosus, and intended to withdraw the clot through a slit in its walls by means of a specially-constructed pump. The patient died from excessive bleeding before the clot could be extracted. The operation was hindered by an adherent pericardium.

When patients who are profoundly anaemic from menorrhagia due to fibroids undergo hysterectomy, it is a useful measure to give them 20 grains of sodium citrate twice daily in order to diminish the abnormal tendency of the blood to coagulate in the vessels. Certainly this drug should be administered if there is the least evidence of thrombosis. I have seen good consequences follow its use.

In conclusion let me reiterate that the history of hysterectomy, as a radical method of treating uterine fibroids, during the last ten years has been a record of continuous improvement in the hospitals of London. Like all valuable operations, especially those connected with the reproductive organs, hysterectomy for fibroids has had to pass through a probational period of prejudice. To-day no major operation can show so small a mortality, and what is more to the point, no major operation produces such permanent good effects.

Surgery can now enable a wife suffering from "an issue

of blood," and spending half her life as a chronic invalid, to become a companion to her husband; or, a spinster afflicted with a bleeding fibroid, and living in humble circumstances, to earn her own living, and cease to be a domestic incubus.

It is a great gratification to realize that hysterectomy for fibroids occupies in London to-day an established position in surgery from which no misrepresentation can raze it.

REFERENCES.

¹ *Journal of Obstetrics and Gynaecology of the British Empire*, May, 1908. ² *Ibid.*, 1906, x, 1. ³ *Lancet*, 1901, i, 452. ⁴ *Archives of the Middlesex Hospital*, 1907, xi, 78.

THE ETIOLOGY OF RICKETS: A CLINICAL AND EXPERIMENTAL STUDY.*

By LEONARD FINDLAY, M.D.,

ASSISTANT TO PROFESSOR OF PATHOLOGY, AND CLINICAL TUTOR WITH PROFESSOR OF CLINICAL MEDICINE, GLASGOW UNIVERSITY; EXTRA DISPENSARY PHYSICIAN, WESTERN INFIRMARY, AND EXTRA HONORARY PHYSICIAN TO OUTDOOR DEPARTMENT OF ROYAL HOSPITAL FOR SICK CHILDREN, GLASGOW.

[WITH SPECIAL PLATE.]

MANY and varied have been the causes assigned for rickets. Almost every prejudicial circumstance in the life of the infant—and in that of the parent, too, for that matter—has been cited as the etiological factor.

ETIOLOGICAL THEORIES.

According to Kassowitz¹ and some of the German school, it is of congenital origin, a theory, however, mainly founded, at least by its earlier adherents, on macroscopic appearances alone, and not on the finding of any characteristic histological changes in the bones. In every infant the bones are soft, the abdomen may be protuberant, the fontanelles will be open, and there will be some swelling of the costo-chondral junctions, and yet it was on the ground of these facts alone that such a theory was based. The later advocates² of this theory, however, did examine the minute histological changes in the bones, but as much of the material studied was obtained from stillborn children, and as the pathological anatomy of osteochondritis syphilitica and rachitis was at that time only imperfectly understood, these two conditions were much confused, and it is not improbable, therefore, that many of the supposed rachitic lesions were in reality of a syphilitic nature. As these two processes came to be differentiated from one another, each succeeding published record of work done on this subject showed a marked diminution in the proportion of congenital rickets. Kassowitz in 1882 considered that 80 per cent. of all children born were rachitic; in 1897 Tschistowitsch³ put the proportion at 12 per cent., while more recently (1902) Escher,⁴ after investigating a large series of cases both clinically and histologically, was only able in one single instance to make a diagnosis of congenital rickets.

There is of course such a condition known and described as "fetal rickets," but it is considered by the majority of observers an entirely different disease to what ensues after birth, and is, moreover, as rare as ordinary rickets is common. It will probably be better then, at least for the present, to classify it apart, and designate it by one of its pseudonyms—for example, "osteogenesis imperfecta."

Closely allied to the above theory (congenital origin of rickets) is that mainly advocated by Siebert,⁵ who holds that the condition is hereditary, and dependent on some inherited weakness or predisposition. This after all is merely a case of begging the question and failing to discuss the main points at issue. Hausen,⁶ however, reports the case of a stallion who begot seven rachitic foals. Later two of the mares, one of which had given birth to three and the other to two of these rachitic foals, became pregnant to another stallion, and bore healthy non-rachitic offspring.

It is during extrauterine life, however, that most present-day authorities believe the etiological factors play their part, and the general consensus of opinion would ascribe the disease to incorrect feeding, bad hygiene, and want of sunlight. Of all these possible malign influences it is some error in feeding which receives most credence and support.

* The work was carried out in the Pathological and Physiological Laboratories, University of Glasgow.