liver. The necrotic transformation may be so complete that the centre of the tumour is of a watery or pultaceous consistence, and this may then be absorbed, leading to the disappearance of the tumour. In other cases only the central parts have undergone degeneration, and then we have the firm white, whorled appearance of the normal fibroid unchanged at the circumference of the tumour, surrounding areas of the mahogany-coloured degeneration. The condition is a degeneration or aseptic necrobiosis of the component parts of the tumour. The colour seems to be due to the diffusion of the colouring matters of blood throughout the degenerating part. Fairbairn,³ who first drew particular attention to the subject and investigated the pathology, states that "there is a marked deficiency in the nuclear staining of the tissue cells, and in the more advanced cases this had resulted in an almost total absence of nuclei in the section, leaving only wavy bundles of fibres, which had taken up the ground stain; in some cases the muscle bundles showed signs of granular or hyaline change." How or why pregnancy plays a part in setting up these changes is not clear, but that it does so is probable. Although the condition occurs in non-pregnant women, it attains its most typical form only during pregnancy. Although it is not considered that any organisms play a part in this degeneration, fibroids undergoing this change may cause quite alarming constitutional and local symptoms. It seems not unlikely that these are the expression of toxic absorption of products from the breaking-down tumours. The following case, bearing on these points, came under my care some months ago:

A woman aged 25, a patient of Dr. Robertshaw of Stockbridge, and four months pregnant, was referred to me, complaining of severe pain in the abdomen with constant vomiting.

History.

Four days before admission into hospital the patient had been seized with very severe pain in the lower and left quadrant of the abdomen. She could not sleep for pain during the next night, and next day the pain became so unbearable that she kept to bed. At the same time she became aware that there was a tumour in the abdomen in the painful region, which was extremely tender to touch. Vomiting also commenced, and became very troublesome for the last twenty-four hours before admission, she having vomited everything that she swallowed. For some four or five months she had had increased frequency of micturition, especially during the night, and this had been getting more troublesome of late. Menstruation up to five months ago had been quite regular, not profuse, but accompanied with a good deal of pain. When I saw her the patient was lying on her back, with the knees drawn up, complaining of very much pain in the lower abdomen. The pulse was 120, temperature 101°, the tongue brown and dry. There was general distension of the abdomen, its walls were slightly rigid, and slight tenderness was everywhere present. The summit of the uterus could be felt some little way above the pubes, and attached to its upper left angle was a tumour, somewhat greater in size than two large fists, very tender to touch and very movable; it could be pushed about the abdomen in various directions. Misled by the mobility, I considered that we had to deal with the twisted pedicle of a solid ovarian tumour.

Operation.

On opening the abdomen, it was found to contain a considerable quantity of serous fluid, and the supposed ovarian tumour turned out to be a fibroid attached by a somewhat wide base to the left of the summit and back of a very movable pregnant uterus. Below it was another small one. The fibroids were easily enucleated and the wounds so made carefully sutured with catgut. The peritoneum covering the tumour showed considerable injection of the small vessels and also a number of small haemorrhages, and the peritoneum of the flank also showed signs of commencing inflammation. The progress of the patient was in every way satisfactory until the tenth day. On this day—the stitches having been removed on the previous day—the patient attempted to vomit, and to our consternation the abdominal wound burst open and a quantity of the serous fluid escaped. The wound, which had been sutured previously in three layers, was again sutured in the same way, with the exception that strong silkworm supporting stitches to the deeper layers were this time used.

After-history.

The patient made an excellent recovery, went home well, and five months afterwards was safely confined of a living child. The tumour, on section, shows over the greater part of the area the well-marked mahogany or liver colour of red degeneration. The capsule is roughened, and shows a number of small haemorrhages in its substance. The patient's abdomen became natural in appearance, and all trace of fluid quickly disappeared.

Dr. Frank Taylor,⁴ in a recent full and excellent paper on the subject of Necrobiosis in Fibroids, together with

many other illustrative cases, relates a very similar one to the above which occurred in the practice of Mr. Bland-Sutton, in which, although the same mistake in the primary diagnosis was made, the same treatment was followed by a similar good result.

The interesting points about my case were the constitutional disturbance, the great abdominal tenderness, and the great quantity of fluid that was present in the peritoneal cavity. The presence of this fluid was indeed curious. It might be that it was the result of the irritation of the peritoneum by the products of the change in the fibroids; indeed, there might have been some infection of the fibroid by the colon bacillus, and the fluid be comparable to that which is sometimes found in large quantity in cases of appendicitis, even when there is no gangrene or perforation of the appendix. Unfortunately, no bacteriological examination was made, so that the question cannot be decided.

The foregoing remarks imperfectly deal with a series of cases which are alike in that in all of them there was the sudden onset of acute and alarming intra-abdominal mischief during the course of pregnancy, and in all the involvement in greater or lesser degree of the peritoneum, with its risks to the lives of the patients and to the continuance of the pregnancy. Incidentally is shown how they all may be successfully dealt with by prompt surgical interference.

REFERENCES.

- ¹ Connell, *Journal of Obstetrics and Gynaecology of the British Empire*, August, 1906. ² Heaton, *BRITISH MEDICAL JOURNAL*, 1905. ³ Fairbairn, *Journal of Obstetrics and Gynaecology of the British Empire*, August, 1903. ⁴ Taylor, *Practitioner*, June, 1906.

ON THE MENTAL NERVE AREA AND ITS RELATION TO THE GREYNESS OF HAIR.

BY

G. LENTHAL CHEATLE, C.B., F.R.C.S.,

SURGEON TO KING'S COLLEGE HOSPITAL AND ITALIAN HOSPITAL.

[WITH SPECIAL PLATE.]

THE mental nerve area is unique in this way: it is the only nerve area which can be completely covered with hair, and in which, at the same time, there is so little overlap from other nerves. All other nerve areas are either more overlapped by neighbouring nerves, or they are not completely covered with hair. Hence the mental nerve area is the most interesting and instructive nerve area in the human body in which to study the incidence and spread of the hair's greyiness or whiteness.

It will be seen in this article that the points of incidence and area of spread of greyiness indicate a neurotrophic cause for both to perfect demonstration.

Figs. 1 and 2 show the mental region (blackened) as worked out by Dr. Henry Head as being the area which becomes tender in certain affections of the organs of the head and neck. The figures are reproduced with his kind permission from his article in *Brain*, vol. xvii, On the Disturbances of Sensation, with Special Reference to the Pain of Visceral Disease. Figs. 3 and 4 are photographs of a man whose beard has become white on the area seen in Figs. 1 and 2. Fig. 3 is very striking in that it shows the absence of whiteness beneath the lip on the man's right hand side, and the absence of whiteness below the lip on the left side is the same, but, unfortunately, the moustache covers the beard on this side. Compare Fig. 3 with Figs. 1 and 2, and it will be seen that the mental nerve area described by Dr. Head does not cover the chin immediately below the mucous membrane of the lower lip.

It can be claimed that the posterior margin of the white area of the beard in Fig. 4 corresponds exactly with the posterior limit of the mental area in Dr. Head's diagram in Fig. 2, hence the area of whiteness is identical with the mental area described by Dr. Head. I have seen the mental areas practically free from white hairs whilst the whiskers have been white, and the line of demarcation between the two corresponded exactly to the posterior line of Dr. Head's mental area. Fig. 5 shows the maximum point of the mental area, and is taken from the same article of Dr. Head to which reference has just been made. Fig. 6 shows the beard of a man whose greyiness began on this maximum point on both sides; the left side began first. The posterior limit to the

Fig. 3.

Fig. 4.

Fig. 9.

Fig. 6.

Fig. 8.

mental nerve supply as seen in Fig. 7 is already marked on the left side of the man's beard. Fig. 7. represents Merkel's areas of distribution of the branches of the fifth cranial nerve, and is reproduced from *Die Oberflächlichen Nerven des Kopfes*, von Dr. Med. Fritz Frohse. Figs. 8 and 9 show this area abruptly limited by the white hairs of these men. The greyiness of the mental nerve area is sometimes associated with greyiness of the superior laryngeal nerve area.

Curiously enough, on the lower lip of the patient from whom Fig. 9 is photographed there were two lesions: on

Fig. 1.

the left side of the middle line was a typical squamous epithelioma, and on the right side of the middle line was a sessile wart of very suspicious appearance.

With regard to the incidence of neurotrophic diseases of the lower lip, I have seen leucoderma, scleroderma, herpes zoster, greyiness of hair, on the lip, and I regard the blackish-yellow patches of pigmentation which appear on the mucous membrane and skin of people who are comparatively advanced in years as a neurotrophic lesion, and this, too, I have seen on the lower lip, as well as on the temples, face, and backs of hands.

The fact that the cancer on the lip of Fig. 9 began on the same point and at the same time of life as some of

Fig. 2.

these lesions is, I consider, a point in favour of the arguments that the genesis of cancer is not due to the local action of an external organism, and that if it is due to the action of an organism it cannot be the sole cause.

As bearing on the prolific growth which can be attained by hair in this region, it is interesting to consider the following facts:

In connexion with my idea upon the nerve influence upon growth, I have often considered the experiment of John Hunter, in which he essayed to demonstrate that increased blood supply led to increased growth of a part.

It will be remembered that Hunter transplanted a cock's spur into a cock's comb. The result to the spur was that it grew to about 6 in. in length; and, according to Hunter, this result was due to the increase of blood supply from the very vascular tissue of the cock's comb.

In thinking over this experiment, it is at once evident that it proves that growth is not influenced only by the amount or quality of the nourishment supplied to a part, for the cock's comb itself had stopped growing, although, according to Hunter, its vascularity was capable of inducing extraordinary growth in another structure. Now,

Fig. 5.

if an enormous blood supply were the only factor influencing growth, the cock's comb ought to grow as well as the cock's spur to which it was supplying nourishment. Being assured by these preliminary conjectures that though the facts were correct, Hunter's explanation of them was probably wrong, in November, 1907, I transplanted the spurs of cocks into the birds' backs as well as into their combs, and since that time the rate of the spur's growth has been the same in all cases. Those in the cocks' combs do not appear to grow faster or bigger than those in the birds' backs. In about two years' time, when the growth has attained greater proportions, I hope to be able to show these birds.

Fig. 7.

In talking over the result of my experiment with Dr. Arthur Whitfield, he at once suggested what appears to me to be the correct explanation of Hunter's and my own experiments. Dr. Whitfield's explanation is that the excessive growth of a spur transplanted to abnormal sites such as the comb and the back is due to the fact that it is not worn away by the friction which usually keeps a cock's spurs short, and which occurs as the result of the bird's social, domestic, and bellicose existence. There are many experiments and facts which support this explanation, but into which I propose to enter not now, but at a future time.