3. The advantages claimed for ether by the open method are all present, though these do not seem to be greater than those of maintaining the anaesthesia with chloroform. The patient may be readily anaesthetized more deeply should this be necessary. The initial induction is as at present practised, except that the small bag only need be carried. The use of the valves as an adjunct to the "small bag method" of gas and ether administration automatically avoids any rebreathing during induction. It is customary to avoid this when valves are not employed by removing the lower part of the facepiece during expiration.

The disadvantages of the method are that the anaesthesia tends to be light, that there is oozing of blood, but it is now red instead of bluish, that the patient is kept on ether for an increased time, often much to his discomfort afterwards, and that the purity of the surrounding air is vitiated.

On the whole the administration of ether by the open method seems to have its uses in a small number of cases, chiefly in very anaemic patients, for short operations, and in those who readily become cyanosed.

ON RECTAL DRAINAGE IN CASES OF PELVIC ABSCESS DUE TO APPENDICITIS.

By BILTON POLLARD, B.S., F.R.C.S.,

SURGEON TO UNIVERSITY COLLEGE HOSPITAL.

In the June number of the *Annals of Surgery*, received in London, July 3rd, I find a short paper by Dr. Archibald MacLaren, Professor of Clinical Surgery in the University of Minnesota, on Pelvic Abscess with Special Reference to Rectal Drainage. I was myself about to publish a short paper on this very point, and notwithstanding the appearance of Dr. MacLaren's paper I think it may be of service to emphasize still further the value of rectal drainage in the pelvic abscesses resulting from appendicitis. I have advocated and practised this method of evacuating pelvic abscesses for several years, but I believe I am right in saying that it is a method rarely adopted by English surgeons—indeed I have heard of its being characterized as "not surgery." It is well known that in the days before appendicitis and appendix abscesses were understood and treated by operation cases occasionally recovered after the spontaneous evacuation of an abscess through the vagina or the rectum, and the same fortunate event from time to time occurs nowadays. Nature's method of cure is often a good one to follow and the subject under consideration is a good instance in point.

Pelvic abscesses in connexion with appendicitis may be arranged in two groups:

1. An appendix abscess has formed above the true pelvis in the iliac fossa, and has been opened through the anterior abdominal wall. At the time of the operation no collection of pus can be detected in the pelvis by rectal or vaginal examination. The temperature falls, and all goes well for a few days. The temperature then goes up again, and the pulse increases in frequency. The drainage of the abscess already opened appears to be quite satisfactory, and consequently a further examination of the pelvis by rectal or vaginal examination is made, and the relapse is explained by the presence of a collection of pus in the basin of the pelvis.

2. The illness has begun in the manner characteristic of appendicitis. The temperature continues high and the pulse rapid, and so perhaps a week goes by. No mass is felt by abdominal palpation, but as the case is not progressing satisfactorily a surgeon is called in, and his routine rectal examination discovers a pelvic swelling bulging the anterior rectal wall, either alone or in association with resistance or swelling behind the right rectus muscle.

In the first group the adhesions of the bowels intervening between the first abscess and the collection of pus in the pelvis have become so firm and the separation of the coils of intestine from one another so indistinct that the attempt to open the pelvic collection through the abscess cavity already opened would be attended with grave risk of injury to the intestine. The difficulties just mentioned have been met by operating in the left linea semilunaris. This plan involves a second abdominal

incision, and the risks attendant upon evacuating an abscess through the abdominal cavity, protected though it may be by gauze packing. It is in these cases that Dr. MacLaren has opened the pelvic abscess through the rectum. I have several times done the same thing, and always with success. I have operated thus in females as well as in males. I have selected the rectal route in preference to the vaginal in females, because in my cases the abscess has bulged more towards the rectum than the vagina, and it has appeared that a rectal opening would be more dependent than a vaginal one. With regard to cases in the second group, Dr. MacLaren writes:

There is another type which I have not operated upon—namely, the man who is desperately sick and who presents a decided fullness in the cul-de-sac. In such a case I will, on my first opportunity, simply open the cul-de-sac and drain, and not make an anterior incision at all.

I have recently dealt with a case of this type precisely in the way advocated by Dr. MacLaren, and I propose to give a short account of the case in order to illustrate my method of operating, which is very similar to that employed by Dr. MacLaren:

G. O., a man aged 26, was admitted to University College Hospital on June 16th, 1908. His illness began nearly three weeks before his admission with severe abdominal pain and sickness. The pain was mostly central at first, but on admission it was worst on the left side below the level of the umbilicus. The lower part of the abdomen was distended, and did not move much during respiration. There was well-marked tenderness all over the lower part of the abdomen, and a distinct mass could be felt coming up from the pelvis, especially on the left side. By rectal examination the pelvis was found to be occupied by a large mass which bulged into the rectum, especially on its anterior and left lateral aspects. The swelling was very tender, and the most prominent part of it, which was about 3½ in. from the anus, was soft. The temperature was 101.4° F., and the pulse 88.

Rectal drainage was decided upon. The patient was anaesthetized with chloroform, and placed in the lithotomy position with the pelvis well raised and rotated forwards. The sphincter ani was well stretched, and the anus was held widely open by three retractors, one on either side and one behind. The bulging swelling was thus completely displayed. A trocar was pushed into it, and pus found. The mucous membrane was divided longitudinally with scissors for 2 in., and the other structures were split and stretched by means of polypus forceps introduced through the trocar puncture. About a pint of stinking pus was evacuated. The opening into the recto-vesical pouch was maintained by means of a tube an inch in length and an inch in diameter, with a flange at each end. One flange was passed into the abscess and the other remained in the rectum. The tube was left in position for eight days, and the abscess cavity was irrigated daily through the tube, and after its removal the irrigation was continued through the wound for three days more. The temperature fell to normal a few hours after the operation and the patient made an uninterrupted and rapid recovery.

If this case had been dealt with by an abdominal incision it would have been necessary to have made it free enough to allow of the introduction of abundant gauze packings in order to protect the rest of the abdomen from infection during the evacuation of the pus. There would have been some risk of infection of the peritoneal cavity; but, putting this at a minimum, the use of the tube and the almost certain infection of the abdominal wound would have been followed by a weak scar, which would probably have been on the left side, and hence at an unsuitable place for the subsequent appendectomy.

In dealing with a well-localized appendix abscess in any situation, it is, in my opinion, unwise to do more than is absolutely necessary for the efficient evacuation and drainage of the abscess. The appendix should not in these cases be removed at the time the abscess is drained, for its complete removal involves the breaking down of limiting adhesions and the risk of peritoneal infection. Holding this view, I am very averse from removal of the appendix in the original operation for pelvic abscess when this is done through an abdominal incision; and so I am not deterred from using the rectal route on account of the impossibility of removing the appendix through it.

Gynaecologists constantly use the vaginal route for evacuating pelvic abscesses, and I believe that surgeons would be wise in adopting the vaginal or rectal route in females and the rectal route in males for the evacuation of large appendix abscesses which are apparently confined to the pelvic cavity.