which possess the power of destroying the uric acid, or a portion of it when once formed. These we may call *uricolytic ferments*.

IV A.

I wish now to draw *very summarily* attention to the relation of the nucleins, nucleo-proteids, the nucleic acids, and particularly the purin bases, to the formation of uric acid as bearing on the pathology of gout, and as affecting those suffering from gout or the gouty diathesis.

Since the time when Dr. (afterwards Sir) Alfred Baring Garrod first drew attention of the medical and scientific world to the statement that during attacks of gout the amount of uric acid in the blood is increased—whilst prior to, and usually during, an attack of acute gout the amount of uric acid eliminated in the urine is diminished, an abnormal amount being excreted after the attack—the medical profession, especially in England, came to the conclusion that uric acid must be looked upon, in a sense, as the cause of gout, the *materies morbi*. As a matter of fact, the conditions are by no means always as stated above. The uric acid in the blood is not always increased in acute gout, and the conditions of the urine do not always agree with that which is stated above—"uric acid storms," that is, abnormal eliminations of uric acid, sometimes occurring during the acme of the gouty paroxysm.

All kinds of theories and various régimes have been based upon this uric acid hypothesis, which, viewed from the wider standpoint of to-day, are seen to be far from expressing the truth—at any rate, *the whole truth*—on the matter. Believing uric acid (at any rate, the exogenous fraction) to be closely related to a meat diet, there have been many authorities on gout who have insisted upon a complete absence of meat in the dietary of the gouty. It has been shown, however, that the gouty manifestations (even the deposits of chalk stones, which are composed of sodium diurate, in the joints) cannot be explained on the mere uric acid hypothesis. In the first place, the facts of retention of uric acid in the system before an attack of gout and the alleged constant excess of uric acid in the blood are far from constant phenomena; in the second place, it appears that a very moderate meat diet exerts singularly little influence on the excretion of uric acid, even in the gouty, *always providing that the nucleo-proteids, the nucleic acids and the purin bases are (so far as possible) eliminated from the diet*. On the latter matter there appears to be now a consensus of opinion. Still we have to account for or, at any rate, provide the most probable hypothesis to explain the phenomena of the gouty diathesis and the occurrence of attacks of gout. Facts appear to indicate that in the gouty there is a diminished power of the system to deal with the purin bases (so that there is a tendency to their accumulating in the system); it appears probable that this is connected with a deficiency or inefficiency of the ferments which are concerned in their normal transformations—a deficiency or inefficiency which may be due to the action of toxins generated in the body.

It may be, too, that there is a deficiency of, or interference with, the enzymes which normally destroy a part of the uric acid formed, a process which has by several distinguished scientific men been supposed to have its principal, though not exclusive, seat in the liver.

In any case, all authorities on gout are agreed that, so far as possible, the purin bases in the diet should be kept at as low a level as possible.

The force of the arguments advanced in this paper in reference to the use or non-use of yeast extracts will escape no one who is in a position to follow them.

To characterize a preparation which (for the sake of argument) we shall assume to contain 18 per cent. of meat extract added to an extract of yeast mixed with considerable quantities of gelatin, as a preparation which excels in value pure meat extracts, is to endanger the health of the large number amongst us who have succeeded to the unfortunate heritage of the gouty diathesis.

REFERENCES.

- ¹ Ruppel, op. cit., p. 175. ² *Zeitschrift für Biologie*, Band iv., 1868, S. 3.
- ³ Hefenextracts, *Zeitschrift für Hygiene u. Infektions-Krankheiten*, vol. xiii, 1903, pp. 461-466. ⁴ Arthur Gamgee and Walter Jones, "Ueber die Nukleoproteide des Pankreas, der Thymus und der Neben-niere mit Besonderer Berücksichtigung ihrer optischen Akti vität," *Hofmeister's Beitrage*, vol. iv, 1903, p. 10. ⁵ Gamgee and Croft Hill in *Hofmeister's Beitrage*, vol. iv, 1903.
- ⁶ Karl Micko, "Untersuchung von Fleisch, Hefen, und anderen Extracten auf Xanthin-körper. (I. Die Xanthin-körper des Fleisch extracts. 2. Die Xanthin-körper der Hefenextracts.) *Zeitschrift d. Nahrungs und Genussmitteln*, vol. vi, 781-91; vol. vii, 257-59; vol. viii, 225-37.

A Third Report

ON

CLINICAL EXPERIENCES WITH SPINAL ANALGESIA:

WITH A

THIRD SERIES OF ONE HUNDRED CASES.

By ARTHUR E. BARKER, F.R.C.S.,

PROFESSOR OF SURGERY AT UNIVERSITY COLLEGE HOSPITAL MEDICAL SCHOOL AND SURGEON TO THE HOSPITAL; CONSULTING SURGEON TO THE QUEEN ALEXANDRA MILITARY HOSPITAL, S.W.

AMONG the large number and variety of operations performed under different anaesthetics which have fallen to my lot at University College Hospital since my first and second 100 spinal analgesias were completed and published in the *BRITISH MEDICAL JOURNAL* of March 23rd, 1907, and February 1st, 1908, a fresh 100, operated on under the same method of analgesia, are now to be found. These have been carefully noted and entered from day to day in the same form of tables as before by Captain H. Browne-Mason, Lieutenant Treves, and Captain H. Simpson, of the R.A.M.C., to whom I am indebted for the accuracy and fullness of the records, as well as to Mr. H. Morriston Davies, who has checked them. A summary of these tables added to the previous series for the sake of contrast is given below in fifties as before.

SUMMARY OF TABLES.

	Series 1.	Series 2.	Series 3.	Series 4.	Series 5.	Series 6.	Total.
Strangulated herniae ...	4	0	1	2	4	1	12
R. cure herniae ing. fem. vent. ...	10	7	16	11	10	14	68
Acute appendicitis ...	2	5	3	3	3	0	16
Appendectomy à froid ...	5	3	2	3	4	4	21
Gastro-enterostomy ...	0	0	2	1	2	0	5
Other laparotomies ...	4	5	2	2	3	5	21
Kidney operations ...	0	0	0	1	0	0	1
Excis. rectum (Kraske) ...	0	0	1	0	0	0	1
Perforated gastric ulcer ...	—	—	—	—	1	0	1
Ectopic gestation ...	—	—	—	—	1	0	1
Gall bladder ...	—	—	—	—	1	0	1
Total abdominal ...	25	20	27	23	29	24	148
Piles (Whitehead) and fistula ...	7	10	6	9	2	11	45
Pubis and genitals ...	7	6	1	2	8	5	29
Dislocation of hip reduction ...	0	0	0	1	0	0	1
Amputation of thigh, leg, foot ...	1	0	1	1	1	0	4
Internal S. cart. and knee ...	2	2	5	5	1	1	16
Wiring fractured patella ...	0	3	2	2	0	2	9
Other operations on lower limb ...	6	8	8	7	8	6	43
Failed to puncture dural sac ...	2	1	0	0	1	1	5
Total abdominal ...	25	30	23	27	21	26	152
Total abdominal ...	25	20	27	23	29	24	148
	50	50	50	50	50	50	300
General anaesthetic required ...	14	0	6	0	3	0	

The study of these 300 consecutive cases, operated on in the same hospital and by the same surgeon, appears to possess a certain value, and the results are now placed at the disposal of others. It would seem to me premature to adopt any decisive attitude towards this comparatively new procedure. It is still *sub judice*, and must long

remain so unless we agree to ignore it altogether, which seems undesirable considering its importance as a possible substitute for general anaesthesia, the risks of which are acknowledged, and the extent to which it is already employed. It has never been claimed, so far as I know, that the procedure is without danger. It has always been simply a question whether its risks, immediate and remote, are greater or less than those of the ordinary accepted forms of general anaesthesia, and whether it offers better conditions for the performance of operation to the surgeon. And these questions cannot be answered until we possess a much larger experience of carefully recorded cases. As far as our observations go, the conditions produced by spinal analgesia appear to be as regards the actual performance of the operation more favourable for the surgeon, and not more disagreeable for the patient than those produced by general anaesthesia. I believe that deaths will unfortunately continue to occur after grave operations under any and every form of anaesthesia. The great difficulty is to determine which factors in a complicated case have contributed most to the fatal result. The general condition of the patient before the operation, the shock of the operation itself, the presence of toxæmia such as is present in many cases of advanced intestinal obstruction, have all to be taken account of in any exact estimate of the cause of death.

The principles and the technique of the procedure have remained the same as those described in the two papers referred to. Doubtless new light will be thrown both on the principles and practice in the future, just as some advance has been shown in our last series in some respects as contrasted with those before them, but we continue to work on the same broad lines at our hospital which have directed us hitherto.

As to the class of case in which spinal analgesia has been resorted to, it will be seen from the tables that in the last hundred there is a larger proportion of very serious abdominal conditions requiring a high analgesia than in the earlier series. This class of cases claim a greater familiarity with and confidence in the procedure, and renders the possibility of failure to produce adequate analgesia greater, besides probably enhancing the risks all round. The few inadequate analgesias in this series belong to the high operations.

It would be impossible to give all the abdominal operations at length, but a few may be mentioned by way of illustration. Among them are several enterectomies—one removal of the splenic flexure of the colon for cancer, one abdominal hysterectomy and pyosalpinx in the same case and at the same sitting, one ectopic gestation, and various exploratory laparotomies. Among these a few deserve a more detailed description.

F., aged 43, was admitted for faecal fistula in the region of the caecum dating from an operation in the provinces for some suppurative condition suspected to be tuberculous. There was a very large hard mass filling the iliac fossa. This turned out to be the caecum, ascending colon, and about 2 ft. of the last part of the ileum involved in the most extensive tuberculous disease I have ever seen in the intestines. The whole of this diseased tract was excised, and the ileum was united to the transverse colon. Recovery was quite uneventful. This operation lasted nearly two hours, involving very intricate dissection. Quite towards the end the patient began to feel the manipulations of the mesentery, but though advised to have a little chloroform for the remaining few minutes, stoutly refused, and apparently to her advantage. There was no shock, and only a little retching after the operation, throughout which she was in excellent condition.

F., aged 47, was sent into University College Hospital with a diagnosis of acute appendicitis. The usual onset had been noted, and there was a lump the size of two fists in the right iliac fossa, which was very tender. The incision for appendicitis on the right side was made, and the lump was found to be a multiple fibroma of the uterus covered with pus. On further examination the pelvis was discovered to be filled up by a large left pyosalpinx which had pushed up the fibromatous mass to the right. This was removed, and subsequently the adherent pyosalpinx was separated *en masse* from the pelvic walls, tied off, and removed. The wound was sutured in layers, and healed perfectly per primam under one dressing. Recovery was absolutely blameless. There was no shock, and no unpleasant after-effects.

F., aged 33, was a case of ectopic gestation, which, beyond the uncertainty of diagnosis and feebleness of the patient, presented no difficulties, and recovery was perfectly smooth under one dressing.

F., aged 31, was brought from the country suffering from perforated gastric ulcer. This case showed high spinal analgesia at its best. The muscles were quite relaxed, and consequently the manipulations and mopping-out of the inflammatory effusion

from the abdomen much easier than under the average general anaesthesia. There was, however, a hint of sensation on mopping out the subphrenic spaces, but not to a serious extent. The patient made an excellent recovery, without any trace of shock during or after operation.

But throughout our series of 300 cases we have noted from time to time cases in which marked pallor came on soon after the spinal injection, though there were no fatal cases resulting from it. This has always appeared to me to be best explained by cerebral anaemia due to the complete relaxation of the abdominal muscles, and consequent filling up of the abdominal veins, which have lost their support. It does not appear where the effects of the drug are designedly confined to the pelvis, as in pile and other pelvic cases. This faintness has usually been followed by slight retching, and then rapid return to comfort. In many cases the pulse has hardly been affected, it has often been slowed; in others it has fallen in tension while the faintness lasted. Had this been due to the general specific toxic action of the drug apart from the relaxation of the abdominal muscles, it might have been expected to last longer, whereas it has always, when present at all, been noted about eight minutes after injection, and passed off speedily after a little retching.

In some of the cases included in this third hundred the condition of the patient before operation was very grave, but all recovered, so that there is no fatality to record in the whole series due to the spinal analgesia, nor is there among private cases done during the same period but not included in our list. It is unnecessary to say that a case of perforated gastric ulcer with extensive extravasation into the abdomen is serious, and yet one of these cases showed the method at its best. Another of gangrenous femoral hernia, for which enterectomy was required, was in a very bad condition, but recovered from this operation. But the worst general condition to be contended against was in a patient of 48, operated on in the country for gangrene of nearly 5 ft. of the small intestine strangled by band. She was pulseless and cyanosed in face and hands, and the last pulse felt by her doctor had been between 120 and 130. About 6 cg. of stovaine placed her in comfort in five minutes, and I excised 4 ft. 9 in. of black gangrenous bowel, anastomosed the healthy gut above and below, with a perfectly blameless recovery. Before I left the house the patient's pulse was good and she had no unpleasant after-effects beyond bringing up at once a cup of tea given on the operating table. Again, in the case of a very stout lady who had nearly succumbed under chloroform a year and a half before when having an operation done on the gall bladder by another surgeon, two operations within six weeks which I found it necessary to do on the same region passed off without any serious trouble.

But from all this I should not like it to be inferred that I for my own part regard the procedure as without serious risk in debilitated patients where the upper part of the abdomen has to be dealt with. This is far from being the case. I think there may be great risks, but whether greater than with other anaesthetics it will be impossible to say until we have far larger experience. And it may be necessary to formulate some rule against operating on such cases later on. It is a remarkable fact that several cases briefly alluded to above have been in a very perilous condition of asthenia before operation, and have taken no harm from the spinal analgesia, while showing in their subsequent recovery an absence of those troubles from shock, vomiting, etc., so often seen after chloroform. And yet in two cases which I have operated on since the series of 300 was completed the result was quite different, and the fatal issue was almost beyond doubt contributed to by the method of spinal analgesia employed. It would be quite certain that this was the case, but that both were in a most serious condition before any injection was made, and had to pass through most trying operations. Though they do not belong to our 300 cases, but to any series to be published later, it is right that they should appear here for the guidance of others possibly inclined to employ this procedure. Whether they would have survived if general anaesthesia had been employed instead is an open question.

M., aged 62, looking older and feeble, suffering from extensive cancer of the stomach with secondary deposits (as it turned out) in the surrounding glands and liver. An ordinary injection of 6 cg. of stovaine was made and acted well. The patient became very faint with shallow breathing before the operation began. He then improved and the exploration was completed. He

seemed dazed but smiled, and was removed to the ward. Here he lapsed back into his former condition, and, in spite of the usual restoratives, sank not long after.

In this case the general effect of the drug, by whatever mechanism it acted, was injurious. But one has seen the same under general anaesthetics in similar cases of advanced disease of the stomach, and I have seen more than one flicker out, as it were, after even a short exploratory operation. The deduction would be, and is generally accepted, that operations on such decrepit aged people with advanced carcinoma in the epigastrium are always accompanied by great risk, and that the hope that these are obviated by spinal analgesia is not so far securely based. Shock played a part in this case, beyond a doubt, as was seen by the immediate effect of the manipulations of the stomach on the pulse. And this would hold for all cases of the kind under any anaesthetic so far known to us. Every chloroformist and surgeon is alive to the depressing effects of handling of the organs in the epigastrium, and, for my own part, I have for some years in many cases hesitated, and in some refused, even to explore such carcinomatous conditions in this region in decrepit individuals.

When carcinoma is accompanied by obstruction with large accumulation of putrid fluid faeces the risks are even greater. We have factors here the full weight of which we are unable as yet to accurately estimate. We have toxic infection with its fever and other depressing effects, also the prolonged reduction of proper alimentation, with consequent debility, great distension, and the special effects of the toxins on the lungs, kidneys, and probably other organs. These factors have to be kept in mind in the following unsuccessful case, though I have little doubt that the anaesthetic directly influenced the result in some way not so easy to explain:

F., aged 48, suffered from intermittent obstruction for three and a half months due to cancer of the descending colon culminating in acute obstruction, with raised temperature ten days or so before operation. She was fairly nourished but looked white and low. There was extreme distension of the colon with great straining against the obstruction, producing much pain and tenderness. At 4 minutes to 3 p.m., 5½ cg. of stovaine were injected without any trouble, and anaesthesia up to the fifth rib followed in a few minutes. The patient had the not unusual faintness in about eight minutes as the abdominal walls relaxed, and retched a little. The operation, which consisted in a left lateral oblique incision and division of the attachments of the meso-colon of the splenic flexure downwards until the growth could be brought out, necessitated much dragging on the peritoneum under the diaphragm. The transverse colon was greatly distended with fluid faeces, and was anastomosed to the sigmoid flexure. During the operation the patient seemed in deep sleep. When roused the sensation was tested and found as before—down to the fifth rib. Shortly after the breathing became irregular and sighing in character, and she began to look blue; the pulse, however, was fair, the conjunctival and pupil reflexes gone. Strychnine was injected and oxygen inhalations, which improved the colour, the pulse remaining fair as before. She was left on the operating table, as the work of the afternoon was practically over. Artificial respiration with the addition of oxygen was kept up for fifteen minutes, and she then took a long breath, and gradually voluntary respiration returned. It was perfectly regular and deep and 7 to 8 to the minute when timed by watch over and over again. The pulse remained fair and the colour without any oxygen was particularly good. An enormous quantity of foul liquid faeces came away per anum from time to time. This condition of things remained while she was carefully watched until about 7.45 p.m., when breathing again became halting, after more than two hours of regular breathing. Artificial respiration and oxygen were again employed, but death took place at about 8 p.m., heart and respiration failing at about the same time. The body temperature remained good all through. The day was very hot. The pupils all through remained of average size, but did not react. It appeared to those watching the case closely all through that the anaesthesia was passing off as well as the motor paralysis. For both the diaphragm and intercostals were working deeply for certainly more than one and a half hours before the end. But the shock apparently was not passing off, and appeared to be accentuated not long before the end by the large discharge of liquid faeces from the anus.

It is well to be careful in reviewing all the facts of the case before coming to a conclusion as to the cause of death. That the anaesthetic contributed to, but was not the sole cause, may be safely inferred. Under any known anaesthetic any severe operation on such a case was bound to involve great risk, as we know from experience. Whether the best form of anaesthesia for such a condition of things was employed is debatable as far as my personal experience of over 300 previous cases justifies me in expressing an opinion. But I have certainly operated on

cases under spinal analgesia which were to all appearance quite as or even more desperate with the best results as mentioned above. In many of them there was no evidence of severe shock; in this case shock was marked.

I have mentioned these two cases here, though they do not come into our third hundred, because they ought to be known and reflected on by all who are studying or practising spinal analgesia. It will be well not to make either too little or too much of them. As I have said in a previous paper, we may have a very valuable addition to surgery in this procedure which makes it worth cultivating, or it may have defects which will compel us to give it up. To be hasty in judgement in either direction would be unwise at present. If the first mischances met with in the earlier history of chloroform had led to its being discarded, we should have been deprived of one of the greatest blessings to humanity ever discovered.

From my own small experience I should be inclined to advise beginners not to use the method in extremely debilitated patients for severe operations on the abdomen requiring high analgesia. Our best results have been in operations below the umbilicus, and best of all on the pelvic organs, though, as shown above, some high severe operations passed off quite happily.

Again, the head and shoulders should be kept well raised at all times throughout the operation and after, and all shifting or jerking of the patient should be interdicted. A minute on the side after injection seems quite long enough, and then the patient should be gently rolled over on the back without any sudden movement or lifting of the pelvis. Finally, the need of care in the exact measurement and preparation of the drug cannot be too much insisted on.

It will be seen that the actual technique has improved in each hundred. In the first there were fourteen cases which required a general anaesthetic to complete the operation, in the second hundred six, and in the third only three cases. As to the actual puncture of the dural sac I failed in two cases in the last hundred. Both were oldish men with irregular spinous processes, and one was almost absolutely stone deaf, and could not be made to understand the position we wanted him to assume and the need of perfect stillness. In 12 cases tropacocaine was employed, but as the effect in some appeared somewhat depressing stovaine was as before resorted to. I therefore desisted after a trial. Both were in the lateral position.

SOME LESSONS FROM ANCIENT FRACTURES.

By FREDERIC WOOD JONES, M.B., B.S. LOND.

THE collection of specimens illustrating the pathology of the people of ancient Nubia, that was made under the directorship of Captain H. G. Lyons, F.R.S., of the Survey Department of the Egyptian Government, by Dr. Elliot Smith and myself, has recently been exhibited at the Royal College of Surgeons, to which body the whole series has been presented by the Egyptian Government.

Contained in this collection are some 200 cases of bones that have been broken in ancient times, and have healed; and, for the practical purposes of surgery, they are perhaps the most interesting of the relics of a bygone population.

These fractures were collected from a series of upwards of six thousand bodies buried in that tract of Nubia that lies between Sheilal and Demhid—that is, immediately south of the First Cataract (Aswan)—and they are from representatives of all periods comprised between early predynastic times (the earliest period from which human remains have come down to us, *circa* 4000 B.C.) and the Christian era, until about the fifth century after Christ. They therefore cover a period of the history of the human race of considerably over four thousand years.

Some brief notices of special cases have already appeared in the two *Bulletins of the Survey of Nubia*, published by the Egyptian Government, and a detailed account of them all will be given in the annual volume of the survey; but at a time when the ideas concerning the treatment of fractures is by no means so stereotyped as it used to be, a short summary of the incidence and result of fracture in an ancient people may be useful, apart from the mere historical interest.

In order to give some idea of the relative frequency of the fractures of different bones, I have taken all our cases