the sequence of repair in the human patient, for an injured animal must always fail entirely to obtain that most important factor in the repair of fracture—that is, rest.

[This report is published with the permission of Captain H. G. Lyons, F.R.S., Director-General of the Survey Department of the Egyptian Government.]

REFERENCE.

¹ Lewis A. Stimson. *A Treatise on Fractures and Dislocations.*

HYPERPYREXIA.

By JAMES SMITH, M.R.C.S.,

NEWCASTLE-UPON-TYNE.

THE following case presents clinical features of more than ordinary interest. On two previous occasions I have taken temperatures of 107° F. and 107.4° F., the former occurring at the crisis of a pneumonia, the latter within a few hours of the commencement of an influenzal attack. In both these cases the hyperpyrexia was of short duration, and recovery followed.

In the present instance the patient (Miss D. E., aged 20 years) developed a sore throat on the homeward journey from St. Moritz in the first week of February, 1908. She remained in London two days, where some local treatment was used, and then came home. I saw her on the second day after her arrival. She was febrile (101° F.), tongue slightly coated, was somewhat hoarse in speech, and complained of her throat. This on examination was unlike anything I had previously observed. Both tonsils (the right particularly) were studded with pustular looking ulcers, some located on the anterior fauces. The pharynx and soft palate were intensely inflamed and swollen. On the latter were several roughened, raised patches. The submaxillary glands were slightly enlarged; there was some difficulty in swallowing, considerable discomfort and pain. I prescribed a quinine mixture, to be taken every four hours, and kept the bowels active by a saline aperient. The local treatment consisted in frequent swabbing with antiseptic solutions, and for three days the application of a solution of silver nitrate. In the course of a week the throat was well, and my patient progressing rapidly. After being up for some days she unfortunately left her bedroom, the prevailing weather at the time being cold and damp. The next day I found her with right ankle and tarsus swollen and painful, and a temperature of 100° F. This was disappointing, seeing that nine years ago she had an attack of acute rheumatism. I now prescribed sodium salicylate gr. xij, every four hours. This, although resulting in free diaphoresis, had apparently little effect in arresting the progress of the disease. Daily one or more joints were involved, until, on February 22nd, practically the whole of the articulations of the lower and upper extremities, even to the fingers of both hands, had been implicated. The temperature varied from 100° F. to 102.2° F., pulse 84 to 88. There was no heart complication.

On February 23rd, at my second visit (7 p.m.), the temperature had risen to 103.4° F.; the pulse, which had been well maintained so far, was 100, regular, though of less volume, and weaker. At 11 p.m. the nurse in charge telephoned to say the temperature was 105° F. I ordered 10-grain tabloid phenacetin and caffeine and an additional dose of the salicylate mixture, and requested to be informed should this not result in a remission of the fever. The next message received was at 1.10 a.m. (February 24th), telling me the temperature was then 107° F. I set off at once and saw my patient about 1.50 a.m.; the temperature was now 108.2° F. She had an anxious, frightened expression, wanted to know what was wrong, her speech was tremulous, and she kept moving the head from side to side; the pupils were dilated; the tongue moist and clean; the radial pulse was quite uncountable—a mere flicker under the finger;

the skin was hot and dry, though to the sense of touch not so marked as one might have looked for; respiration was very short and frequent. Within twenty minutes of my arrival she became unconscious.

A cool pack was immediately applied, and replaced by a cold pack in fifteen minutes—the latter being carefully renewed every ten or fifteen minutes during the first hour. The temperature about 2 a.m. was 108.4° F. Within the next half-hour I took it myself and found it 108.6° F. Shortly after 3 a.m. the temperature had fallen to 107.2° F., and with a slightly improved pulse it seemed just possible the patient might survive. This hope, however, was speedily dissipated. I noticed small jerkings of the facial muscles, the patient vomited, and this was followed by severe general convulsions. A more desperate condition it would be difficult to imagine, each succeeding seizure jeopardizing life.

Dr. H. Paige (Gosforth) was now summoned, and with his concurrence a hypodermic injection of morphine, gr. $\frac{1}{4}$, and atropine $\frac{1}{100}$, was given immediately after his arrival. The cold packs were regularly continued, with cold applications to the head. Within half an hour after giving the hypodermic injection an abatement of the convulsions was noticeable, they gradually become less violent, with longer intervals. Between 3 and 4 a.m. the patient slept a short time. The temperature was still receding and pulse improving. About 4 a.m. she had several slight convulsions, and then fell into a sleep which lasted until 6.30 a.m., when, on awaking, there was some consciousness; she went asleep again quickly, and an hour after woke up perfectly conscious.

The temperature meantime was rapidly falling—107° F. at

4 a.m. it had dropped to 102° F. at 7 a.m.; the pulse-rate 100. After passing through this period of the utmost gravity and stress, it was very surprising to find my patient so far recovered. She was inclined to talk and took liquid nourishment freely.

During the forenoon the temperature again began to rise—104.8° F. at 2.30 p.m., it rapidly ascended to 106° F. at 4 p.m. The iced pack was applied, and at 6 p.m. the temperature stood at 102.6° F. Dr. Beattie (Newcastle) saw the patient this evening; it was decided to continue the salicylate treatment—15-grain doses every four hours, and, in addition, to give an occasional 2-grain dose of quinine sulph., which I had found useful in controlling the fever.

On the afternoon of February 25th and evenings of

February 26th and 27th the fever rose to 103.8° F., 103° F., 103° F. respectively, but tepid sponging in each instance effected reduction. From this time until February 29th a gradual declension of the fever took place; at 6 p.m. on this day the record was 99° F. The subsequent progress of the case was steady and without further complication. The joint inflammations gradually subsided, the heart escaping injury. Eventually, I am happy to say, my patient made an excellent recovery. I saw her last week (June 12th). She had just returned from Bath, after a month's residence there, is much improved, and able to get about with considerable freedom.

Appended is an hourly chart during the critical period of this attack.

In concluding this report, I must gratefully acknowledge my indebtedness to Dr. H. Paige for his ready and valued help in a position of the utmost urgency and responsibility, and for his co-operation in the after-conduct of the case. I must also recognize the efficient and untiring services of Nurse Worby and my patient's sister, which at the critical period were invaluable.

Taking a general survey of this case as a whole, we have displayed a clinical picture striking and suggestive. Commencing with a septic throat, followed after a short interval (during convalescence) by acute rheumatism, and culminating in extreme pyrexia and convulsions, does it not seem probable that more than one toxin was active and contributory?

I have in mind a case of fully-developed facial erysipelas seen in the evening for the first time. I noted a rash on the neck identical with that of scarlet fever and con-

tinuous with the erysipelatous rash. The following day the punctated rash was gone, but within ten days two members of the same family developed scarlet fever. The erysipelas had practically subsided on the third day; the patient was comparatively well, and both diseases were arrested, a result, one may surely infer, of toxins acting antagonistically. May we not, then, suppose the converse of this likewise to hold—that some toxins acting concurrently are supplementary, and combine in the production of an intensified resultant, and thus be an explanation of those sudden, irregular, and alarming manifestations of occasional occurrence during the course of febrile disorders?

SEVENTY-SIXTH ANNUAL MEETING

OF THE

British Medical Association.

Held at Sheffield on July 24th, 25th, 27th, 28th, 29th, 30th, and 31st.

PROCEEDINGS OF SECTIONS.

SECTION OF DENTAL SURGERY.

FRANK HARRISON, M.R.C.S.Eng., L.D.S.Edin.,
President.

PRESIDENT'S ADDRESS.

AFTER a few introductory remarks the PRESIDENT said: The investigation of the physical condition of school children by virtue of the Medical Inspection Act is welcome. The examination is to be made by medical men, who, together with other defects and diseases, will have to observe the condition of the teeth and report thereon. This will do good, because it will be a means of bringing home to the minds of a large number of medical men, education committees, and the public the really dreadful condition of the teeth of the rising generation and the diseases consequent upon that condition.

Let me emphasize the fact that dental disease is the most widespread and commonest of all diseases to which flesh is heir; not only in this country, but statistics show that where investigations have been made the condition is much the same over the greater part of Europe, and we have no reason to suppose that the rest of the world is any exception.

The micro-organic flora in dental caries runs wild, rendering the oral cavity intensely septic. It is repulsive to think of the soiling and poisoning of the pabulum in its very first process of digestion. That this oral sepsis cannot exist without its producing lowered vitality must be admitted. The proof is very evident by the rapid recovery of health and vigour after radical treatment. If it is granted that this minus dental function and its concomitant evils are a loss to the individual, and that this degenerate dental condition is so widespread, what must the loss mean to the community in the aggregate? Medical men and guardians of the young are every day recognizing more fully the advantages to be derived from early care of the teeth.

Is it not passing strange that at this meeting I am able to make the statement that the real cause of the commonest of all diseases is still unknown? What an opportunity presents itself for one of our modern universities, fully equipped with all the latest scientific instruments and expert professors, to solve this problem of the cause of dental caries, and what a benefit it would confer upon humanity at large!

It is to our physiologists and pathologists who have such advantages of systematic research work that we must look to give us the "open sesame!" to these locked-up secrets. At present we can only treat effects, and much good may be done thereby; preventive treatment, such as we have, and care of the young, are very beneficial. Still, the methods, although very elaborated, are more or less primitive, and can only touch the fringe of the trouble. We want to get at the cause and treat it.

To my mind, the problem would not prove so involved as at first sight might appear, and would most likely be solved by an exhaustive examination of the function or want of function of the salivary glands. However, without surmising further, it is clear that such investigation cannot be carried on by medical practitioners or dentists, but should be undertaken by research workers under the most advantageous conditions.

DISCUSSION ON

ANTRAL DISEASE IN RELATION TO SPECIAL AND GENERAL SURGERY.

I.—HERBERT TILLEY, M.D.Lond., F.R.C.S.Eng.,

Surgeon, Ear and Throat Department, University College Hospital.

THE breadth of the title chosen for this discussion is a matter for congratulation, for it is only when workers in the special branches of medicine and surgery recognize that these studies are merely offshoots of a great science and art, and that all the fundamental laws upon which the greater edifice has been built apply equally to special departments, that real scientific and lasting progress can be made.

The diseases of the maxillary antrum appeal to the general as well as the special surgeon, and the latter comprise the rhinologist or surgeon with special knowledge of nasal diseases, and the dental surgeon whose studies have been focussed on pathological conditions of the teeth. Speaking generally, it may be said that the general surgeon will be more familiar with those diseases which cause obvious external swelling or deformity of the upper jaw or of the adjacent soft parts, and so he will be consulted for large alveolar abscesses, dental cysts, malignant disease of the antrum, and necrosis of the upper jaw. Since these affections are also frequently seen by the rhinologist and dental surgeon, we may as well discuss them at once.

Alveolar Abscess.—When this occurs in the upper jaw its clinical manifestations are so familiar to us all that I need not waste time in discussing them. It would be well, however, to remind you that when an abscess forms at the root of one of the incisor teeth the swelling may appear in the floor of the nose, just within or beyond the vestibule, and cause nasal obstruction of a marked degree. When the canine tooth is affected the tumefaction may appear externally at the side of the nose below the nasal bone, and may seem at first sight to have little to do with a diseased tooth. If such abscesses become chronic, cholesterol crystals are often found in their contents. Treatment must be directed to removal of the diseased tooth and opening the alveolar abscess.

Dental Cysts.—These, by their expansion, often cause considerable external deformity, and may so encroach upon the antral cavity that the sinus is almost obliterated. Pain, as a rule, is slight or absent, the swelling over the anterior antral wall within the mouth frequently presents the so-called "egg-shell cracking" when pressed upon, the hard palate is never pressed downwards, and if the transillumination test be applied, the affected antrum is only slightly less translucent than the normal one, or at any rate it is not quite opaque. To be successful treatment must be radical, for nothing is more wearisome to surgeon and patient than the constant packing of the cyst cavity after it has been merely opened from the mouth. The most effectual, speedy, and certain treatment is to reflect the mucous membrane from the buccal aspect of the cyst wall, freely open the cyst, remove all its thin, bony wall, especially that portion which projects into and often almost entirely obliterates the true antral cavity, then remove the inner antral wall together with the anterior half of the inferior turbinal, so that a large permanent opening into the nasal cavity is provided. Finally, the bucco-antral incision is sutured with a couple of horsehair stitches, and immediate union of the wound will take place. Practically there is no after-treatment beyond the use of a warm alkaline nasal douche for two or three weeks.

Malignant Disease of the Antrum.—Sarcoma and epithelioma are the commonest forms of malignant disease which affect the antrum. Pain of an intense and boring character is often an early symptom in epithelioma. With