only be made with difficulty. A little urine was drawn from the bladder by catheter, and found to be intensely albuminous. Labour proceeded with tolerable rapidity, and at 1 p.m. she was delivered of a healthy living child. The placenta came away soon after the birth, and with it a gush of blood, which caused syncope. The uterus refused to contract under the influence of any stimulus; a slow oozing of blood continued; and she never spoke after the expulsion of the child, but died at 2 A.M.

In speaking of this case to the midwifery class, Dr. Smith mentioned that he had been called to another case a short time previously, in which a woman suffering from albuminuria and dropsy, went with labour, and nearly died of alarming hæmorrhage after delivery, without any symptoms of convulsions.

The first case offers some points of interest as regards the curability of certain cases of albuminuria and pregnancy, and the cause of this affection. Some writers, especially Dr. Litzmann, maintain that the renal disorder cannot be relieved before labour has occurred. With reference to the cause of albuminuria in pregnancy, it has been referred to a depraved state of the blood occurring in pregnancy, the renal affection being a secondary condition; and to pressure exerted on the kidney and its vessels by the gravid uterus, especially in primipara. The first view is that entertained by Professor Simpson; the second has been insisted upon by Dr. Cormack. In the case of E. T., and in others of a similar kind, occurring in the middle part of pregnancy, the disorder can scarcely be referred to pressure. The uterus in her case was not of large size: the muscles of the abdomen were very relaxed: in fact, there was no sign whatever of abdominal tension. Dr. Tyler Smith's own opinion is that, in many cases of albuminuria in pregnancy, the congestion of the kidney which produces the disorder is caused by uterine sympathetic irritation; the uterus irritating the kidney in these cases in the same way that it irritates the mamma, stomach, salivary glands, thyroid, etc., in other cases.

Great attention has been paid to the connexion between albuminuria and convulsions, but very little notice has been taken of uterine hæmorrhage in albuminuria. The two last cases are remarkable, as showing no tendency to convulsions, but as accompanied by marked inertia of the uterus after delivery. It is not difficult to understand that, in the poisoning of the blood and nervous system, which obtains in albuminuria, from the retention of urea in the circulation, and the loss of albumen, the nervous and muscular actions upon which the contraction of the uterus depends may be seriously impaired. The tendency to uterine inertia being recognised, practitioners can be on their guard against flooding as well as convulsions in these cases.

St. George's Hospital.

COMPOUND FRACTURE OF THE SKULL.

Communicated by GEORGE POLLOCK, Esq.

[Continued from page 1006.]

A MAN, aged 43 years, was admitted into the hospital on the 19th October 1854, under the care of Mr. Tatum, with a lacerated wound about an inch and a half in length over the anterior part of the left parietal bone. At the bottom of this wound a ridge of bone was readily felt, elevated about half an inch above the surface of the adjacent and surrounding fractured portions. The injury was occasioned by a blow from a piece of iron. The man was reported to have been insensible for half an hour after the accident; but subsequently recovered perfectly from the state of concussion. He then walked to the house of a medical man, then to a railway station (fifteen miles from town), came to London by the next train, and after this walked between two and three miles in town before arriving at the hospital. He did not recollect the blow upon the head;

but was perfectly collected in all other respects. He said he had lost a large quantity of blood; but all hæmorrhage had ceased before his admission.

Mr. Tatum exposed the fractured surface by an incision through the scalp, when some hæmorrhage was observed—blood making its escape from between the edges of the fractured bone. A small trephine was now applied; and on the outer table being cut through, it separated from the inner one; and now the inner one was found to be more extensively fractured than the external table. The depressed fragment of the inner table could not be raised without a second portion of bone being removed by the trephine. This was done at a lower angle of the fracture, and a circular piece of the whole thickness of the skull was taken away. The depressed portions were then readily elevated, and some removed. One piece was marked by the groove of the meningeal artery, and this vessel itself was observed bleeding, though it soon stopped. There was not much blood under the skull, but the dura mater was perceptibly separated from the bone near this portion of the artery. The man was quite sensible throughout the time of the operation.

Oct. 20th. There has been very slight oozing from the wound. In other respects, he appears in a satisfactory condition, with no symptoms of discomfort or distress. Tongue clean; pulse quiet. Fever diet.

Oct. 23rd. He continued improving till to-day. His manner is now rather more oppressed, and he complains of severe pain in the head. He vomited last evening, and this morning was attacked with a rigor. Pulse quiet and soft; tongue clean. The wound looking healthy.

9 P.M. The pain in the head continuing, he was ordered two grains of calomel every four hours.

Oct. 24th. Some puffiness to be observed over the left temple; the headache gone; and there appear no other active symptoms. The calomel to be omitted.

Oct. 26th. He has slept rather badly. Pulse 80, weak. Complaints of pain over the eyebrow and back of the head. The wound does not look quite healthy, and the discharge rather foul. He was ordered beef-tea and a small quantity of wine.

Oct. 28th. There is an erysipelatous blush over the forehead and scalp, accompanied with some œdema. Bowels not being open, he was ordered a dose of calomel and castor oil.

Oct. 29th. Last night the œdema having increased, some punctures were made in the scalp. He slept badly. Pulse 68, weak. He appears rather low, and speaks somewhat confusedly. There is much œdema at the back of the head. More incisions were requisite, and were made. He was ordered to take six ounces of red wine.

Oct. 30th. He appeared in all respects better. The wine was increased to ten ounces.

Subsequent to this date, he continued progressively improving, the wound healing satisfactorily, and no unfavourable symptoms showing themselves. He left the hospital quite recovered in the middle of December.

I have to thank Mr. Tatum for permission to publish this case.

The above case, with the case of White, which was published last week in this Journal, appear to me to possess many points of interest worthy of observation. On a careful perusal of their progress and treatment, it will be noticed that, notwithstanding their differences in age, the difference in the amount of the forces which occasioned the fractures, and the perceptible difference in the amount of concussion, the plan of treatment considered necessary for the one was also equally requisite for the other, although the conditions which indicated this, differed very materially in these two cases; and as the results in both were most favourable, their record will not have been the less interesting or less instructive.

The extensive amount of injury to the skull in the case of White, the difficulty in replacing into their natural

positions those portions of the fractured bones which were not removed, and the severe amount of shock which the brain itself sustained by the violence of the blow on the head, made this case appear very formidable in the first instance; nor was anxiety diminished whilst the disturbed condition of the brain was manifested by the constantly irritable and noisy state of the sufferer.

Compound fractures of the skull, with depression of bone, and the loss of a portion of the cranium, consequent upon either the comminution of the fractured pieces, or the use of the trephine, though always to be regarded with the most serious consideration, are generally found to be less grave in their results in the younger than in the older subjects that come under the care of the surgeon. The age of this boy greatly favoured the chances of recovery. But, in all probability, the greater sensitiveness of the growing brain of the child maintained, if it did not increase, the rather long continued irritability and restlessness of the patient.

This condition might have been partly influenced or modified by the loss of blood sustained at the time of the accident, which in one so young might produce serious effects for the time being. In this case, I believe it influenced the character of the symptoms to as great an extent as it modified the plan of treatment.

It is almost impossible to suppose that, with the amount of injury seen in this case, such a complete and long continued state of concussion depended merely on the shaking of the brain; and yet, in the absence of positive proof, we have no very definite rule by which we can decide how far such a state of functional derangement could be independent of organic lesion. Though he continued irritable, noisy, and in a half-insensible state, for some days, his pulse was never once in a condition indicating excess of action or increase of power; yet, in consequence of the heat of skin, and the very restless and noisy state, which rather increased some few hours after the operation, it was considered desirable to put him slightly under the influence of calomel. But the following day, as he appeared low, and as nothing but fluid had passed through his lips for three days, the calomel was stopped. The loss of blood, the starvation, and the action of the calomel, were all to be considered, separately and together; and it was evident there was no occasion to reduce him. The pulse also indicated weakness; and he was consequently ordered beef-tea. The next day he was still so weak that wine and water was ordered, in addition to the beef-tea; and yet the head symptoms were but little altered or improved. It was his general aspect and the pulse which indicated the importance of more stimulus.

With food, and with the cautious administration of wine, but with no other treatment as regards medicine, this boy recovered without a bad symptom.

He was seen some weeks after leaving the hospital, well in health, and with no discomfort in the head. The brain was no longer seen to pulsate, and the wound was sound. The eyeball was almost in its natural position, and the eye-sight good.

Though the second case differs materially from that of White in the amount of violence which produced the fracture, and also in the amount of cerebral disturbance, still it points out in an equal degree the care that is required in the treatment of symptoms arising in a case of compound fracture of the skull. After some days of progressive improvement, and without any previously active symptoms, pain in the head was complained of; and Mr. Tatum considered it desirable to place him under the action of calomel. At the end of twenty-four hours, the pain of the head was relieved, and the general symptoms indicated that calomel was no longer required. The pulse was weak, and the discharge from the wound rather foul. There were indications of constitutional disturbance threatening, of a character opposite to that of power; and he was consequently ordered beef-tea and wine.

The following day, erysipelas and œdema were observed

on the scalp, and his wine was increased. With the increase of the wine and food, but with no other medical treatment, he gradually improved, and left the hospital recovered from all the effects of the accident.

In the case of White, the use of stimulus was indicated at an early period after the accident; and this was apparently influenced to a great extent by the loss of blood in one so young. The want of stimulus was equally evident in the other case, but at a later period; and that from a condition more immediately resulting from the poison of erysipelas damaging the system—a poison, the effects of which in the wards of a London hospital, rarely admit of any other treatment than one of support and stimulus.

ORIGINAL COMMUNICATIONS.

CASES OF PROLAPSUS AND PROCIDENTIA UTERI.

By W. J. ANDERSON, F.R.C.S., Obstetric Surgeon to the St. George's and St. James' Dispensary.

IN the volume of the ASSOCIATION JOURNAL for 1854, at page 904, I made some remarks upon the treatment of procidentia uteri; and I am induced to revert to this subject again, as a new pessary was there mentioned somewhat diffidently, on account of my not having had sufficient experience as to its efficacy. A year has elapsed since that time, which has afforded me ample opportunities not only of witnessing its utility in the possibly bigoted hands of its inventor, but also in those of others. The instrument is intended to supply the place of the boxwood, or other ball pessaries; it is composed of an air ball of vulcanised India rubber, and serves all the purposes of the others, without any of their inconveniences; it is "of the same shape, able to give the same kind of support, but sufficiently elastic to avoid all pain when passing in and out of the vagina, and incapable of becoming saturated with the discharges, or of retaining them in its cavity; it is light, elastic, and firm; fills the vagina without producing injurious pressure, and is beautifully clean, from the fact of its being waterproof even to its string, and, therefore, when washed, is as fresh as new." In my former remarks I deprecated most strongly the use of the perinæal pad in procidentia, as well as the impropriety of using a pessary in cases of simple prolapsus, and upon these points I still hold the same opinion; the few subjoined cases will, however, show my views upon this subject quite as accurately as the most elaborate description could possibly do.

CASE I. Mrs. T., aged 30 years, February 1st, 1854. Procidentia uteri of three months standing; she is now about six weeks advanced in her fifth pregnancy; the uterus, anterior wall of the vagina, and bladder came down, but are easily replaced in the recumbent posture; there is profuse leucorrhœal discharge and dragging pain in the back and abdomen. She has suffered lately from anxiety and fatigue. Perfect rest was enjoined, and astringent injections, tonics, and mild aperients were employed. As pregnancy advanced, the uterus rose and became fixed above the pelvis, its size preventing its falling down, but a sensation of bearing down continued until she was delivered on 21st September. The child was healthy, and she had a good time; and up to January 1855, when I last saw her, there had been no bearing down.

As a general rule, where procidentia occurs in the early months of pregnancy, a pessary is more likely to do harm